

# ANIMAL PRODUCTION SCIENCE

**CONTENTS** Volume 53, Issue 1, 2013, 1–86

## PORK CRC

Foreword

*Roger Campbell* i

Methyl donor supplementation of gestating sow diets improves pregnancy outcomes and litter size

*W. H. E. J. van Wettere, R. J. Smits and P. E. Hughes* 1–7

Dietary ractopamine promotes growth, feed efficiency and carcass responses over a wide range of available lysine levels in finisher boars and gilts

*C. V. Rikard-Bell, J. R. Pluske, R. J. van Barneveld, B. P. Mullan, A. C. Edwards, N. J. Gannon, D. J. Henman and F. R. Dunshea* 8–17

Fibre in late gestation gilt diets, glucose tolerance, and feed intake and bodyweight loss during subsequent lactation

*P. Langendijk and T.-Y. Chen* 18–22

Increasing the energy content of lactation diets fed to first-litter sows reduces weight loss and improves productivity over two parities

*R. J. Smits, D. J. Henman and R. H. King* 23–29

Feeding level and dietary energy source have no effect on embryo survival in gilts, despite changes in systemic progesterone levels

*R. Z. Athorn, P. Stott, E. G. Bouwman, A. C. Edwards, M. A. Blackberry, G. B. Martin and P. Langendijk* 30–37

Current recommended levels of dietary lysine in finisher pig diets are sufficient to maximise the response to ractopamine over 28 days but are insufficient in the first 7 days

*C. V. Rikard-Bell, J. R. Pluske, R. J. van Barneveld, B. P. Mullan, A. C. Edwards, N. J. Gannon, D. J. Henman and F. R. Dunshea* 38–45

Piglet performance and immunity is determined by the parity of both the birth dam and the rearing dam

*Y. J. Miller, A. M. Collins, D. Emery, D. J. Begg, R. J. Smits and P. K. Holyoake* 46–51

Blend-feeding or feeding a single diet to pigs has no impact on growth performance or carcass quality

*K. L. Moore, B. P. Mullan and J. C. Kim* 52–56

Embryo survival, but not first-parity litter size, is increased when gilts are fed diets supplemented with omega-3 fatty acids from fish oil

*R. J. Smits, B. G. Luxford, M. Mitchell and M. B. Nottle* 57–66

The response of entire male and female pigs from 20 to 100-kg liveweight to dietary available lysine

*K. L. Moore, B. P. Mullan, R. G. Campbell and J. C. Kim* 67–74

Spray-dried porcine plasma and yeast derived protein meal influence the adaption to weaning of primiparous and multiparous sow progeny in different ways

*M. V. Edwards, R. G. Campbell, T. Chapman, H. Brouwers, S. G. Pierzynowski, B. R. Weström, O. Prykhod'ko, L. Gabor and M. Chocł* 75–86