

[10.1071/MF23161](https://doi.org/10.1071/MF23161)

Marine and Freshwater Research

Supplementary Material

Immune responses of sea cucumber (*Apostichopus japonicus*) to combined environmental stress from high temperature and oil pollution

Zhonglei Ju^A, Xishan Li^{A,}, Wei Yang^A, and Deqi Xiong^{A,*}*

^ACollege of Environmental Science and Engineering, Dalian Maritime University, Dalian, 116026, PR China.

*Correspondence to:

Xishan Li College of Environmental Science and Engineering, Dalian Maritime University, Dalian, 116026, PR China
Email: lxsdmu@dlnu.edu.cn

Deqi Xiong College of Environmental Science and Engineering, Dalian Maritime University, Dalian, 116026, PR China
Email: xiongdq@dlnu.edu.cn




Fig. S1. Concentrations of 16 priority PAHs (Σ PAHs) listed by the US Environmental Protection Agency (1984) in the water-accommodated fraction (WAF) of Oman crude oil exposure solution at an optimal temperature of 16°C (WAF + 16°C) and Oman crude oil WAF exposure solution at an elevated temperature of 26°C (WAF + 26°C).

Reference

US Environmental Protection Agency (1984) Method 610: polynuclear aromatic hydrocarbons. US EPA, Washington, DC, USA.