

Supplementary Material

Where there's smoke, there's cats: long-unburnt habitat is crucial to mitigating the impacts of cats on the Ngarlgumirdi, greater bilby (*Macrotis lagotis*)

Harry A. Moore^{A,B,}*

^ADepartment of Biodiversity, Conservation and Attractions, Bentley Delivery Centre, Locked Bag 104, Perth, WA, Australia.

^BSchool of Agriculture and Environmental Sciences, University of Western Australia, Crawley, WA 6009, Australia.

^CNyamba Buru Yawuru, Environmental Services Unit, 55 Reid Road, Cable Beach, WA 6726, Australia.

Lot 19–20, First Street, Ardyaloon, PO Box 2145, Broome, WA 6725, Australia.

C/o Beagle Bay Community, PO Box 2145, Broome, WA 6725, Australia.

Walalakoo Aboriginal Corporation, 47 Loch St, Derby, WA 6728, Australia.

School of Biological Sciences, University of Western Australia, Crawley, WA 6009, Australia.

Research Institute for Environment and Livelihoods, Charles Darwin University, Darwin, NT 0800, Australia.

*Correspondence to: Harry A. Moore Department of Biodiversity, Conservation and Attractions, Bentley Delivery Centre, Locked Bag 104, Perth, WA, Australia Email: harry.moore@dbca.wa.gov.au



Figure S1. Typical post fire successional stages of observed within acacia shrublands and eucalypt woodlands on the Dampier Peninsula, <3 years fire stages dominated by annual grasses.

Table S1. Model summary from generalised linear mixed-effects models testing the influence of fire attributes on the presence of greater bilbies and feral cats at the 1000-m scale on the Dampier peninsula in north-west Australia.

Variable	Bilby			Feral cat		
	Estimate	P	s.e.	Estimate	P	s.e.
Intercept	-4.15	0.00	0.94	-1.20	0.00	0.22
Fire Frequency	-0.72	0.07	0.40	0.55	0.01	0.20
Recent rainfall	0.03	0.92	0.30	0.19	0.30	0.18
Acacia shrublands	0.76	0.48	1.07	-0.07	0.89	0.54
Intercept	-5.04	0.02	2.23	-1.17	0.00	0.23
Pyrodiversity	0.03	0.95	0.48	-0.10	0.60	0.18
Recent rainfall	-0.03	0.93	0.34	0.21	0.25	0.18
Acacia shrublands	0.73	0.59	1.37	-0.20	0.72	0.56
Intercept	-4.83	0.04	2.34	-1.16	0.00	0.22
Proportion recently burnt	-0.07	0.88	0.46	0.21	0.20	0.17
Recent rainfall	-0.02	0.94	0.34	0.23	0.20	0.18
Acacia shrublands	0.73	0.58	1.31	-0.14	0.80	0.55
Intercept	-4.09	0.00	0.84	-1.21	0.00	0.22
Proportion long unburnt	0.87	0.02	0.37	-0.57	0.00	0.20
Recent rainfall	0.07	0.81	0.29	0.20	0.26	0.18
Acacia shrublands	0.59	0.57	1.03	0.00	0.99	0.55

Table S2. Model summary from generalised linear mixed-effects models testing the influence of fire attributes on the presence of greater bilbies and feral cats at the 3000-m scale on the Dampier peninsula in north-west Australia.

Variable	Bilby			Feral cat		
	Estimate	P	s.e.	Estimate	P	s.e.
Intercept	-3.89	0.00	0.77	-1.24	0.00	0.22
Fire Frequency	-0.81	0.02	0.35	0.41	0.04	0.20
Recent rainfall	0.06	0.84	0.28	0.17	0.34	0.18
Acacia shrublands	0.70	0.48	0.99	-0.12	0.84	0.59
Intercept	-4.15	0.00	1.02	-1.21	0.00	0.22
Pyrodiversity	-0.48	0.30	0.46	-0.08	0.66	0.19
Recent rainfall	-0.06	0.84	0.30	0.20	0.27	0.18
Acacia shrublands	1.05	0.35	1.13	-0.27	0.65	0.59
Intercept	-4.10	0.00	1.09	-1.21	0.00	0.22
Proportion recently burnt	-0.17	0.66	0.38	0.29	0.08	0.17
Recent rainfall	-0.04	0.89	0.30	0.22	0.22	0.18
Acacia shrublands	0.83	0.45	1.11	-0.16	0.78	0.58
Intercept	-3.85	0.00	0.73	-1.27	0.00	0.22
Proportion long unburnt	0.85	0.01	0.34	-0.63	0.00	0.21
Recent rainfall	0.06	0.82	0.27	0.18	0.34	0.18
Acacia shrublands	0.57	0.56	0.98	0.02	0.97	0.58

Table S3. Model summary from generalised linear mixed-effects models testing the influence of fire attributes on the presence of greater bilbies and feral cats at the 5000-m scale on the Dampier peninsula in north-west Australia.

Variable	Bilby			Feral cat		
	Estimate	P	s.e.	Estimate	P	s.e.
Intercept	-3.92	0.00	0.78	-1.23	0.00	0.22
Fire Frequency	-0.80	0.02	0.34	0.40	0.05	0.20
Recent rainfall	0.09	0.75	0.29	0.17	0.34	0.18
Acacia shrublands	0.72	0.48	1.00	-0.15	0.80	0.58
Intercept	-4.22	0.00	1.15	-1.23	0.00	0.22
Pyrodiversity	-0.19	0.67	0.45	-0.20	0.31	0.20
Recent rainfall	-0.05	0.86	0.30	0.18	0.31	0.18
Acacia shrublands	1.04	0.38	1.19	-0.13	0.83	0.61
Intercept	-4.11	0.00	1.13	-1.23	0.00	0.22
Proportion recently burnt	-0.14	0.71	0.38	0.38	0.02	0.17
Recent rainfall	-0.04	0.89	0.30	0.22	0.21	0.18
Acacia shrublands	0.84	0.45	1.12	-0.12	0.84	0.58
Intercept	-3.85	0.00	0.73	-1.30	0.00	0.23
Proportion long unburnt	0.88	0.01	0.34	-0.70	0.00	0.22
Recent rainfall	0.09	0.74	0.28	0.18	0.34	0.18
Acacia shrublands	0.54	0.58	0.98	0.10	0.86	0.59

Table S4. Model summary from generalised linear mixed-effects models testing the influence of fire attributes on the presence of greater bilbies and feral cats at the 10000-m scale on the Dampier peninsula in north-west Australia.

Variable	Bilby			Feral cat		
	Estimate	P	s.e.	Estimate	P	s.e.
Intercept	-4.16	0.00	1.00	-1.16	0.00	0.22
Fire Frequency	-0.65	0.09	0.39	0.19	0.33	0.19
Recent rainfall	0.10	0.75	0.31	0.20	0.26	0.18
Acacia shrublands	0.04	0.97	1.11	-0.42	0.41	0.52
Intercept	-4.47	0.00	1.33	-1.15	0.00	0.22
Pyrodiversity	-0.49	0.30	0.47	0.02	0.93	0.20
Recent rainfall	-0.08	0.79	0.31	0.24	0.19	0.18
Acacia shrublands	0.84	0.50	1.26	-0.54	0.34	0.56
Intercept	-4.61	0.11	2.89	-1.18	0.00	0.21
Proportion recently burnt	0.00	1.00	0.63	0.33	0.05	0.17
Recent rainfall	-0.07	0.85	0.33	0.24	0.17	0.18
Acacia shrublands	0.26	0.83	1.22	-0.28	0.59	0.51
Intercept	-3.79	0.00	0.72	-1.26	0.00	0.22
Proportion long unburnt	0.97	0.01	0.35	-0.64	0.00	0.22
Recent rainfall	0.17	0.56	0.28	0.18	0.32	0.18
Acacia shrublands	-0.28	0.77	0.96	0.04	0.94	0.53