# CS R Publishing



# Molluscan Research

Volume 22, 2002 © Malacological Society of Australasia

# All enquiries should be directed to:



Molluscan Research CSIRO Publishing PO Box 1139 (150 Oxford St) Collingwood, Vic. 3066, Australia

Telephone: +61 3 9662 7629 C S I RO PUBLISHING Telephone: +61 3 9662 7629 Fax: +61 3 9662 7611 Email: publishing.mr@csiro.au

Published by CSIRO Publishing for the Malacological Society of Australasia

www.publish.csiro.au/journals/mr

# Some Recent Thraciidae, Periplomatidae, Myochamidae, Cuspidariidae and Spheniopsidae (Anomalodesmata) from the New Zealand region and referral of *Thracia reinga* Crozier, 1966 and *Scintillona benthicola* Dell, 1956 to *Tellimya* Brown, 1827 (Montacutidae) (Mollusca : Bivalvia)

## Bruce A. Marshall

Museum of New Zealand Te Papa Tongarewa, PO Box 467, Wellington, New Zealand. Email: brucem@tepapa.govt.nz

#### Abstract

Thracia vitrea (Hutton, 1873) is redescribed and referred to Thracia Blainville, 1824, subgenus Odoncineta O. G. Costa, 1829. The circum-Antarctic species Thracia meridionalis E. A. Smith, 1885 is newly recorded from New Zealand and referred to subgenus Crassithracia Soot-Ryen, 1941. New thraciids are described belonging to Asthenothaerus Carpenter, 1864 (one), Parvithracia Finlay, 1926 (s. str.) (one), Parvithracia (Pseudoasthenothaerus) Kamanev, 2002 (two) and Trigonothracia Yamamoto & Habe, 1959 (one). Myadora biconvexa Powell, 1927 and Parvithracia cuneata Powell, 1937 are referred to, Barythaerus, a new genus of Thraciidae. Thracia reinga Crozier, 1966 and Scintillona benthicola Dell, 1956 are referred to Tellimya Brown, 1827 in Montacutidae. The periplomatid Pendaloma micans (Hedley, 1901) is newly recorded as widely distributed off New Zealand. New data on shell morphology and distributions are provided for the myochamids Hunkydora novozelandica (Reeve, 1859) and Myochama tasmanica (Tenison Woods, 1877) and new species of Hunkydora Fleming, 1948 and Myadoropsis Habe, 1960 are described. Cuspidaria aupouria Dell, 1950 is referred to Rhinoclama Dall & E. A. Smith in Dall, 1886 (s. str.). A new and minute species of the Cuspidariidae from the Norfolk Ridge is referred to a new genus, namely Pseudogrippina, and cuspidariids belonging in Plectodon Carpenter, 1864 (three) and Rhinoclama (Austroneaera) Powell, 1937 (two) are described. The three Rhinoclama (Austroneaera) species previously recorded from the New Zealand region are reviewed. Spheniopsidae is transferred from Heterodonta to Anomalodesmata and tentatively grouped in Cuspidarioidea. Six new spheniopsids of the genus Grippina Dall, 1912 are described and G. aupouria (Powell, 1937) is redescribed.

Additional keywords: Asthenothaerus, Austroneaera, Barythaerus, Crassithracia, Grippina, Hunkydora, Myadoropsis, Myochama, new taxa, Odoncineta, Plectodon, Pseudoasthenothaerus, Pseudogrippina, Rhinoclama, Trigonothracia.

# Introduction

The subclass Anomalodesmata is a large group of highly derived marine bivalves characterised by diverse and distinctive anatomies and shells. The gills are modified to form a unique muscular pump in the carnivorous species (Cuspidariidae, Poromyidae and, to some extent, Verticordiidae) or have a reduced and dorsally reflected outer demibranch in the deposit feeders and filter feeders (other families; Prezant 1998 and references therein). The hinge typically lacks teeth, although secondary teeth are present in some taxa. The primary ligament may be internal or external, supplemented in some taxa by a dorsal secondary ligament that helps maintain alignment of the valves. The inner primary ligament is characteristically elongated and calcified as a lithodesma, which is involved with load distribution in the hinge area and alignment and opening of the valves (Yonge and Morton 1980).

The present contribution was initiated during routine identification in preparation for databasing, when I recognised six undescribed *Grippina* species, which almost doubles the number of living Spheniopsidae known from the world. Besides describing these, the

© Malacological Society of Australasia 2002

10.1071/MR02011 1323-5818/02/030221

opportunity is taken to disentangle the three distinct species belonging to three genera in two families currently confused under *Hunkydora novozelandica* (Reeve, 1859) and to record additional related species.

# Materials and methods

Shells smaller than 10 mm in maximum dimension were selected, soaked in aqueous wetting agent (trisodium phosphate solution), briefly soaked in strong commercial bleach (to remove traces of soft tissue), washed and cleaned ultrasonically in distilled water and stored in 90% ethanol. They were then dried and affixed to aluminium stubs with conductive carbon tape, 'wired' with conductive carbon paint, sputter-coated with carbon and gold-palladium and examined and digitally imaged with a scanning electron microscope. Larger shells were cleaned and photographed uncoated under artificial light with a digital camera. Distribution maps were prepared by downloading personally verified coordinates for personally verified lots from the museum's database (Te Kahui) onto appropriate maps using a commercial mapping program and final plates were prepared using a commercial digital image manipulation program. Shells were measured with a calibrated optical graticule in a stereomicroscope. Shell terminology follows Cox (1969).

Institutional abbreviations and text conventions

AIM	Auckland Institute and Museum, Auckland
AMS	The Australia Museum, Sydney
BMNH	The Natural History Museum, London
MNHN	Muséum National d'Histoire Naturelle, Paris
NMNZ	Museum of New Zealand Te Papa Tongarewa, Wellington
NSMT	National Science Museum, Tokyo
NZGS	Institute of Geological and Nuclear Sciences, Lower Hutt
NZOI	National Institute of Water and Atmospheric Research, Wellington
PI	Prodissoconch I
PII	Prodissoconch II
pr	pair of valves of single individual
RRS	Royal Research Ship
RV	Research Vessel
SS	Sailing Ship

- v valve
- ZMB Zoological Museum, Humboldt University, Berlin

Unless specified, all material examined is at NMNZ (registration numbers preceded by 'M.').

In descriptions of ligamental structures, I have used the term 'sunken' for resilifers that have migrated (in an evolutionary sense) from the hinge line to a position behind/beneath the hinge line as viewed from the valve plane. A sunken resilifer (= subumbonal plate of Kamanev 2002) is associated with a secondarily enlarged ligament that may be calcified (= lithodesma of many Anomalodesmata and some Montacutidae).

#### Systematics

# Class **BIVALVIA** Linnaeus, 1758

# Subclass ANOMALODESMATA Dall, 1890

# Superfamily THRACIOIDEA Stoliczka, 1870

# Family THRACIIDAE Stoliczka, 1870

# Genus *Thracia* Blainville, 1824

*Thracia* Blainville, 1824: 347. Type species (by monotypy: two species group taxa originally included, one of which is a *nomen nudum*): *Mya pubescens* Pulteney, 1799; Recent, Europe. For discussion of authorship of *Thracia* and its type species, see Coan (1990*a*).

Osteodesma Blainville, 1827: 659. Type species (by subsequent designation of Dall, 1903: 1522): Anatina myalis Lamarck, 1818 = M. pubescens.

# Remarks

Three Recent species from the New Zealand region are referrable to *Thracia* (*s. lat.*): *Lyonsia vitrea* Hutton, 1873, here resurrected from synonymy and referred to subgenus *Odoncineta* O. G. Costa, 1829; *Thracia meridionalis* E. A. Smith, 1885, a new record for the region, here referred to *Crassithracia* Soot-Ryen, 1941; and *Thracia* (*s. lat.*) *vegrandis* P. Marshall & R. Murdoch, 1919 (not discussed herein: see Powell 1974: 200, 1979: 433).

*Thracia reinga* Crozier, 1966 from off the Three Kings Islands, northern New Zealand, is closely similar to *Tellimya ferruginosa* (Montagu, 1808), type species of *Tellimya* Brown, 1827 in gross shell morphology, although larger, and evidently belongs in Montacutidae. *Thracia reinga* is here referred to *Tellimya* together with the similar New Zealand species *Scintillona benthicola* Dell, 1956. Although strongly developed, the ligaments in *Montacuta substriata* (Montagu, 1808) and *M. tenella* (Lovén, 1846) (type species of *Montacuta* Turton, 1822 and *Decipula* Friele, 1875 respectively) are weaker than in *T. ferruginosa*, with resilifers less sunken and less oblique (Ockelmann 1965). *Tellimya ferruginosa* has a strongly developed, somewhat butterfly shaped, brown (non-calcified) ligament set on rather strongly oblique sunken resilifers (Fig. 1*L*). *Tellimya reinga* and *T. benthicola* both have large, strongly oblique, sunken resilifers, as does the similar North American species *Montacuta floridana* Dall, 1899 (type species of *Orobitella* Dall, 1900) and they must have substantial ligaments too. The enlarged ligament in the montacutid genus *Montacutona* Yamamoto & Habe, 1959 is partly calcified (i.e. a lithodesma; Morton 1980).

#### Subgenus Odoncineta O. G. Costa, 1829

*Odoncineta* Costa, 1829: 14. Type species (by monotypy): *Tellina papyracea* Poli, 1791 (not Gmelin, 1791) = *Amphidesma phaseolina* Lamarck, 1818; Recent, Europe.

#### Thracia (Odoncineta) vitrea (Hutton, 1873) n. comb.

# Figs 1A,D,H,K, 3A, 5A, 21A

Lyonsia vitrea Hutton, 1873a: 61; Hutton, 1878: 44; Hutton, 1880: 136.

- *Thracia vitrea.* Hutton, 1884: 514 (in part = '*Parilimya' neozelanica* (Suter, 1914)); Hutton, 1893: 75 (in part = *Asthenothaerus maxwelli* n. sp. (pl. 9, fig. 80) + '*P' neozelanica*); Suter, 1913: 1024 (in part: pl. 62, fig. 15 = *A. maxwelli*); Dell, 1956: 41; Powell, 1957: 84 (in part: pl. 11, fig. 24 = *A. maxwelli*); Powell, 1962: 125 (in part: pl. 11, fig. 24 = *A. maxwelli*); Marshall, 1998: 145.
- *Eximiothracia vitrea*. Powell, 1937*b*: 61 (in part: pl.11, fig. 24 = *A. maxwelli*); Powell, 1946: 63 (in part: pl. 11, fig. 24 = *A. maxwelli*).
- *Thracia (Hunkydora) australica novozelandica.* Powell, 1976: 131 (in part not Reeve, 1859: pl. 18, fig. 24 = *A. maxwelli*); Powell, 1979: 433 (in part not Reeve: pl. 79, figs 6, 7 = *A. maxwelli*).

NOT Thracia vitrea. – Odhner, 1924: 84 (= A. maxwelli).

#### Material examined

*Syntypes.* No locality data; type locality here selected as Wellington Harbour, New Zealand (4 fragmentary v, M.281).

*Other material examined.* SW of Cape Maria van Diemen,  $34^{\circ}41.9'S$ ,  $172^{\circ}33.5'E$ , alive, 103 m (1pr, M.76043). Bay of Islands:  $35^{\circ}11.5'S$ ,  $174^{\circ}15.5'E$ , 54-58 m (1v, M.95832); off Onewhero Bay,  $35^{\circ}14'S$ ,  $174^{\circ}06'E$ , 18 m (1v, M.35568); off Onewhero Bay,  $35^{\circ}14'S$ ,  $174^{\circ}05.5'E$ , 15 m (1v, M.44218); off Onewhero Bay,  $35^{\circ}14'S$ ,  $174^{\circ}06'E$ , 18 m (1v, M.35568); off Onewhero Bay,  $35^{\circ}14'S$ ,  $174^{\circ}05.5'E$ , 15 m (1v, M.44218); off Onewhero Bay,  $35^{\circ}14'S$ ,  $174^{\circ}06'E$ , alive, 18 m (1pr, 1v, M.49478). Off Hen Island,  $36^{\circ}00.5'S$ ,  $174^{\circ}43'E$ , alive, 59 m (1pr, M.43819). W of Plate Island,  $37^{\circ}39.1'S$ ,  $176^{\circ}31.5'E$ , 64-59 m (1pr, 1v, M.67588). Off Motiti Island, alive, 46 m (7pr, M.39370). SE of Plate Island,  $37^{\circ}43.4'S$ ,  $176^{\circ}38.5'E$ , alive, 59 m (3pr, M.65313). Off Motuhora Island, 25 m (2pr, 3v, M.39929). ENE of Tolaga Bay,  $38^{\circ}15.2'S$ ,  $178^{\circ}38.6'E$ , 139 m (1v,

M.59820). Wellington Harbour, various stations (8pr, M.8000; 1pr, M.15940; 6v, M.15939; 8v, M.15941; 3pr, M.62798; 17pr, M.144103; 8pr, M.144123; 25pr, M.144136; 3pr, M.144160; 1v, M.14401). NE of Stephens Island, 40°31'S, 174°15'E, alive, 117–119 m (1pr, M.53756); 40°33'S, 174°07'E, alive, 132 m (1pr, M.53033). W of Kapiti Island, 40°44'S, 174°34'E, 146 m (1v, M.11306). Marlborough Sounds: Catherine Cove, Admiralty Bay, 40°53'S, 173°54'E, 40-44 m (2pr, M.55110); Admiralty Bay, 40°54'S, 173°56'E, alive, 44 m (2pr, 1v, M.55049); between West Entry Point and Duffers Reef, Pelorus Sound mouth, 40°57.5'S, 174°01.5'E, alive, 29 m (8pr, M.51506); Orchard Bay, Pelorus Sound mouth, 40°57.5'S, 174°03.5'E, alive, 26–29 m (3pr, 1v, M.54699); Orchard Bay, 40°58'S, 174°04'E, alive, 29 m (2pr, 1v, M.45133); Waitui Bay, 41°00.5'S, 174°11.0'E, alive, 15 m (2pr, 1v, M.51549); Port Gore entrance, 41°00.5'S, 174°15.0'E, alive, 15 m (7pr, M.51630); between Motuara Island and White Rocks, Queen Charlotte Sound, 41°05.5'S, 174°19.0'E, 15 m (8pr, 1v, M.54326); Titirangi Bay, Guards Bay, 41°01'S, 174°08'E, 4-9 m (1pr, M.149603); head of Anakoha Bay, Guards Bay, 41°02'S, 174°06'E, alive, 15-18 m (2pr, M.52177); Manaroa Bay, Pelorus Sound, alive, 5-18 m (10pr, M.144402); Otanerau Bay in East Bay, Arapawa Island, 41°11'S, 174°20'E, alive, 20–29 m (18pr, M.50830); Otanerau Bay in East Bay, 41°11'S, 174°20'E, 22–37 m (1v, M.54383); Double Bay, Kenepuru Sound (3v, M.18069). Akaroa Harbour entrance, 43°54'S, 172°57'E, alive, 13–15 m (1pr, M.108881). Stewart Island: Maori Beach (1v, M.3460); off Vaila Voe (2v, M.19293); Glory Inlet, 18-36 m (3v, M.19295).

#### Description

Shell up to 30 mm long, thin and brittle, whitish; left valve almost as inflated as the right, dorsoventrally shorter, and seated largely within its perimeter; umbones low, rounded, at approximately posterior one-third; periostracum very thin, translucent pale buff. Prodissoconch approximately 200 µm wide, strongly inflated (surface eroded). Dissoconch margins smoothly merging; anterior margin well rounded, posterior margin subtruncate, other margins broadly rounded; exterior posterior area distinctly flattened, elsewhere rather evenly rounded. Hinge line interrupted by narrow U-shaped space occupied by small lithodesma. Ligament strong, projecting externally, seated on large, prominent, rounded, similar chondrophores that descend from hinge line. Lithodesma butterfly shaped, small (width 0.8 mm in shell 13.7 mm long; 0.9 mm in shell 21.5 mm long), development slowing as shell enlarges, its resilifers elevated, oblique, steadily enlarging together with chondrophores as part of their supporting buttresses. Pallial sinus U-shaped, deep, depth 48%–51% of shell length. Anterior adductor scar considerably larger than posterior. Interior traversed by weak radial lines. Interior iridescent due to thin layer of nacre prisms. Exterior with low, irregular commarginal growth lines and wrinkles; flattened posterior area covered with minute, crowded granules, elsewhere a dulling microsculpture resembling irregular paving stones.

# Distribution

North, South and Stewart islands, 0-146 m; taken alive at 15-132 m from mud (locally common; Fig. 5*A*).

# Remarks

Practically since its introduction, the New Zealand species *Lyonsia vitrea* Hutton, 1873, has been variously confused with the parilimyid '*Parilimya*' *neozelanica* (Suter, 1914) and the superficially similar species *Asthenothaerus maxwelli* n. sp. (Thraciidae) and *Hunkydora novozelandica* (Reeve, 1859) (Myochamidae). The type material of *Lyonsia vitrea* (NMNZ) originally comprised four valves gummed to a glass plate. These valves are now fragmented, but the species involved is clearly indicated by valve outline in the gum on the plate, shell thinness, iridescence and microsculpture and the characteristic chondrophores. *Lyonsia vitrea* is here tentatively referred to *Thracia*, subgenus *Odoncineta*, because of close similarity to the European type species, *T.* (*O.*) *phaseolina* (Lamarck, 1818) in gross



**Fig. 1.** Shells of *Thracia, Asthenothaerus* and *Hunkydora* species and details of hinge areas. *A*,*D*,*H*,*K*, *Thracia* (*Odoncineta*) vitrea (Hutton, 1873): *A*,*H*, right (*A* upper) and left valve, length 18 mm, off Kau Point, Wellington Harbour (M.62798); *D*, dorsal view of pair, length 20 mm, between Somes Island and Eastbourne, Wellington Harbour (M.144136); *K*, ventral view of hinge area, lithodesma arrowed. *B*,*G*, *Thracia* (*Crassithracia*) meridionalis E. A. Smith, 1885: right valve (*B* upper), length 22.0 mm, and left valve, length 26.0 mm, Papanui Canyon, off Otago Peninsula, New Zealand, 490–540 m (M.158244). *C*,*E*,*I*,*J*, *Asthenothaerus maxwelli* n. sp.: *C*,*I*, right (*C* upper) and left valve, length 30.5 mm, Horseshoe Bay, Stewart Island (holotype, M.152678); *J*, juvenile, length 2.50 mm, off Patea, 40 m (M.53531). *F*, *Hunkydora novozelandica* (Reeve, 1859): dorsal view of pair, length 15 mm, SE of Three Kings Islands, 121 m (M.150896). *L*, *Tellimya ferruginosa* (Montagu, 1808): ventral view of hinge area, showing large, non-calcified ligament (arrow), Eldfisk Oil Field, North Sea, 70 m (Swedish Museum of Natural History, Stockholm). *M*, *Pendaloma micans* (Hedley, 1901): ventral view of hinge area, lithodesma arrowed, Pegasus Canyon, NE of Banks Peninsula, 622 m (M.90220). Scale bars: *K*,*M*, 1 mm; *L*, 200 µm.

shell morphology. However, the New Zealand species differs markedly in being thinner and in that the interior is slightly, but distinctly, iridescent due the presence of a thin nacreous layer (thickness approximately 30 µm; Fig. 3A). Although nacre has not been recorded from Thraciidae hitherto (Keen 1969b: 850; Taylor et al. 1973: 282), interior iridescence (presumably due to nacre) has, nevertheless, been recorded in T. (s. lat.) myopsis Beck in Møller, 1842 (Greenland) (Soot-Ryen 1941), T. (Odoncineta) roumei Cosel, 1995 (West Africa) (confirmed as interior by R. von Cosel, personal communication), and T. (Eximiothracia) speciosa Angas, 1869 (Australia) (Coan 1990a). The latter species is type of Eximiothracia Iredale, 1924, which Coan (1990a) has suggested is possibly synonymous with Odoncineta, despite indication that T. (O.) phaseolina is not iridescent. Whether or not T. (O.) phaseolina and other thraciids that are non-irridescent actually lack a thin nacreous layer (perhaps sandwiched within the shell wall) remains to be established by examination of shell cross-sections. Uncertain of relationships in this group, I follow Coan (1990a) in treating Odoncineta as a subgenus of Thracia, at rank equal with Ixartia Leach, 1852, Homoeodesma P. Fischer, 1887, Crassithracia Soot-Ryen, 1941 and Cetothrax Iredale, 1949.

*Thracia* (*Odoncineta*) *vitrea* is illustrated here for the first time (Figs 1*A*,*D*,*H*,*K*, 3*A*, 21*A*); all previously published illustrations purporting to represent it are actually of *Asthenothaerus maxwelli* n. sp. *Thracia* (*O*.) *vitrea* is separable from the previously named species *Hunkydora novozelandica* by the thinner shell, more posterior umbones, more strongly convex left valve, much deeper pallial sinus, relatively smaller lithodesma, exterior microsculpture of microscopic granules rather than crisp, commarginally elongate prisms and in having prominent matching chondrophores hanging from the hinge line of both valves.

# Subgenus Crassithracia Soot-Ryen, 1941

*Crassithracia* Soot-Ryen, 1941: 19. Type species (by original designation): *Thracia crassa* Becher, 1886 = *T. septentrionalis* Jeffreys, 1872 (following Coan *et al.* 2000); Recent, North Atlantic. Introduced as a subgenus of *Thracia* Blainville, 1824.

# Thracia (Crassithracia) meridionalis E. A. Smith, 1885 n. comb.

# Figs 1B,G, 5C

Thracia meridionalis Smith, 1885: 68, pl. 6, figs 4–4b; Dell, 1990: 63, figs 109–111 (early references); Hain, 1990: 101, pl. 15, figs 7a,b; Branch et al. 1991: 51; Linse, 1997: 61; Linse & Brandt, 1998: 884; Cattaneo-Vietti et al. 2000: 176; Troncoso et al. 2001: 112, fig. 42.
Mysella? truncata Thiele, 1912: 230, pl. 18, fig. 18.
Mysella? frigida Thiele, 1912: 230, pl. 18, fig. 19.

#### Material examined

*Syntypes of Thracia meridionalis.* Balfour Bay, Royal Sound, Kerguelen Is. (BMNH 1887.2.9.2473, 1887.2.9.2473a-b and 2474, 2475).

Holotype of Mysella truncata. Gauss Station, Davis Sea, Antarctica (1v, ZMB 63110).
Holotype of Mysella frigida. Gauss Station, Davis Sea, Antarctica (ZMB 63111).
Other material examined. Antarctica: several hundred specimens (34 lots NMNZ); New Zealand:
Papanui Canyon, off Otago Peninsula, 45°50′S, 171°01′E, 490–540 m (5v, M.158244).

#### Distribution

Circum-Antarctic and off Otago Peninsula, New Zealand, 5–752 m; taken alive at 5–220 m (Fig. 5*C*).

# Remarks

The Papanui Canyon valves fall within the range of variation of Antarctic specimens of this common, widely distributed species and represent both a new record for the New Zealand region and the northern-most record for the species. No other specimens have been recorded from scores of rich samples dredged from Papanui Canyon or other submarine canyons off Otago Peninsula, suggesting that the shells possibly date from a period when sea temperatures were lower. The sample containing them was extremely and unusually rich, both in number of species and individuals, although almost entirely empty shells, and seems likely to have accumulated over a long period of time, probably hundreds or even thousands of years (but certainly post-dating last glacial maximum).

*Thracia meridionalis* is here referred to subgenus *Crassithracia* because of its similarity to *T. septentrionalis* in shell morphology. It is even closer to the circum-Arctic species *T. devexa* G. O. Sars, 1878, which also seems likely to belong here. The ligament in *T. septentrionalis* and *T. devexa* is much stronger than in *T. vitrea*, with resilifer rims that are long and upturned rather than short and downturned, the ligament in *T. vitrea* projecting further ventrally.

#### Genus Asthenothaerus Carpenter, 1864

Asthenothaerus Carpenter, 1864: 311. Type species (by monotypy): Asthenothaerus villosior Carpenter, 1864; Recent, eastern Pacific.

# Asthenothaerus maxwelli n. sp.

#### Figs 1*C*,*E*,*I*,*J*, 3*B*, 4*A*, 5*B*, 21*B*

*Thracia vitrea.* – Hutton, 1893: 75 (in part not Hutton, 1873 + '*Parilimya*' *neozelanica* (Suter, 1914)), pl. 9, fig. 80; Suter, 1913: 1024 (in part not Hutton), pl. 62, fig. 15; Odhner, 1924: 84 (not Hutton).

*Eximiothracia vitrea*. – Powell, 1937*b*: 61 (in part not Hutton), pl. 02, fig. 13, 64 milet, 1924. 61 (in part not Hutton), pl. 11, fig. 24; Powell, 1946: 63 (in part not Hutton), pl. 11, fig. 24

*Thracia vitrea*. – Dell, 1956: 41 (in part not Hutton); Powell, 1957: 84 (in part not Hutton), pl. 11, fig. 24; Powell, 1962: 125 (in part not Hutton), pl. 11, fig. 24.

*Thracia (Hunkydora) australica novozelandica.* – Powell, 1974: 200, figs 13, 14 (not Reeve, 1859); Powell, 1976:131 (in part + *T. vitrea*), pl. 18, fig. 24; Powell, 1979: 433 (in part + *T. vitrea*), pl. 79 figs 6, 7 only.

*Parvithracia cuneata*. – Powell, 1979: 434 (Auckland Islands record). Thraciidae sp. 1 Spencer *et al.* (in press).

#### Material examined

Holotype. Horseshoe Bay, Stewart Island, New Zealand, Apr. 1963, E. C. Smith (pr, M.152678).

*Paratypes.* New Zealand, Stewart Island, Horseshoe Bay, Apr. 1963, E. C. Smith (5pr, 5v, M.16872); Horseshoe Bay (3pr, M.15302; 2pr, 10v, M.30985; 12pr, M.132839).

Other material examined. Fossil: Landguard Bluff, Wanganui, Late Pleistocene (Castlecliffian) (1pr, 8v, M.13080). Recent: Off Rangaunu Bay,  $34^{\circ}49.6'$ S,  $173^{\circ}15.0'$ E, alive, 23 m (1pr, M.150895). Off Takau Bay,  $35^{\circ}10'$ S,  $174^{\circ}11'$ E, 80 m (2v, M.43695). Bay of Islands: off Twin Rocks,  $35^{\circ}10'$ S,  $174^{\circ}18'$ E, 46-73 m (1v, M.154732); off island at entrance to Deep Water Cove,  $35^{\circ}11.9'$ S,  $174^{\circ}17.1'$ E, 47-49 m (2v, M.152674); between Hope Passage and Deepwater Cove,  $35^{\circ}12'$ S,  $174^{\circ}18'$ E, 37-40 m (2v, M.41416); Bamboo Bay, Moturua Island,  $35^{\circ}13.9'$ S,  $174^{\circ}11.3'$ E, 4-6 m (1v, M.40840). Schooner Bay, Great Barrier Island, 15 m (1pr, M.155466); off Cuvier Island, 49-73 m (1v, M.3458). Off Rurima Rocks, 26 m (2, M.152676). Ranfurly Bank, East Cape,  $37^{\circ}32.8'$ S,  $178^{\circ}48.7'$ E, 94 m (1pr, M.60728);  $37^{\circ}33.1'$ S,  $178^{\circ}49.5'$ E, 94-89 m (3v, M.64820). N of Motuhora Island,  $37^{\circ}35.9'$ S,  $176^{\circ}59.5'$ E, 179-139 m (1v, M.60572). N of New Plymouth,  $38^{\circ}55.0'$ S,  $174^{\circ}09.3'$ E, 48 m (1pr, M.150894). S of Patea, alive,  $39^{\circ}56'$ S,

174°26'E, 40 m (11pr, M.53531). SW of Wanganui, 40°11'S, 174°49'E, 58-64 m (2pr, M.50776). Off D'Urville Island: off Puangiangi Island, 15-22 m (1pr, M.100530); Long Beach (3v, M.11936). Titirangi Bay, Guards Bay, 41°01'S, 174°08'E, 4-9 m (5v, M.54071). Head of Titirangi Bay, beach (18v, M.40656). Queen Charlotte Sound, 15 m (1pr, M.132954). Manaroa Bay, Pelorus Sound, 22 m (4pr, 5v, M.18354). Mernoo Bank, W Chatham Rise, 43°06.1'S, 175°20.5'E, 153 m (1v, M.65072); 42°51.9'S, 175°26.5'E, 183 m (1v, M.15437); 42°59.4'S, 175°30.5'E, 112 m (2v, M.15438). Chatham Islands: N of The Sisters, 43°32.5'S, 176°47.5'W, 60 m (1v, M.12410); E of Forty Fours, 44°04'S, 175°23.5'W, 238 m (1v, M.10872). Head of Waitaki Canyon, off Oamaru, 45°10'S, 171°30'E, 293-256 m (1v, M.51265). SW of Hare's Ears, Doubtful Sound, 45°17.2S, 166°49.3'E, 146 m (4v, M.58743). Off Taiaroa Head: 45°44'S, 171°02'E, 137 m (2v, M.11083); Taiaroa Canyon, 45°45.4'S, 171°05'E, 549 m (2v, M.152675); NE of Cape Saunders, 45°50'S, 170°56'E, 105 m (3v, M.45373); Papanui Canyon, 45°51'S, 171°00'E, 348–220 m (1v, M.74868); 45°51'S, 170°59'E, 220 m (1v, M.131304). Off NW end of Cooper Island, Dusky Sound, 45°44.0'S, 166°47.5'E, 12 m (1v, M.135529). Opposite Sandy Point, Long Sound, 46°02.16'S, 166°45.88'E, alive, 20 m (1pr M.141156). Off Taieri, 46°07.0'S, 170°39.0'E, 87 m (8v, M.58904). Stewart Island: Easy Harbour, 47°09'S, 167°34'E, 7-20 m (1v, M.26561); North Arm, Port Pegasus, 47°11'S, 167°41'E, 37-44 m (2v, M.26725); North Arm, 47°11'S, 167°41'E, 40-46 m (1v, M.42401); S of Port Pegasus, 47°18.5'S, 167°42.5'E, 119 m (1v, M.26514); Mason Bay (1pr, M.19297); Halfmoon Bay (5v, M.19296); off Vaila Voe (3pr, M.155168); Maori Beach (2v, M.3457); Golden Bay (1pr, M.19294). Off Bounty Islands, 49°40.19'S, 178°44.30'E, alive, 113 m (1pr, 8v, M.150075). Off Antipodes Islands, 49°40'S, 178°52'E, 86-95 m (3v, M.39558); 49°40'S, 178°53'E, 103 m (5v, M.39558). N of Auckland Islands, 174 m (2v, AIM 131367). Between Dees Head and Tucker Point, Auckland Islands, 26-27 m (4v, M.16744).

#### Description

Shell up to 46 mm long, longer than high, thin, white, hinge edentulous, no external ligament, lithodesma well developed; periostracum very thin, olive green. Prodissoconch 200 µm wide, PI globular, 150-170 µm wide. Right valve approximately 1% larger than left, approximately 8% thicker, left valve sitting within its perimeter; beaks at approximately posterior one-third, opisthogyrous. Posterodorsal margin straight, anterior margin rounded, anterodorsal and ventral margins more broadly rounded, margins smoothly merging; posterior margin obliquely truncate, separated by rounded angulations. Externally convex, right valve more inflated than left; posterior area defined by rounded angulation, weakly concave or flat; sculpture of weak, irregular, rounded commarginal ridges; microsculpture of minute, crisp, crowded granules resembling irregular paving stones (not separated by wavy radial striae), coarser on posterior area. Lithodesma very large relative to those in most other thraciids, strong, somewhat butterfly shaped; resilifers oblique, narrowly subtriangular, posterior margin straight or broadly concave; anterior margin broadly convex, elevated, rim undercut. Anterior adductor scar curved, narrowly elliptical; posterior scar subcircular; pallial sinus depth 44%-48% of adult shell length, anterior margin rounded.

# Distribution

Late Pleistocene (Castlecliffian) – Recent. North, South, Stewart, Auckland, Antipodes, Bounty and Chatham Islands and Chatham Rise, 0–293 m; taken alive (juveniles) at 20–40 m from sand and sandy mud (Fig. 5*B*).

# Remarks

*Asthenothaerus maxwelli* closely resembles the type species of *Asthenothaerus* (Coan 1990*a*: 39, figs 32, 33) in gross facies, including the characteristic lack of an external ligament. However, the New Zealand species differs markedly in attaining more than three times the size. Specimens from off the Auckland and Antipodes Islands (length up to 46 mm) attain larger size than specimens from elsewhere in the New Zealand region, but

are otherwise indistinguishable. As indicated by the synonymy, *A. maxwelli* has been confused with both *Thracia vitrea* and *Hunkydora novozelandica*, the three species being sympatric, but apparently asyntopic. Although quite similar to *T. vitrea* in shell shape, *A. maxwelli* differs markedly in that the ligament is fully internal, in lacking chondrophores and in that the lithodesma is very much larger and fully functional at all stages of growth (development retarded from an early stage of growth in *T. vitrea*). *Asthenothaerus maxwelli* differs further in attaining a larger size (length up to 46 mm as against 30 mm) and in that the shell is thicker and considerably stronger. *Asthenothaerus maxwelli* differs principally from *H. novozelandica* in having a more strongly convex left valve with beaks further posterior, in lacking a posterodorsal groove on the right valve for seating the corresponding margin of the left valve and in that the external microscopic granules are thinner, mostly longer and separated by wavy radial striae.

Hutton (1893) treated *Thracia granulosa* Hutton, 1873 (1873*b*) as a synonym of *T. vitrea* and illustrated a specimen of what is clearly *A. maxwelli*, which is not uncommon at Landguard Bluff, Wanganui. When providing the substitute name *T. neozelanica* for *T. granulosa* Hutton, 1873 (not A. Adams & Reeve, 1850), Suter (1914) correctly indicated that *T. neozelanica* is an entirely different species, as indeed is clear from Hutton's original description. *Thracia neozelanica*, which also occurs Recent (16 lots NMNZ), belongs in the Parilimyidae and is currently grouped in *Parilimya* Melvill & Standen, 1899 for want of a more appropriate location (Beu and Maxwell 1990).

Asthenothaerus maxwelli is similar to the eastern Australian myochamid Hunkydora australica (Reeve, 1859) in shape (two syntypes BMNH 1962695, Moreton Bay, Queensland; the larger figured by Fleming 1951, figs 1, 2), but the latter is immediately separable by the characteristic microsculpture (external shell layer) of fine commarginal striae and wavy radial striae.

# Etymology

After Philip Maxwell of Waimate (formerly NZGS), in appreciation of his friendship and outstanding contributions to molluscan systematics and palaeontology.

# Genus Barythaerus n. gen.

Type species: Myadora biconvexa Powell, 1927; Recent, southern New Zealand.

# Diagnosis

Shell small (maximum lengths 5.10 and 5.50 mm), stout, beaks at posterior one-third to one-quarter, no external ligament, resilifers sunken, lithodesma ovate, prominent tooth in front of beak on right valve only, no corresponding socket in left valve. Interior not nacreous, pallial sinus depth 33%–41% of shell length. Externally with irregular, paving stone-like microsculpture.

# Remarks

The two *Barythaerus* species resemble *Parvithracia* species in having a prominent anterior tooth on the right valve, but differ in being stouter and more oblique and in having an irregular paving stone-like exterior dissoconch microsculpture and markedly opisthogyrous beaks. *Barythaerus* and *Parvithracia* species differ further in the shape of the lithodesma,

which is more or less ovate in *Barythaerus*, rather than butterfly shaped as in *Parvithracia*. *Asthenothaerus* species are similar in shape but lack the prominent tooth in front of the beak on the right valve. *Parvithracia* Finlay, 1926 and *Lampeia* MacGinitie, 1959 are less stout, have considerably shorter anterior ends and a butterfly-shaped lithodesma. The resilifers are simple and not buttressed as in *Lampeia* (Kamanev and Nadtochy 1998). A few valves from off the Chatham Islands and East Otago (Thraciidae sp. 4 of Spencer *et al.* in press; e.g. M.10860, M.48859) represent an additional species. See additional remarks under *Grippina* (below).

#### Etymology

From the Greek barys (heavy) and thaeros (hinge); gender masculine.

#### Barythaerus biconvexus (Powell, 1927)

# Figs 2A,C, 3C, 4B, 5D, 21C

Myadora biconvexa Powell, 1927: 123, pl. 23, fig. 9; Powell, 1979: 432.

#### Material examined

*Syntypes.* Off Puysegur Point, New Zealand, 46°11′S, 166°30′E, 183 m, Dec. 1908, SS *Rakiura* (12v, CM M.861, M.862)

*Other material examined.* Off Taieri,  $46^{\circ}07.0'$ S,  $170^{\circ}39.0'$ E, 87 m (1v, M.66069);  $46^{\circ}15.0'$ S,  $170^{\circ}29.0'$ E, 91 m (3v, M.66168). Off Puysegur Point,  $46^{\circ}11'$ S,  $166^{\circ}30'$ E, alive, 183 m (and 311 m – Powell 1927) (1pr, many v, M.7407). SE of Nugget Point,  $46^{\circ}40'$ S,  $170^{\circ}00'$ E, 140 m (2v, M.65941). Stewart Island: off Tehatatahi Island, 46-64 m (1v, M.19877); off Halfmoon Bay, 27-55 m (2v, M.19873); Halfmoon Bay, 27 m (1v, M.19874); off Garden Point, 22 m (2v, M.19875); off Poutama Island,  $47^{\circ}16'$ S,  $167^{\circ}23'$ E, alive, 55 m (1pr, 15v, M.19878); off Big Island, South Cape, 73-82 m (2v, M.19876).

# Description

Shell up to 5.10 mm long, strongly inaequilateral, oblique, umbones at approximately posterior one-sixth, weakly to very strongly thickened, moderately inflated,  $1.48-1.57\times$  longer than high, translucent white. Prodissoconch approximately 230 µm wide, smooth; PI approximately 170 µm wide, globular. Dissoconch with shallow but distinct lunule, anterodorsal margin more or less flat or very weakly convex, posterodorsal margin broadly concave, anterior and posterior margins well rounded, ventral margin very broadly rounded or more or less flat; exterior broadly convex. Right valve with strong anterior hinge tooth, strongly shelved anteriorly and posteriorly to seat elevated antero- and posterodorsal margins of left valve; hinge area of both valves divided by broad triangular space. Lithodesma stout, short, thick; supporting plates simple, oblique, elliptical. Interior glossy, with weak radial lines; pallial sinus and pallial line well defined, sinus depth 33%–34% of shell length. Adductor and anterior pedal retractor muscle scars sharply defined. Exterior with low, rounded, anterodorsal angulations that bound concave lunule; tightly rounded to concave escutchion; microsculpture resembling irregular paving stones, imparting a dull sheen.

# Distribution

Southern South Island and Stewart Island, 27–311 m; taken alive at 55–183 m from bryozoan/shell substrata (Fig. 5*D*).



**Fig. 2.** Shells of *Barythaerus* and *Parvithracia* species. *A,C, Barythaerus biconvexus* (Powell, 1927): left (*A* upper) and right valve, off Poutama Island, Big South Cape Island, Stewart Island, 55 m (M.19878). *B,D, Barythaerus cuneatus* (Powell, 1937): left valve (*B* upper), length 5.30 mm, and two right valves, lengths 4.60 mm (*B* lower) and 5.00 mm (*D*), 22 km ENE of Great Island, Three Kings Islands, 200 m (M.144363). *E,G, Parvithracia* (*s. str.*) *suteri* (Finlay, 1927): left (*G* upper) and right valves, length 3.25 mm, North Arm, Port Pegasus, Stewart Island, 37–44 m (M.44790). *F,H, Parvithracia* (*s. str.*) *ampla* n. sp.: holotype, left (*H* upper) and right valve, length 6.60 mm, Saunders Canyon, off Otago Peninsula, 457 m (M.9125).

# Remarks

*Barythaerus biconvexus* is redescribed here to facilitate direct comparison with *B. cuneatus. Barythaerus biconvexus* attains more than twice the size of its type specimens (length up to 5.1 mm).

# Barythaerus cuneatus (Powell, 1937)

Figs 2B,D, 3D, 4C, 5E, 21D

*Parvithracia cuneata* Powell, 1937*a*: 174, pl. 46, fig. 9; Powell, 1979: 434, fig. 117/4 (in part: Auckland Islands record = *Asthenothaerus maxwelli*).

#### Material examined

*Holotype.* Off Three Kings Islands, New Zealand, 34°11.6'S, 172°10.9'E, 92 m, 17 Aug. 1932, *RRS Discovery II* (1v, BMNH 1962949).

*Other material examined.* Off Three Kings Islands: King Bank, 33°57.0'S, 172°19.0'E, 128 m (8v, M.149501); King Bank, 33°57.4'S, 172°19.4'E, alive, 128–123 m (1pr, 34v, M.149568); 37 km NE of Great Island, 33°58.0'S, 172°30.6'E, 550 m (1v, M.149505); 22 km ENE of Great Island, 34°05.0'S, 172°24.6'E, 200 m (2v, M.59551); Three Kings Trough, 34°00'S, 171°55'E, 805 m (6v, M.17519); 34°01'S, 172°07'E, 622 m (2v, M.35000); 13 km N of Great Island, 34°01.8'S, 172°12.0'E, 508 m (36v, M.59548); Middlesex Bank, 34°02.0'S, 171°44.0'E, 246–291 m (3v, M.149566); Middlesex Bank, 34°02.1'S, 171°45.8'E, 221–206 m (3v, M.149503); 22 km ENE of Great Island, 34°05.0'S, 172°24.6'E, 200 m (3v, M.144363); off North-east Island, 34°08.5'S, 172°11'E, 102 m (3v, M.34549); 34°11'S, 172°10'E, 91 m (1v, M.144364). NW of Cape Reinga, 34°14.2'S, 172°32.4'E, 100 m (6v, M.59552). Off Spirits Bay, 34°18.36'S, 172°49.39'E, alive, 68 m (1pr, many v, M.149504). N of Cape Reinga, 34°21.60'S, 172°43.19'E, 48 m (2v, M.153096). NW of Cape Reinga, 34°22.8'S, 172°24.6'E, 121 m (1v, M.149567).

# Description

Shell up to 5.5 mm long, strongly inequilateral, oblique, umbones at approximately posterior one-fifth, modestly thickened, moderately inflated,  $1.41-1.50 \times$  longer than high, translucent white. Prodissoconch approximately 250 µm wide, smooth; PI approximately 170 µm wide, globular. Dissoconch with shallow but distinct lunule, anterodorsal margin very broadly convex, posterodorsal margin shallowly concave, anterior margin well rounded, posterior and ventral margins broadly rounded, exterior broadly convex. Right valve with strong anterior hinge tooth, shelved anteriorly and posteriorly to seat elevated antero- and posterodorsal margins of left valve; hinge area of both valves divided by broad triangular space. Lithodesma stout, broad; supporting plates simple, oblique, elliptical. Interior glossy, with weak radial lines; pallial sinus and pallial line well defined, sinus depth 38%–41% of shell length. Adductor and anterior pedal retractor muscle scars sharply defined. Exterior with low, rounded, anterodorsal angulations that bound concave lunule; tightly rounded to concave escutchion; microsculpture resembling irregular paving stones, imparting a dull sheen.

#### Distribution

Off Three Kings Islands and off Spirits Bay, 68-805 m; taken alive at 68-128 m from comminuted bryozoan and shell substrata (Fig. 5*E*).

# Remarks

*Barythaerus cuneatus* is extremely similar to *B. biconvexus* in gross facies, differing principally in having a lighter hinge area, a deeper pallial sinus and in the shape of the lithodesma; the great majority of specimens differ further in that the hinge tooth on the right valve is shorter, yet projects further ventrally, the shell is higher relative to the length and the shell wall is thinner. The two species are allopatric.



Fig. 3. Dissoconch shell structure and microsculpture of Thraciidae, Myochamidae and Cuspidariidae (exterior, right valve and adult unless indicated). A, Thracia (Odoncineta) vitrea (Hutton, 1873), inner surface of central dissoconch, obliquely fractured to show nacre prisms; Guards Bay, Marlborough, 15-18 m (M.52177). B, Asthenothaerus maxwelli n. sp., juvenile, off Patea, 40 m (M.53531). C, Barythaerus biconvexus (Powell, 1927), off Poutama Island, Big South Cape Island, Stewart Island, 55 m (M.19878). D, Barythaerus cuneatus (Powell, 1937) (M.144363). E, Trigonthracia mimica n. sp., holotype, Doubtful Sound entrance, 117-128 m (M.155019). F, Myadoropsis wairua n. sp., holotype, N of Three Kings Islands, 622 m (M.152686). G, Myochama tasmanica (Tenison Woods, 1877), left valve, costate area of pre-cementation stage, Arch Pinnacle, Prince's Islands, Three Kings Islands, 40 m (M.117130). H,I, Pseudogrippina wanganellica n. sp., holotype (I = interior), Wanganella Bank summit, Norfolk Ridge, 113 m (M.152681). J,K, Plectodon lepidus n. sp., edge of prodissoconch (arrow) and early dissoconch (J) and adult (K), off E side of Mayor Island, 59-74 m (M.152682). L, Plectodon pruinosus n. sp., holotype, Middlesex Bank, NW of Three Kings Islands, 98-103 m (M.152683). M, Plectodon regalis n. sp., holotype, off Three Kings Islands, 91 m (M.152684). N, Rhinoclama (Austroneaera) brevirostris (Powell, 1937), SE of Great Island, 173-178 m (M.144426). O, Rhinoclama (Austroneaera) raoulensis (Powell, 1958), off Meyer Island, Raoul Island, Kermadec Islands, 27-22 m (M.225814). Scale bar: A, 20 µm; others, 50 µm.

# Genus Parvithracia Finlay, 1926

Parvithracia Finlay, 1926: 461. Type species (by original designation): Montacuta triquetra Suter, 1913, not Verrill & Bush, 1898 (preoccupied) = Parvithracia suteri Finlay, 1927; Recent, New Zealand.

# Remarks

Kamanev (2002) has recently introduced *Pseudoasthenothaerus* as a new subgenus of *Parvithracia* for species with a lunule that lacks an internal groove beside the posterodorsal margin and that have resilifers supported by columnar buttresses. Whereas *P. ampla* n. sp. has characteristics of *Parvithracia* (*s. str.*), *P. melchior* n. sp. and *P. fragilissima* n. sp. resemble *Pseudoasthenothaerus* species in lacking an internal groove beside the posterodorsal margin, but the resilifer plate is not supported by pillars and there is no lunule.

# Parvithracia (s. str.) suteri Finlay, 1927

# Figs 2*E*,*G*, 4*E*, 7*A*, 21*E*

*Montacuta triquetra* Suter, 1913: 915, pl. 53, fig. 7*a* (not Verrill & Bush, 1898 (preoccupied)). *Parvithracia triquetra*. – Finlay, 1926: 461.

Parvithracia suteri Finlay, 1927: 529 (replacement name for Montacuta triquetra Suter, 1913); Dell, 1956: 167 (in part = P. ampla n. sp.); Powell, 1979: 434, fig. 117/3.

Parvithracia (Parvithracia) suteri. - Kamanev, 2002: 109, figs 2-10.

#### Material examined

Lectotype (Boreham 1959: 21). Port Pegasus, Stewart Island, New Zealand, 33 m (NZGS TM 399). Other material examined. SE of Great Island, Three Kings Islands, 34°14.8'S, 172°13.6'E, 173–178 m (8v, M.149537). N of North Cape: 34°18'S, 173°02'E, 137 m (1v, M.3060); SE of Three Kings Islands, 34°20.2'S, 172°21.8'E, 121 m (7v, M.149532); 34°20.0'S, 173°06.6'E, 163-168 m (1v, M.149545). N of Kerr Point, Tom Bowling Bay, 34°22.01'S, 173°00.20'E, 89 m (1v, M.155120). NW of Cape Reinga, 34°22.8'S, 172°24.6'E, 121 m (4v, M.149542). E of North Cape, 34°23.0'S, 173°12.6'E, 447–357 m (1v, M.149538). E of North Cape, 34°25.0'S, 173°13.1'E, 327–257 m (30v, M.149536). Great Exhibition Bay: 34°33.4'S, 173°04.8'E, 63 m (2v, M.113191); 34°33.6'S, 173°04.9'E, 66 m (1v, M.149540). SW of Cape Maria van Diemen, 34°41.9'S, 172°33.5'E, 103 m (1v, M.149541). Off Rangaunu Bay, 34°42.8'S, 173°14.5'E, 63 m (10v, M.155117). Off Whangaroa Bay, 34°54.0'S, 173°42.6'E, alive, 83 m (2pr, 30v, M.155118). Off Cape Brett, 35°08'S, 174°12.5'E, alive, 80 m (2pr, 30v, M.35716). Off Twin Rocks, 35°10′S, 174°18′E, 46–73 m (21v, M.42182). Off Takau Bay, 35°10′S, 174°11′E, 80 m (13v, M.43650). Off Ahipara, 35°10.4'S, 172°35.4'E, 233 m (2v, M.149546). Bay of Islands: 35°11.2'S, 174°16.7'E, alive, 58-64 m (15pr, 6v, M.96116); 35°11.5'S, 174°15.5'E, 54-58 m (1v, M.131108); 35°11.6'S, 174°16.5'E, 54 m (2v, M.95923); Deepwater Cove entrance, 35°12'S, 174°18'E, 46 m (3v, M.41697). Off Poor Knights Islands: 35°22'S, 174°43'E, 146 m (30v, M.35141); 35°28'S, 175°03'E, 282 m (1v, M.36066); 35°29'S, 174°43.5′E, 110 m (4v, M.74370); 35°32′S, 174°41′E, 121-113 m (18v, M.35829). Off Hen Island: 35°58.5'S, 174°44'E, 62 m (3pr, 7v, M.43937); 36°00.5'S, 174°43'E, 59 m (5pr, 8v, M.43821). Off Aldermen Islands: 37°00.8'S, 176°12.3'E, alive, 178-248 m (10pr, 15v, M.66634); 165 m (1v, M.44387). Off Mayor Island: 37°11.5'S, 176°10.0'E, alive, 198–273 m (2pr, M.66600); 37°16.7'S, 176°17.5'E, 104-109 m (1pr, 2v, M.67389); 37°18.9'S, 176°16.2'E, 59-74 m (1v, M.65729). Rungapapa Knoll, 37°33.8'S, 176°59.0'E, W of White Island, 188-228 m (1v, M.113535). Off Plate Island, 37°39.4'S, 176°34.4′E, 82 m (1v, M.67744). Ranfurly Bank, East Cape: 37°32.8′S, 178°48.7′E, 94 m (many v, M.60799); 37°33.1'S, 178°49.5'E, alive, 94-89 m (2pr, 30v, M.71422); 37°33.4'S, 178°48.3'E, alive, 106-103 m (1pr, 40v, M.71654); 37°36.7'S, 178°51.6'E, 56-63 m (2v, M.149539); 37°37.8'S, 178°52.4'E, 50-72 m (1v, M.96195); 37°38.5'S, 178°56.4'E, 153-143 m (21v, M.149534). Off Raglan Harbour: 37°48.0'S, 174°14.7'E, 103 m (1v, M.149535); 37°48.0'S, 174°14.0'E, 103 m (3v, M.149533). N of New Plymouth, 38°40.2'S, 174°03.9'E, alive, 88 m (4pr, 3v, M.149544). Off Cape Kidnappers, 39°52.8'S, 177°36.5'E, alive, 785-882 m (3pr, 2v, M.155115). S of Waverley, 40°10'S, 174°40'E, 77-82 m (4v, M.50645). W of Rangitikei R. mouth, 40°16'S, 174°58.5'E, alive, 75 m (4pr, 1v, M.53286). NE of Stephens Island, 40°33'S, 174°07'E, 132 m (1pr, M.53071). N of Kapiti Island, 40°38'S, 174°54.5'E, alive, 91 m (2v,

2pr, M.52902). E of Stephens Island, 40°42'S, 174°10.6'E, 106 m (2pr, M.7771). Off Palliser Bay, 41°35'S, 175°04'E, 128–146 m (1pr, 2v, M.44307); SE of Cape Campbell, 42°00.8'S, 174°41.0'E, alive, 939–1019 m (3pr, M.60459). Pegasus Canvon, 43°31.0'S, 173°30.5'E, alive, 256-293 m (1pr, M.94783). Milford Sound: 44°36.0'S, 167°49.0'E, 123 m (6v, M.10359); entrance sill, 44°36.42'S, 167°50.47'E, 117 m (6v, M.11423). Head of Waitaki Canyon, off Oamaru, 45°10'S, 171°30'E, 293-256 m (2v, M.51322). Thompson Basin floor, Thompson Sound, 45°14.38'S, 166°58.87'E, 340-362 m (3, M.150518). Doubtful Sound entrance, 45°16', 166°48', alive, 117-128 m (13pr, M.58625). Bradshaw Basin floor, Bradshaw Sound, 45°17.3'S, 167°02.6'E, 415 m (1pr, 4v, M.138387). Doubtful Sound: Utah Basin floor, 45°17.9'S, 166°55.5'E, 400 m (2v, M.138681); Seymour Basin, Malaspina Reach, 45°18.15'S, 167°00.64'E, 107 m (2pr, M.147072); Seymour Basin, 45°18.90'S, 167°00.64'E, 94 m (1v, M.147099); Seymour Basin, 45°18.21'S, 167°00.50'E, 64 m (1v, M.147128); Seymour Basin, 45°18.21'S, 167°00.87'E, alive, 94 m (6pr, 5v, M.147169); Seymour Basin, 45°18.27'S, 167°00.69'E, 115 m (4v, M.147231). Off Chatham Point, upper Breaksea Sound, 45°33'S, 166°53'E, 42 m (1v, M.139727). Off Tairoa Head: Karitane Canyon, 45°37.5'S, 171°03'E, 420 m (2v, M.45618); head of Karitane Canyon, 45°38.5'S, 171°05.0'E, alive, 585-530 m (4pr, 4v, M.51091). Cook Channel, Dusky Sound, 45°46.0'S, 166°43.0'E, 44 m (1v, M.135465). North Arm, Port Pegasus, Stewart Island, 47°11'S, 167°41'E, alive, 37-44 m (many pr, M.44790).

#### Distribution

Three Kings, North, South, Stewart and Chatham Islands, 33-1019 m; taken alive at 37-549 m from mud (Fig. 7*A*).

#### Remarks

*Parvithracia suteri* is distinctive in its small size (length rarely greater than 4 mm), subtrigonal outline, subcentral beaks, obliquely subtruncate posterior end and strong anterior hinge tooth on the right valve. The prodissoconch (Fig. 4E) is smooth, approximately 180 µm wide and the PI is approximately 170 µm wide and globular.

# Parvithracia (s. str.) ampla n. sp.

Figs 2*F*,*H*, 4*F*, 7*B*, 21*F* 

Parvithracia suteri. - Dell, 1956: 167 (in part).

#### Material examined

*Holotype.* Saunders Canyon, off Otago Peninsula, New Zealand, 45°55.2'S, 170°56.4'E, alive, 457 m, 16 Jan. 1954, MV *Alert* (pr, M.9125).

*Paratypes.* Off Gable End Foreland, 38°38.7'S, 178°41.1'E, alive, 755–725 m (1pr, M.155114). Off Cape Kidnappers, 39°52.8'S, 177°36.5'E, alive, 785–882 m (4pr, M.155115). SE of Cape Campbell, 42°00.8'S, 174°41.0'E, alive, 939–1019 m (3pr, M.60459). Wall of Pegasus Canyon, NE of Banks Peninsula, 43°14'S, 173°39'E, alive, 1006–512 m (2pr, 2v, M.52890). Chatham Rise: 43°32'S, 178°38'E, 549 m (1v, M.10866); 43°38'S, 177°19'E, 531 m (2v, M.10865); E of Forty Fours, 44°04'S, 175°23.5'E, alive, 238 m (1pr, M.42286). Head of Waitaki Canyon, off Oamaru, 45°10'S, 171°30'E, alive, 293–256 m (8v, M.155152). Off Taiaroa Head: Karitane Canyon, 45°37.5'S, 171°03'E, 420 m (1v, M.155151); head of Karitane Canyon, 45°38.5'S, 171°05.0'E, 585–530 m (1v, M.155153); Taiaroa Canyon, 45°45.4'S, 171°05'E, 549 m (2v, M.9063); Papanui Canyon, 45°46'S, 171°03'E, 660 m (1v, M.58603); Saunders Canyon, 45°55'S, 170°56'E, 421 m (1v, M.26296).

*Other material examined.* Off Three Kings Islands: 22 km ENE of Great Island, 34°05.0'S, 172°24.6'E, 200 m (2v, M.149547); 34°10'S, 172°12'E, 252 m (1v, M.34235); SE of Great Island, 34°14.8'S, 172°13.6'E, alive, 173–178 m (2pr, 6v, M.155170); 37 km SW of Great Island, 34°20.4'S, 171°48.2'E, 440 m (1v, M.155121).

#### Description

Shell up to 6.50 mm long, longer than high, slightly but distinctly oblique, thin, white, no external ligament, left valve slightly smaller than right valve and seated within much of the



Fig. 4. Prodissoconchs of Thraciidae, Myochamidae and Cuspidariidae (prodissoconch I and II boundaries or prodissoconch boundary indicated by arrows). A, Asthenothaerus maxwelli n. sp., off Patea, 40 m (M.53531); B, Barythaerus biconvexus (Powell, 1927), off Puysegur Point, 183 m (M.7407). C, Barythaerus cuneatus (Powell, 1937), off Spirits Bay, 68 m (M.149504). D, Parvithracia (Pseudoasthenothaerus) melchior n. sp., 22 km ENE of Great Island, 200 m (M.59557). E, Parvithracia (s. str.) suteri (Finlay, 1927), North Arm, Port Pegasus, Stewart Island, 37-44 m (M.44790). F, Parvithracia (s. str.) ampla n. sp., head of Waitaki Canyon, off Oamaru, 293-256 m (M.155152); G, Trigonothracia mimica n. sp., holotype, Doubtful Sound entrance, 117-128 m (M.155019). H-J, Myochama tasmanica (Tenison Woods, 1877): H, Bradshaw Basin floor, Bradshaw Sound, Fiordland, 415 m (M.138438); I, SE of Three Kings Islands, 121 m (M.152688); J, Arch Pinnacle, Prince's Islands, Three Kings Islands, 40 m (M.117130). K, Hunkydora novozelandica (Reeve, 1859), North Arm, Port Pegasus, Stewart Island, 37-44 m (M.44788). L, Hunkydora rakiura n. sp., holotype, channel between Ulva Island and Bradshaw Peninsula, Paterson Inlet, Stewart Island, 33 m (M.152724). M, Pseudogrippina wanganellica n. sp., Wanganella Bank summit, Norfolk Ridge, 113 m (M.152681). N, Plectodon lepidus n. sp., holotype, off E side of Mayor Island, 59–74 m (M.152682). O, Plectodon pruinosus n. sp., Middlesex Bank, NW of Three Kings Islands, 98-103 m (M.152683). Scale bars: 50 µm.

perimeter of the right valve. Prodissoconch approximately 200 µm wide, smooth; PI approximately 180 µm wide, sharply defined, more strongly inflated than PII. Beaks roundly angulate, posterior occupying approximately 40% of shell length. Posterodorsal margin straight; posterior margin broadly rounded, oblique, subtruncate, bounded above and below by tightly rounded angulations; anterodorsal margin weakly convex, anterior margin well rounded; ventral margin broadly rounded. Inflation of both valves moderate, externally convex, posterior area weakly flattened, posterodorsal margins of both valves externally bounded by sharp angulations. Right valve with prominent hinge tooth immediately below and before prodissoconch, no corresponding socket in left valve; longer, low posterior tooth separated from dorsal margin by shallow groove; long, welldefined internal shelf beside anterodorsal and posterodorsal margins that seats left valve. Resilifers oblique, narrowly ovate, anterior rim weakly elevated and undercut, lithodesma present. Adductor muscle scars and pallial line well defined; pallial sinus depth 42%-43% of adult shell length, anterior margin rounded. Exterior with irregular commarginal ridges and growth lines, dull due to exceedingly fine, granular surface texture; each valve with flattened anterodorsal and posterodorsal areas, posterior area bounded by long, low, rounded angulation, anterior area boundary rounded.

#### Distribution

Off Three Kings Islands, south-eastern North Island, eastern South Island, and Chatham Rise, 173–1019 m; taken alive at 238–1019 m from foraminiferal ooze and mud (Fig. 7*B*).

# Remarks

Of the three *Parvithracia* species here recorded from the New Zealand region, *P. ampla* most closely resembles *P. suteri*, from which it differs in attaining larger size (length up to 6.50 v. 4.10 mm) and in having a longer anterior end. *Parvithracia suteri* and *P. ampla* are sympatric but asyntopic, *P. suteri* ranging into much shallower depths than *P. ampla*. I am unable to detect any differences between specimens from disjunct northern and southern populations and consider them to be conspecific.

# Etymology

From the Latin amplus (large).

# Subgenus Pseudoasthenothaerus Kamanev, 2002

Pseudoasthenothaerus Kamanev, 2002: 111. Type species (by original designation): Parvithracia (Psedoasthenothaerus) lukini Kamanev, 2002; Recent, NW Pacific.

#### Parvithracia (Pseudoasthenothaerus) melchior n. sp.

Figs 4D, 5F, 6A,C

Thraciidae sp. 2 Spencer et al. (in press).

# Material examined

*Holotype.* SE of Great Island, Three Kings Islands, New Zealand, 34°14.8'S, 172°13.6'E, 173–178 m, 2 Feb. 1981, RV *Tangaroa* (1v, M.155110).

*Paratypes.* Off Three Kings Islands: Middlesex Bank, 33°57.0'S, 171°45.4'E, 98–103 m (1v, M.149495); 37 km NE of Great Island, 33°58.0'S, 172°30.6'E, 550 m (2v, M.59550); Middlesex Bank, 33°59.9'S, 171°45.3'E, 186–196 m (3v, M.149496); Three Kings Trough, 34°00'S, 171°55'E, 805 m (2v,



Fig. 5. Maps of the New Zealand region showing distributions of *Thracia* and *Barythaerus* species (200 and 1000 m isobaths indicated). *A*, *Thracia* (*Odoncineta*) vitrea (Hutton, 1873). *B*, *Asthenothaerus* maxwelli n. sp. C, *Thracia* (*Crassithracia*) meridionalis E. A. Smith, 1885. *D*, *Barythaerus biconvexus* (Powell, 1927). *E*, *Barythaerus cuneatus* (Powell, 1937). *F*, *Parvithracia* (*Pseudoasthenothaerus*) melchior n. sp.

M.17518); 34°01'S, 172°07'E, 622 m (2v, M.149493); Middlesex Bank, 34°01.2'S, 171°44.4'E, 206–211m (2v, M.149494); 13 km N of Great Island, 34°01.8'S, 172°12.0'E, 508 m (1v, M.59547); Middlesex Bank, 34°02.0'S, 171°44.0'E, 246–291 m (2v, M.149499); 37 km NW of Great Island, 34°02.0'S, 171°48.4'E, 188 m (2v, M.59549); Middlesex Bank, 34°02.1'S, 171°45.8'E, 221–206 m (1v, M.149500); 22 km ENE of Great Island, 34°05.0'S, 172°24.6'E, 200 m (8v, M.59557); S of Great Island, 34°14.1'S, 172°09.0'E, 192–202 m (4v, M.149492); SE of Great Island, 34°14.8'S, 172°13.6'E, 173–178 m, 2 Feb. 1981, RV *Tangaroa* (6v, M.144395); 39 km SW of Great Island, 34°17.6'S, 171°45.3'E, 427 m (3v, M.149497); 34°18.8'S, 172°18.5'E, 93–88 m (1v, M.149498); 37 km SW of Great Island, 34°20.4'S, 171°48.2'E, 440

m (3v, M.59554); 28 km S of Great Island,  $34^{\circ}24.0'$ S,  $172^{\circ}16.8'$ E, 120 m (3v, M.59553). NW of Cape Reinga,  $34^{\circ}14.2'$ S,  $172^{\circ}32.4'$ E, 100 m (4v, M.144365).

*Other material examined.* 29°33'S, 168°07'E, SSE of Philip Island, Norfolk Island, 143 m (5v, M.224815); 29°20'S, 168°09'E, E of Norfolk Island, 201 m (4v, M.224736).

#### Description

Shell up to 11.5 mm long, longer than high, slightly oblique, of moderate thickness, white, no external ligament, lithodesma not seen. Prodissoconch approximately 230 µm wide, smooth; globular PI approximately 200 µm wide. Beaks at posterior one-third to onequarter, opisthogyrous. Posterodorsal margin straight or very weakly concave, roundly angulated to broadly rounded, subtruncate posterior margin; anterodorsal margin flat or weakly convex, anterior margin well rounded, ventral margin broadly rounded, margins smoothly merging. Inflation of both valves moderate, externally convex, posterior area flattened. Right valve with prominent hinge tooth immediately below and before prodissoconch, no corresponding socket in left valve; no trace of posterior hinge tooth and no posterodorsal groove. Resilifers oblique, narrowly ovate, anterior rim elevated and undercut. Interior glossy, not nacreous, sculptured with shallow radial grooves; adductor muscle scars and pallial line well defined; pallial sinus depth 39%-45% of adult shell length, anterior margin rounded. Exterior with well-defined posterodorsal angulations that bound concave escutchion, no lunule; irregular commarginal ridges and growth lines; dulled by microsculpture of extremely minute, crisp, crowded granules resembling irregular paving stones (not separated by wavy radial striae).

# Distribution

Off Norfolk Island and off Three Kings Islands, 88–805 m (shells only; Fig. 5F).

#### Remarks

*Parvithracia melchior* is similar to *P. ampla* in gross facies, but differs in attaining yet larger size, in having a longer anterior end and in having a more prominent tooth on the right valve. It is tentatively referred to subgenus *Pseudoasthenothaerus* because of similarity to north-west Pacific species in shape and lack of an internal groove beside the posterodorsal margin, but it differs in that the resilifer plates are not supported by pillars.

Specimens from off Norfolk Island and the Three Kings Islands have extremely similar shell morphology and appear to be conspecific.

# Etymology

After Melchior, one of the Three Kings.

# Parvithracia (Pseudoasthenothaerus) fragilissima n. sp.

Figs 6B,D, 7E, 21G

# Material examined

Holotype. Challenger Plateau, New Zealand, 40°50.1'S, 168°14.8'E, alive, 1005–1009 m, 18 Apr. 1980, RV Tangaroa (pr, NZOI H–808).

*Paratype.* Challenger Plateau, 40°50.1'S, 168°14.8'E, alive, 1005–1009 m, 18 Apr. 1980, RV *Tangaroa* (M.158220). Off Kahurangi Point, 41°00.6'S, 169°06.0'E, alive, 914 m (2pr, M.155144; 2pr, 1v, NZOI P942). Off Westport, 41°08.1'S, 170°21.0'E, alive, 695 m (1pr, NZOI Q699). Bounty Trough, 45°21.2'S, 173°35.8'E, alive, 1386 m (1pr, NZOI S153). Saunders Canyon, off Otago Peninsula, 45°55'S, 170°56'E, 421 m (1v, M.155154).



**Fig. 6.** Shells of *Parvithracia*, *Trigonothracia* and *Pendaloma* species. *A,C*, *Parvithracia* (*Pseudoasthenothaerus*) *melchior* n. sp.: holotype right valve (*C*, *A* lower), length 8.40 mm (M.155110) and paratype left valve (*A* upper), length 8.00 mm (M.144395), off Three Kings Islands, 200 m. *B,D*, *Parvithracia* (*Pseudoasthenothaerus*) *fragilissima* n. sp.: holotype, left (*B* upper) and right valve, length 6.70 mm, Challenger Plateau, 1005–1009 m (NZOI H–808). *E,G*, *Trigonothracia mimica* n. sp.: holotype, left (*G* upper) and right valve (*G* lower), length 2.05 mm (M.155019); paratype (*E*), length 2.35 mm (M.144393), Doubtful Sound entrance, 117–128 m. *F,H*, *Pendaloma micans* (Hedley, 1901): left (*H* upper) and right valve, length 18.0 mm, Pegasus Canyon, NE of Banks Peninsula, 622 m (M.90220).

# Description

Shell up to 7.90 mm long, longer than high, very thin and fragile, white, no external ligament, rather strongly inflated, left valve slightly smaller than right valve and seated within much of the perimeter of the right valve. Prodissoconch approximately 200 µm wide, smooth, rounded, no obvious PII (development presumably non-planktotrophic). Beaks roundly angulate, posterior occupying 43%-46% of adult shell length. Posterodorsal margin straight, tightly rounded to long, very broadly convex, slightly oblique, subtruncate, posterior margin, smoothly rounded to ventral margin; anterodorsal and ventral margins broadly convex, anterior margin well rounded. Externally posterior area shallowly concave, posterodorsal margins of both valves externally bounded by weak angulations. Right valve with weak hinge tooth immediately below and before prodissoconch, no corresponding socket in left valve; posterior tooth very weak or scarcely defined at all, separated from dorsal margin by shallow groove; narrow but well-defined internal shelf beside anterodorsal and anterodorsal margins that seats left valve. Resilifers oblique, narrowly ovate, anterior rim elevated and undercut, lithodesma present. Adductor muscle scars and pallial line well defined; pallial sinus depth 34%–38% of adult shell length, anterior margin rounded. Exterior with irregular commarginal ridges and commarginal and radial growth lines, dull but without defined microsculpture; each valve with very weak rounded angulation close to posterodorsal margin, anterodorsal area evenly rounded.

#### Distribution

Off north-western and south-eastern South Island, 421–1386 m; taken alive at 695–1386 m from foraminiferal ooze (Fig. 7*E*).

# Remarks

*Parvithracia fragilissima* differs from other *Parvithracia* species in the New Zealand region by its thinner, more inflated shell, higher posterior truncation and, relatively, much weaker hinge teeth. It is placed in subgenus *Pseudoasthenothaerus* because of lack of a groove beside the posterodorsal margin, but it differs from north-western Pacific *Pseudoasthenothaerus* species in lacking any trace of either lunule or resilifer plate pillars.

#### Etymology

From the Latin *fragilis* (fragile) and *issimus* (adjectival superlative; very fragile).

#### Genus Trigonothracia Yamamoto & Habe, 1959

Trigonothracia Yamamoto & Habe, 1959: 117. Type species (by original designation): Thracia (Trigonothracia) nomurai Yamamoto & Habe, 1959; Recent, Japan.

# Trigonothracia mimica n. sp.

Figs 3E, 4G, 6E, G, 7C

# Material examined

*Holotype*. Doubtful Sound entrance, New Zealand, 45°16'S, 166°48'E, alive, 117–128 m, 1 Jan. 1977, RV *Acheron* (pr, M.155019).

*Paratypes.* Doubtful Sound. Off Rangaunu Bay, 34°42.8'S, 173°14.5'E, alive, 63 m (1pr, M.155123). Off Whangaroa Bay, 34°54.0'S, 173°42.6'E, alive, 83 m (1pr, 3v, M.155122). Off Cape Brett, 35°08'S, 174°12.5'E, 80 m (2v, M.67869). Off Takau Bay, 35°10'S, 174°11'E, 80 m (5v, M.152723). W of Poor

Knights Islands, 35°32′S, 174°41′E, 121–113 m (1v, M.67871). Off Otarawairere Bay, Ohope, alive, 15 m (1pr, 2v, M.44086). N of New Plymouth, 38°40.2′S, 174°03.9′E, 88 m (2v, M.144392). NE of Kaingaroa, Chatham Islands, 43°35′S, 176°03.5′W, 220–229 m (2v, M.10863). Thompson Basin floor, Thompson Sound, 45°14.38′S, 166°58.87′E, 340–362 m (1v, M.150486). Bradshaw Basin floor, Bradshaw Sound, 45°17.3′S, 167°02.6′E, 415 m (2v, M.144394; 1v, M.138705). Seymour Basin, Malaspina Reach, Doubtful Sound: entrance, 45°16′S, 166°48′E, alive, 117–128 m, 1 Jan. 1977, RV *Acheron* (2, M.144393); 45°18.15′S, 167°00.64′E, 107 m (2v, M.147085); Seymour Basin, 45°18.27′S, 167°00.69′E, 115 m (3v, M.147248); 45°18.15′S, 166°43.5′E, 22–27 m (1v, M.14709); Cook Channel, 45°46.0′S, 166°43.0′E, 44 m (2v, M.144391).

#### Description

Shell small (length up to 3.95 mm), slightly longer than high, thin, white; periostracum extremely thin, pale cream. Prodissoconch approximately 200 µm wide, subcircular, smooth, PI approximately 180 µm wide. Right valve approximately 1% larger than left, both valves similar in thickness. Posterior length approximately 40% of shell length, beaks slightly opisthogyrous. Posterodorsal margin shallowly concave, anterodorsal margin straight, anterior margin rounded, ventral margin more broadly rounded, posterior margin strongly and slightly obliquely truncate. Prominent, slightly curved chondrophores behind well-developed lithodesma, ligament weakly developed; small, bluntly rounded tooth on right valve before lithodesma that sits against inner side of left valve (no corresponding socket). Lithodesma attachment area well defined, small, subcircular, directly below umbo. Interior glossy, not nacreous, anterior adductor scar curved, narrowly elliptical; posterior scar subtrigonal; pallial sinus depth approximately 40% of adult shell length, anterior margin rounded. Externally broadly convex; posterior area well defined, flattened; sculpture of low, irregular, rounded commarginal ridges; minute, crowded granules, considerably coarser and sharper on posterior area.

# Distribution

Off NE North Island, New Plymouth, Fiordland, and Chatham Islands, 15-415 m; taken alive at 15-128 m from mud (Fig. 7*C*).

# Remarks

*Trigonothracia mimica* is extremely distinctive among thraciids from the New Zealand region in its small size, subtrigonal outline, external sculpture and the conspicuous chondrophores. Although similar to the Japanese type species of *Trigonothracia* (holotype NSMT 39880; Yamamoto and Habe 1959: pl. 12, figs 9, 10; Higo *et al.* 2001: 182) in gross facies it differs in a number of details, including deeper pallial sinus, weaker inflation, more tightly rounded anterior end and a more strongly truncate posterior end, and in attaining approximately one-third the size (length 3.95 *v.* 10.10 mm). *Trigonothracia mimica* is remarkably similar to *Parvithracia suteri* in shape and size, but the two locally sympatric species differ markedly in hinge morphology and external sculpture.

# Etymology

From the Greek *mimicus* (imitative), alluding to its resemblance to *Parvithracia suteri* in shape and size.



Fig. 7. Maps of the New Zealand region showing distributions of *Parvithracia*, *Trigonothracia*, *Pendaloma* and *Myadoropsis* species (200 and 1000 m isobaths indicated). *A*, *Parvithracia* (s. str.) suteri (Finlay, 1927). *B*, *Parvithracia* (s. str.) ampla n. sp. C, *Trigonothracia mimica* n. sp. D, *Pendaloma micans* (Hedley, 1901). *E*, *Parvithracia* (*Pseudoasthenothaerus*) fragilissima n. sp. F, Myadoropsis wairua n. sp.

# Family PERIPLOMATIDAE Dall, 1895

#### Genus Pendaloma Iredale, 1930

*Pendaloma* Iredale, 1930: 387. Type species (by original designation): *Periploma micans* Hedley, 1901; Recent, New South Wales and New Zealand.

# Remarks

In his review of Pacific and Indo-Pacific Periplomatidae, Bernard (1989) made no mention of *Pendaloma* and merely listed the type species among the three recognised living species of Offadesma Iredale, 1930. Bernard's (1989) failure to compare the type species of his new genus Takashia Bernard, 1989 (Periploma plane Ozaki, 1958; Pleistocene and Recent, Japan) with P. micans suggests that he may have been unaware of Pendaloma, because it is unclear how Takashia actually differs from Pendaloma. The type species of Pendaloma differs from that of Offadesma (Cochlodesma angasi Crosse & P. Fischer, 1864; Recent, Australia and New Zealand) in attaining smaller size, in being equivalved and in being subcircular rather than subovate without a distinct rostrum. Both type species differ from the type species of Periploma Schumacher, 1817 (Periploma inaequivalvis Schumacher, 1817 = Corbula margaritacea Lamarck, 1801; Recent, Atlantic and Caribbean) markedly in shape and in lacking a lithodesma at maturity. The statements of Morton (1981) and Bernard (1989) that Offadesma angasi lacks a lithodesma are true of adults, although, in fact, a minute lithodesma is present in juveniles (width 270 µm in an 8-mm long individual), so evidently the structure is lost following arrested development at an early stage of growth. A lithodesma is also present in *Pendaloma micans* (Fig. 1M) and retained throughout life, although, again, development is arrested in early ontogeny and the lithodesma is eventually minute relative to the chondrophores. Another possible Pendaloma species is Periploma camerunensis Cosel, 1995 (off Cameroon), which is extremely similar to P. micans.

# Pendaloma micans (Hedley, 1901)

Figs 1*M*, 6*F*,*H*, 7*D* 

Periploma micans Hedley, 1901: 25, fig. 7. Pendaloma micans. – Iredale, 1930: 387. Periploma (Pendaloma) micans. – Keen, 1969b: 850, fig. F26/5a. Offadesma micans. – Bernard, 1989: 9. Pendaloma sp. Spencer et al. (in press).

#### Material examined

Holotype. 8 km E of Sydney, New South Wales, Australia, 82 m (AMS C8960).

Other material examined. Australia: New South Wales, shelf between Sydney and Cronulla, alive, 60–100 m (3pr, M.219809). New Zealand: E of North Cape,  $34^{\circ}23.0'S$ ,  $173^{\circ}12.6'E$ , 447-357 m (1v, M.144377). SW of Cape Maria van Diemen,  $34^{\circ}41.9'S$ ,  $172^{\circ}33.5'E$ , alive, 103 m (1pr, M.144360). W of Plate Island,  $37^{\circ}39.1'S$ ,  $176^{\circ}31.5'E$ , 64-59 m (1v, M.67585). ENE of Tokomaru Bay,  $37^{\circ}59.7'S$ ,  $178^{\circ}40.0'E$ , alive, 144 m (1pr, M.59857). N of New Plymouth,  $38^{\circ}40.2S$ ,  $174^{\circ}03.9'E$ , alive, 88 m (1pr, 1v, M.144370). NW of Manawatu R. mouth,  $40^{\circ}22.5'S$ ,  $174^{\circ}59.5'E$ , 86 m (1v, M.53863). NE of Stephens Island,  $40^{\circ}24'S$ ,  $174^{\circ}17'E$ , alive, 110 m (1pr, M.53095). E of Stephens Island,  $40^{\circ}38.5'S$ ,  $174^{\circ}12.0'E$ , 128 m (1v, M.53221). Between West Entry Point and Duffers Reef, Pelorus Sound mouth,  $40^{\circ}57.5'S$ ,  $174^{\circ}01.5'E$ , alive, 29 m (1pr, M.51510). Off Greymouth,  $42^{\circ}04.3'S$ ,  $170^{\circ}12.5'E$ , alive, 750 m (1pr, NZOI Q719). Pegasus Canyon, NE of Banks Peninsula:  $43^{\circ}20.3'S$ ,  $173^{\circ}31.4'E$ , alive, 622 m (4pr, M.90220);  $43^{\circ}24'S$ ,  $173^{\circ}27'E$ , 439-433 m (1v, M.49769);  $43^{\circ}24'S$ ,  $173^{\circ}26'E$ , alive, 366-311 m (10pr, M.51017);  $43^{\circ}26.9'S$ ,  $173^{\circ}29.5'E$ , alive, 505 m (1pr, M.144361). Bradshaw Basin floor, Bradshaw Sound,  $45^{\circ}17.3'S$ ,  $167^{\circ}02.6'E$ , alive, 415 m (1pr, M.138381).

# Description

Shell very thin and fragile, translucent white, subcircular or laterally flexed, length up to 29.5 mm, slightly longer than high, beaks opisthogyrous, valves similar, natural radial umbonal crack forming when shell approximately 4.5 mm long, posterior length approximately 38% of shell length; periostracum very thin, pale buff. Prodissoconch unknown. Posterodorsal margin straight, posterior margin subtruncate, anterodorsal margin weakly convex, ventral margin broadly convex; larger specimens typically shallowly indented anteroventrally and posteroventrally, marginal indentations coinciding with shallow radial concavities. Chondrophores large, stout, ligamental pit concave. Lithodesma before and against chondrophores, in juveniles smaller than chondrophores, although conspicuous; development arrested at an early stage of growth (probably at approximately 4.5 mm shell length, when umbonal crack begins to form) and, thus, relatively minute and inconspicuous in adults. Interior nacreous white, pallial sinus depth 39%-40% of shell length, posterior adductor scar subcircular, anterior adductor scar narrowly elliptical. Exterior dull, convex with shallow posterior and anterior radial concavities; fine, irregular, commarginal sculpture throughout, posterodorsal area typically with addition of eight to 12 fine radial threads; posterior radial concavity delineated by more or less well-defined lines, interspace of which is flat or becomes shallowly concave with increasing size; anterior radial concavity becoming apparent with increasing shell size, margins not sharply defined.

# Distribution

New South Wales. North and South Islands, New Zealand, living in mud at 29–750 m (Fig. 7D).

# Remarks

Compared with *Offadesma angasi* (Crosse & P. Fisher, 1864), the only other Recent periplomatid known from the New Zealand region, *Pendaloma micans*, differs principally in attaining smaller size (length of largest specimen seen 29.5 v. 102.5 mm) and in having a deeper, more broadly rounded pallial sinus, a well-defined band that curves posteriorly from the umbo to the posteroventral margin and fine collabral wrinkles and, typically, radial threads instead of minute but distinct, sharp granules on the posterodorsal area. Whereas *O. micans* lives at 29–750 m, *O. angasi* lives at low tide level to probably as deep as 64 m (very fresh paired juvenile from floor of Seymour Basin, Fiordland, New Zealand and New South Wales specimens are indistinguishable in shell morphology).

Other Anomalodesmata common to Australia and New Zealand are *Offadesma* angasi, Myochama tasmanica (Tenison Woods, 1877), *Cleidothaerus albidus* (Lamarck, 1819), *Spinosipella ericia* (Hedley, 1911) and, possibly, *Poromya australis* E. A. Smith, 1885 (recorded as *P. undosa* Hedley and Petterd, 1906 by Dell (1995); see Krylova (2001)). Genetic interchange between these disjunct populations is presumably maintained by larval stages of sufficient duration for planktotrophic larvae to reach New Zealand from Australia in eastward-moving water masses. However, genetic interchange may be intermittent and the rate of evolution in these taxa may simply be slow.

# Superfamily PANDOROIDEA Rafinesque, 1815

# Family **MYOCHAMIDAE** Bronn, 1862

#### Genus Myadoropsis Habe, 1960

Myadoropsis Habe, 1960: 293. Type species (by original designation): Myadora transmontana Yokoyama, 1922; Recent, Japan.

# Myadoropsis wairua n. sp.

Figs 3*F*, 7*F*, 8*A*,*D* 

Myadoropsis sp. Spencer et al. (in press).

#### Material examined

Holotype. N of Three Kings Islands, New Zealand, 34°01'S, 172°07'E, 622 m, 18 Feb. 1974, RV Acheron (left v, M.152686).

*Paratypes.* 37 km NE of Great Island, Three Kings Islands, 33°58.0'S, 172°30.6'E, 550 m (2v, M.149507). N of Three Kings Islands, 34°01'S, 172°07E, 622 m, (3v, M.34999). 22 km ENE of Great Island, 34°05.0'S, 172°24.6'E, 200 m (1v, M.93352). Off Spirits Bay, 34°21.60'S, 172°43.19'E, 48 m (2v, M.149506).

# Description

Shell up to 5.10 mm long, translucent white, massively secondarily thickened internally, strongly inequilateral, both valves modestly inflated, right valve slightly less inflated than left,  $1.62-1.64 \times$  longer than high; umbones strongly opisthogyrous, at posterior one-sixth to one-quarter; concave escutchion bounded by angulation. Prodissoconch approximately 270 µm wide, almost circular, convex, surface etched but apparently naturally smooth. Dissoconch anterodorsal and ventral margins very broadly rounded, posterodorsal margin concave, anterior margin well rounded, posterior margin obliquely subtruncate. Hinge plate extremely thick, anterior and posterior parts separated by deep, oblique ligamental groove; right valve with two strong, although poorly delineated, divergent teeth, antero- and posterodorsal margins grooved to interlock with corresponding, sharp, elevated margins of left valve. Pallial sinus depth 34%-36% of shell length. Adductor and anterior pedal retractor muscle scars well defined. Exterior typically with strong, rounded commarginal ridges, strongest over posterior half, obsolete in some specimens. Periostracum set with minute, crowded, commarginally elongate, hexagonal calcareous prisms that form the outermost shell layer.

# Distribution

Off Three Kings Islands and Spirits Bay, 48–550 m from comminuted bryozoan/shell substrata (shells only; Fig. 7*F*).

#### Remarks

*Myadoropsis wairua* differs from other nominal species of *Myadoropsis* principally in having a dorsoventrally lower posterior rostrum. Congeners are *M. elongata* (May, 1915) from south-eastern Australia and the Japanese species *M. transmontana* (Yokoyama, 1922), *M. dissimilis* Habe, 1960 and *M. brevispinosa* Habe, 1962. *Myadoropsis wairua* superficially resembles the thraciids *Barythaerus biconvexus* (Powell, 1927) and *B. cuneatus* (Powell, 1937; see above), but is immediately separable by the much heavier hinge area, the lack of a well-defined hinge tooth on the right valve and the distinctive microsculpture.



**Fig. 8.** Shells of *Myadoropsis* and *Myochama* species. *A,D, Myadoropsis wairua* n. sp.: holotype, left valve (*A* upper, *D*), length 4.70 mm, N of Three Kings Islands, 622 m (M.152686); paratype, right valve (*A* lower), length 4.40 mm, 37 km NE of Great Island, Three Kings Islands, 550 m (M.149507). *B,C,E–G, Myochama tasmanica* (Tenison Woods, 1877): *B,F,G*, juvenile, precementation, left (*B* upper, *G*) and right valve, length 2.90 mm, SE of Three Kings Islands, 121 m (M.152688); *C*, juvenile, early cemented stage, left valve, length 4.30 mm, Arch Pinnacle, Prince's Islands, Three Kings Islands, 40 m (M.117130). *E*, juvenile, precementation, right valve, length 2.55 mm, Bradshaw Basin floor, Bradshaw Sound, Fiordland, 415 m (M.138438).

# Etymology

From the Maori *wairua* (spirit), alluding to departure place of spirits in the vicinity of the type locality.

#### Genus Myochama Stutchbury, 1830

*Myochama* Stutchbury, 1830: 96. Type species (by monotypy): *Myochama anomioides* Stutchbury, 1830; Recent, Australia.

# Myochama tasmanica (Tenison Woods, 1877)

Figs 3G, 4H–J, 8B,C,E–G, 11A

Gouldia tasmanica Tenison Woods, 1877: 158.

Myochama woodsii Petterd, 1884: 145.

Myochama tasmanica. – Tate & May, 1901: 461, fig. 13; May, 1921: 14; May, 1923: pl. 5, fig. 11; Cotton
 & Godfrey, 1938: 140, fig. 135; Cotton, 1961: 149, fig. 140; Macpherson, 1958: 9, pl. 5, fig. 11;
 Powell, 1974: 197, figs 1–8; Powell, 1979: 432, text fig. 115; Lamprell & Healy, 1998: 216, fig. 632.

#### Material examined

*Syntypes of Myochama tasmanica*: Long Bay, Tasmania; Tasmanian Museum, Hobart (2v, E63/7404). *Myochama woodsii*: repository of type material unknown.

Other material examined. Off Three Kings Islands: Middlesex Bank, 33°57.0'S, 171°45.4'E, 98–103 m, 31 Jan. 1981 (1v, M.152690); King Bank, 33°57.0'S, 172°19.0'E, 128 m, 1 Feb. 1981 (3v, M.152689); King Bank, 33°57.4'S, 172°19.4'E, 128–123 m, 1 Feb. 1981 (1v, M.152691); Middlesex Bank, 33°59.8'S, 171°46.8'E, 143-163 m, 31 Jan. 1981 (1v, M.152687); off North-east Island, 34°08.5'S, 172°11'E, 102 m, 18 Feb. 1974 (1v, M.34578); E of Great Island, 34°10.5'S, 172°11.4'E, 153 m, 1 Feb. 1981 (1v, M.152692); Arch Pinnacle, Princes Rocks, 34°10.5'S, 172°03.0'E, 38 m, 1 Feb. 1983 (3v, M.154165); off Arch Pinnacle, 40 m, 10 Mar. 1983 (1v, M.117130); off West Island, *Elingamite* wreck, 34°11'S, 172°03'E, 37 m, 16 Mar. 1981 (3v, M.152693); 34°20.2'S, 172°21.8'E, 121 m, 2 Feb. 1981 (2pr, M.152688). Off Spirits Bay: 34°23.87'S, 172°45.62'E, 30 m, 24 Jan. 1999 (1v, M.153099); 34°24.31'S, 172°49.95'E, 29 m, 24 Jan. 1999 (2pr, 1v, M.153081). Parengarenga Harbour: 6-7 m (6v, M.117599); off Dog Island, 1979 (1pr, M.155465). Immediately outside Whangaroa Harbour entrance, 35°00.35'S, 173°45.7'E, 25 m, 26 Jan. 1981 (3v, M.158226). Bay of Islands: 35°11.5'S, 174°15.5'E, 54–58 m, 9 Feb. 1976 (1v, M.131122); Deepwater Cove entrance, 35°12'S, 174°18'E, 46 m, 10 Dec. 1973 (1v, M.41698). Off Ngaio Rock, Poor Knights Islands, 35°29.5'S, 174°44.0'E, alive, 22 m, 16 Apr. 1989 (1pr, 1v, M.158229). Bradshaw Basin floor, Bradshaw Sound, 45°17.3'S, 167°02.6'E, 415 m, 31 May 1997 (1v, M.138438). Off Taiaroa Head: head of Karitane Canyon, 45°38.5'S, 171°05.0'E, 585–530 m, 19 Mar. 1976 (1v, M.51115); 45°50.85'S, 171°01.71'E, alive, 555-604 m, 4 Jun. 1992 (2pr, 13v, M.130631); Saunders Canyon, 45°55'S, 170°56'E, 549 m, 2 Mar. 1954 (1v, M.9121). Stewart Island: off South Cape, 73-91 m (1v, M.63805); off Poutama Island, Big South Cape Island, 47°16'S, 167°23'E, 55 m, Jun. 1955 (1v, M.63787).

#### Description

(Supplemental to that of Powell, 1974). Prodissoconch approximately 250  $\mu$ m wide, smooth; PI approximately 230  $\mu$ m wide, globular. Periostracum set with minute, crowded, commarginally elongate, hexagonal calcareous prisms that form outermost shell layer. Following shell length of approximately 3 mm, when right valve becomes cemented to substratum, periostracum becoming fully calcified and rigid (rather than flexible as before) and essentially 'draped' from rib summit to summit of the commarginal ribs, so covering interstitial voids.

#### Distribution

South Australia, Tasmania and New Zealand: Three Kings Islands, Parengarenga Harbour, Matai Bay (Gardner 1976), Bay of Islands, Poor Knights Islands, East Otago, Fiordland and Stewart Island, living at 6–604 m cemented by right valve to shells (Fig. 11*A*).

# Remarks

The original description of *M. tasmanica* is based on juveniles in the pre-attachment stage (Tenison Woods, 1877). Cotton and Godfrey (1938: fig. 135) illustrated a juvenile and

provided an abridged version of the original description (both copied by Cotton (1961)), although, from the given dimensions  $(13 \times 13 \text{ mm})$ , it is clear that they were well aware that the type specimens were immature. Lamprell and Healy (1998) illustrated an adult, but their description seems to have been derived from their reproduction of the illustration of Cotton and Godfrey (1938: fig. 135). Both juvenile and adult Tasmanian specimens were illustrated by May (1923: pl. 5, fig. 11; reproduced by Macpherson (1958)). *Myochama tasmanica* differs from *M. anomioides* in attaining smaller size (width up to 18 v. 25 mm), in having radial grooves on the interior surface, in the considerably weaker inflation of the left (free) valve and in having commarginal rather than radial sculpture on the the left valve after settlement.

The prodissoconch of *Myochama tasmanica* has an extremely narrow PII stage (Fig. 4*H–J*), suggesting that larval development is non-planktotrophic, presumably lecithotrophic. I am unable to detect any differences between prodissoconchs of Australian and New Zealand specimens. Although development is apparently non-planktotrophic, the strongly disjunct distribution off the northern North Island and the southern South Island of New Zealand suggests initial origin from eastern Australia via north-eastward and south-eastward moving parts of the East Australian Current. The fact that *M. tasmanica* occurs at considerable depth off southern New Zealand, with a record there dating back to 1954, indicates that it is neither a 'warm water' species nor a recent invader, as findings in Parengarenga Harbour suggested to Powell (1976).

Shell structure, anatomy, cementation and substratum preference in *Myochama* ostreoides has been discussed by Harper and Morton (2000).

#### Genus Hunkydora Fleming, 1948

Hunkydora Fleming, 1948: 80. Type species (by original designation): Thracia transenna Suter, 1913 = T. novozelandica Reeve, 1859; Recent, New Zealand.

# Remarks

Fleming (1948) correctly referred Hunkydora to Myochamidae on the basis of characteristically myochamid exterior shell microsculpture and hinge morphology and clearly differentiated it from other myochamid genera. Keen (1969b) retained the genus in Myochamidae. Powell's (1974, 1976, 1979) subsequent placement of Hunkydora in Thraciidae (as a subgenus of *Thracia*) was due to confusion of *H. novozelandica* with the thraciids Asthenothaerus maxwelli n. sp., and Thracia vitrea (Hutton, 1873) (see above). Morton (1977) judged that Hunkydora belonged in Thraciidae from shell morphology of a specimen he identified as 'Hunkydora australica (Fleming, 1951)', although it is unclear whether this was H. novozelandica, T. vitrea or, seemingly most likely (because it is the most common of the three species in collections), A. maxwelli. Hunkydora species certainly are strikingly similar to several thraciids in gross shell facies, but are separable by the commarginally elongate, hexagonal, calcareous periostracal prisms, which, characteristically of myochamids, are arranged in sharply defined, irregular, wavy radial lines (unlike the superficially similar microsculpture of some thraciids, which comprise rounded elongate granules that are not grouped in radial lines; personal observation). Moreover, the left valve is considerably less inflated than the right valve, as in the majority of myochamids (i.e. markedly inequivalve), but, unlike thraciids, in which the left valve is, at most, only slightly less inflated than the right (note that myochamids of the genus Myadoropsis species are equivalve; see above). The outer shell layer on the interior is nacreous, as is characteristic of myochamids, although a few undoubted thraciids also have a nacreous layer (see above).

# Hunkydora novozelandica (Reeve, 1859)

Figs 1F, 4K, 9A–F, 11B, 21H

Thracia novozelandica Reeve, 1859: 12, pl. 3, fig. 19.

Thracia novaezeelandiae [sic]. - Martens, 1873: 41.

Thracia novaezealandiae [sic]. – Hutton, 1873a: 61; Hutton, 1880: 136.

Myadora australica. - Smith, 1885: 67 (in part not Reeve, 1859).

Thracia transenna Suter, 1913: 1023, pl. 55, figs 9, a.

Eximiothracia transenna. – Finlay, 1926: 474.

Hunkydora transenna. - Fleming, 1948: 80, pl. 4, fig. 8, pl. 7, figs 6-9.

Hunkydora australica novozelandica. - Fleming, 1951: 135, pl. 20, figs 3-5.

*Thracia vitrea.* – Dell, 1956: 41 (in part + *Asthenothaerus maxwelli*); Powell, 1957: 84 (in part: pl. 11, fig. 24 = *A. maxwelli*); Powell, 1962: 125 (in part: pl. 11, fig. 24 = *A. maxwelli*).

Myadora (Hunkydora) novozelandica. - Keen, 1969b: 847, figs 4a-c.

*Thracia (Hunkydora) australica novozelandica.* – Powell, 1976: 131 (in part = *Thracia vitrea* + *A. maxwelli* (pl. 18, fig. 24)); Powell, 1979: 433, pl. 79, fig. 9 only (in part = *T. vitrea* + *A. maxwelli* (pl. 79, figs 6, 7)).

NOT Thracia (Hunkydora) australica novozelandica. – Powell, 1974: 200, figs 13, 14 (= A. maxwelli).

#### Material examined

*Lectotype of Thracia novozelandica* (Fleming 1951). 'New Zealand', type locality here designated as Port Pegasus, Stewart Island (BMNH 20010411/1).

Paralectotype of Thracia novozelandica. Same data as lectotype (BMNH 20010411/2).

*Lectotype of Thracia transenna* (Boreham 1959). Port Pegasus, Stewart Island, 22 m (NZGS TM 404) *Paralectotype of Thracia transenna*. Same data as lectotype (1v, M.10984).

Other material examined. Fossil: large bend on Burma Road, between Whakatane and Ohope, brown siltstone, Late Pleistocene (Castlecliffian) (2v, M.96478); Languard Bluff, Wanganui (Castlecliffian) (2v, M.15757). Recent: off Three Kings Islands: King Bank, 33°57.0'S, 172°19.0'E, 128 m (4v, M.150900); King Bank, 33°57.4'S, 172°19.4'E, 128–123 m (2v, M.150875); Middlesex Bank, 33°59.8'S, 171°46.8'E, 143-163 m (1v, M.150883); 22 km ENE of Great Island, 34°05.0'S, 172°24.6'E, 200 m (1v, M.150881); E of Great Island, 34°10.5'S, 172°11.4'E, 153 m (1v, M.150885); S of Great Island, 34°14.1'S, 172°09.0'E, 192-202 m (2v, M.150876). NW of Cape Reinga, 34°14.2'S, 172°32.4'E, 100 m (2v, M.150884); off Spirits Bay, 34°17.38'S, 172°44.00'E, 85 m (2v, M.148268); off Spirits Bay, 34°18.36'S, 172°49.39'E, 68 m (1v, M.148258); N of North Cape, 34°20.0'S, 173°06.6'E, 163-168 m (2v, M.150893); SE of Three Kings Islands, 34°20.2'S, 172°21.8'E, alive, 121 m (2pr, 1v, M.150896). N of Cape Reinga, 34°21'S, 172°37'E, alive, 88 m (1pr, M.36018). NW of Cape Reinga, 34°22.8'S, 172°24.6'E, alive, 121 m (1pr, 1v, M.150890). Great Exhibition Bay: 34°33.4'S, 173°04.8'E, 63 m (4v, M.113152); 34°33.8'S, 173°04.6'E, 63–65 m (1v, M.150897). SW of Cape Maria van Diemen, 34°41.9'S, 172°33.5'E, 103 m (5v, M.150891). Off Doubtless Bay, 34°56.0'S, 173°34.0'E, 47 m (12v, M.25817). Bay in Stephensons Island, opposite Whangaroa Heads, 34°58'S, 173°47'E, 22–24 m (1v, M.41351; 2v, M.41553); Whangaroa Harbour entrance, main channel, 35°02'S, 173°45'E, 13 m (2pr, M.42013). Whangaroa Harbour, centre of Kaouou Bay, 35°02'S, 173°45'E, 13 m (2v, M.41766). Off Cape Brett, 35°08'S, 174°12.5'E, alive, 80 m (1pr, 6v, M.35731). Bay of Islands: NE of Ninepin Rock, 35°08.8'S, 174°10.9'E, 79 m (7v, M.28471); NE of Ninepin Rock, 35°08.8'S, 174°10.3'E, 75 m (18v, 30720); between Hope Passage and Twin Rocks, 35°10'S, 174°18'E, 46–55 m (2v, M.41260); off Takau Bay, 35°10'S, 174°11'E, 80 m (6v, M.43594); off Twin Rocks, 35°10'S, 174°18'E, 46-73 m (3v, M.42149); off Takau Bay, 35°10.5'S, 174°10'E, 37 m (3v, M.43485); 35°10.5'S, 174°19.3'E, 36-53 m (1v, M.95672); 35°10.6'S, 174°15.4'E, 67-73 m (3v, M.96009); 35°10.6'S, 174°19.6'E, 20-27 m (1v, M.95589); 35°11.2'S, 174°16.7'E, 58–64 m (2v, M.96082); off island at entrance to Deep Water Cove, 35°11.9'S, 174°17.1'E, alive, 47–49 m (1pr, 1v, M.95704); Deepwater Cove entrance, 35°12'S, 174°18'E, 46 m (3v, M.41714); Deepwater Cove entrance, 35°12'S, 174°18'E, 33–46 m (1v, M.41477); between Hope Passage and Twin Rocks, 35°12'S, 174°18'E, 37 m and 37-40 m (1v, M.41395; 1v, M.41289); W of Motuwheteke Island, 35°12.3'S, 174°16.4'E, 43 m (1v, M.96057); Waewaetorea Passage, 35°12.5'S, 174°12.9'E, 4 m (1v, M.39671); 35°12.6'S, 174°16.4'E, 32–40 m (2v, M.95877); Bamboo Bay, Moturua Island, 35°13.9'S, 174°11.3'E, 4-6 m (1v, M.44478; 1v, M.40823); Rawhiti Bay, 35°14.0'S, 174°15.3'E, 2-4 m (1v, M.40801); 35°15.4'S, 174°11.5'E, near Knob Point, 4 m (1v, M.44640); Orakawa Bay, 35°15.3'S, 174°11.8'E, 4 m (1v, M.44451); Western Manawara Bay, 35°16.0'S, 174°11.5'E, 6–7 m (1v, M.44756). Off Poor Knights Islands: 35°22'S, 174°43'E, 146 m (4v, M.35105); 35°29'S, 174°43.5'E, 110 m (1v, M.35198); 35°32'S, 174°41'E, 121-113 m (6v, M.35807). Whangarei Harbour, off point SW of Munro Bay, 35°47.8'S, 174°28.9'E, alive, 7–10 m (1pr, M.152673); Castle Rock, Taurikura Bay, Whangarei Heads, 6-8 m (1v, M.43265); Taurikura Bay (2pr, 4v, M.18353). S of Lady Alice Island, Hen and Chicken Islands, 35°55.0'S, 174°43.5'E, 47–55 m (1v, M.150880). Off Hen Island, 36°00.5'S, 174°43'E, 59 m (6v, M.43789). SE of Te Arai Point, Mangawhai, Northland, 36°10.5'S, 174°42.0'E, 38-40 m, sand and shell suction dredged for beach replenishment at Mission Bay, Auckland (6v, M.138169; 3v, M.130830). NE of Takatu Point, 51–57 m (1v, M.21674); off Cuvier Island, 49–73 m (1v, M.3461). Omaha Bay: 36°20.41'S, 174°49.38'E, alive, 17 m (1pr, M.136165); 36°21.38'S, 174°51.03'E, alive, 28 m (1pr, M.136164). N of Aldermen Islands, 36°47.5'S, 176°00.0'E, 108-113 m (1v, M.150898). Off E side of Mayor Island, 37°18.9'S, 176°16.2'E, 59–74 m (1v, M.65726). Off White Island: 37°30.4'S, 177°09.7'E, 83–92 m (1v, M.150892); 37°30.6'S, 177°09.7'E, 73-59 m (1v, M.150886). Rungapapa Knoll, W of White Island, 37°33.5'S, 176°59.4'E, 198–188 m (2v, M.59995). Ranfurly Bank, East Cape: 37°32.8'S, 178°48.7'E, 94 m (6v, M.60708); 37°33.1'S, 178°49.5'E, alive, 94–89 m (4pr, 10v, M.71765); 37°33.4'S, 178°48.3'E, alive, 106-103 m (1pr. 3v, M.71784); 37°35.8'S, 178°47.0'E, 118-119 m (1v, M.150889); 37°38.5'S, 178°56.4'E, alive, 153-143 m (1pr, 3v, M.150899). Off Plate Island, 37°39.4'S, 176°34.4'E, 82 m (6v, M.67685). Between Motuhora and White Islands, 37°40.2'S, 178°53.6'E, 117 m (1v, M.112426). W of Plate Island, 37°39.1'S, 176°31.5'E, alive, 64-59 m (1pr, 9v, M.79735). NNE of Hicks Bay, 37°32'S, 178°25'E, alive, 108-134 m (1pr, M.16936). SE of Plate Island, 37°43.4'S, 176°38.5'E, 59 m (1v, M.65256). N of Motuhora Island, 37°45.8'S, 177°00.8'E, 72-84 m (1v, M.60616); ENE of Tolaga Bay, 38°15.2'S, 178°38.6'E, alive, 139 m (1pr, M.59803). S of Patea, 39°56'S, 174°26'E, 40 m (1v, M.95574). S of Waverley, 40°10'S, 174°40'E, 77-82 m (3v, M.50644). W of Rangitikei R. mouth, 40°16'S, 174°58.5'E, 75 m (2v, M.53363). NW of Manawatu R. mouth, 40°22.5'S, 174°59.5'E, alive, 86 m (2pr, M.53881). NE of Stephens Island, 40°33'S, 174°07'E, 132 m (2v, M.52998). Off shelf of flats inside Fairwell Spit, 40°36.5'S, 173°00.5'E, 26-24 m (2v, M.50022). N of Kapiti Island, 40°38'S, 174°54.5'E, 91 m (2v, M.52935). E of Stephens Island: 40°38.5'S, 174°12.0'E, 128 m (3v, M.53214); 40°42'S, 174°10.6'E, 106 m (1v, M.12929). Main channel, Port Hardy, D'Urville Island, 40°46.0'S, 173°52.5'E, alive, 18 m (1pr, 4v, M.58956). W of Kapiti Island, 40°46.2'S, 174°27.0'E, 137 m (1v, M.12927). N of Alligator Head and E of Rangitoto Islands, Marlborough Sounds, 40°47'S, 174°10.5'E, 73 m (6v, M.54089). E of D'Urville Island, Cook Strait, 40°48.6'S, 174°06.6'E, 60 m (1v, M.150888). Kapiti Channel, 40°51'S, 174°58'E, 64 m (8v, M.52233). W of Kapiti Island, 40°52.6'S, 174°49.5'E, 137 m (4v, M.150879). Between West Entry Point and Duffers Reef, Pelorus Sound mouth, 40°57.5'S, 174°01.5'E, 29 m (1v, M.51517). Off White Rocks, Queen Charlotte Sound, 41°04.3'S, 174°22.5'E, 46 m (1v, M.150882). Pegasus Canyon wall, NE of Banks Peninsula, 43°25.0'S, 173°26.0'E, alive, 329-183 m (1pr, M.71298). Off Oamaru: 110 m (1v, M.11361); 55-88 m (7v, M.111171); head of Waitaki Canyon, 45°10'S, 171°30'E, 293-256 m (12v, M.51261). Open Cove, Thompson Sound, 45°10.5'S, 167°05.0'E, 37 m (1v, M.84086). Doubtful Sound: entrance, 45°16'S, 166°48'E, 117-128 m (3v, M.144398); 0.8 km SW of Hare's Ears, 45°17.2'S, 166°49.3'E, 146 m (10v, M.58770); Kellard Basin floor, 45°21.30'S, 167°03.36'E, alive, 376 m (1pr, M.138600). Dusky Sound, 55 m (1v, M.15935). Off Taiaroa Heads: 45°36.55'S, 170°51.88'E, 41-59 m (2v, M.112307); Karitane Canyon, 45°37.5'S, 171°03'E, 420 m (1v, M.45553); Karitane Canyon, 45°38.5'S, 171°01'E, 200 m (1v, M.45434); 45°41.85'S, 170°58.72'E, 100–97 m (1v, M.112332); Taiaroa Canyon, 45°45.4'S, 171°05'E, alive, 549 m (1pr, 2v, M.9040); 45°50'S, 170°56'E, 105 m (9v, M.45376); Papanui Canyon, 45°51'S, 171°00'E, 348-220 m (1v, M.94878). Kisbee Bay, Preservation Inlet, 46°07.1'S, 166°44.0'E, 5–10 m (1v, M.59271). Off Taieri, S of Dunedin, 46°09.0'S, 170°27.0E, 60 m (3v, M.66937); 46°15.0'S, 170°29.0'E, 91 m (4v, 66171). Stewart Island: Easy Harbour, 47°09'S, 167°34'E, 7-20 m (1v, M.26518); North Arm, Port Pegasus, 47°11'S, 167°41'E, alive, 37-44 m (2pr, 2v, M.44788); North Arm, 47°11'S, 167°41'E, 40-46 m (2v, M.26619).

# Description

Shell up to 42 mm long, of moderate thickness; left valve smaller than right valve although more elongate, approximately half as inflated, lying within its perimeter; interior nacreous white, margins opaque white, some specimens with a yellowish patch near centre of valve. Prodissoconch 220–250 µm wide; PI approximately 200 µm wide, minutely pitted, circular, strongly inflated. Dissoconch anterodorsal margin flat or broadly convex, posterodorsal margin broadly concave, anterior margin tightly rounded, ventral margin broadly rounded; right valve posterior margin set off from posterodorsal margin by angulation, flat, weakly convex or shallowly concave, markedly arched relative to valve plane; left valve posterior margin subtruncate, almost vertical, more or less flat, smoothly merging with posterodorsal

margin, plane flexed to contact corresponding arched part of right valve. Exterior: narrow, flat or shallowly concave anterodorsal area of left valve defined by angulation, sharp throughout or progressively weakening; ovate, shallowly concave escutchion bounded by angulation on left valve only, margin on right valve tightly rounded; posterior one-quarter of left valve distinctly flattened, elsewhere broadly and rather evenly convex; posterior onequarter of right valve defined by low rounded radial swelling, in most specimens followed posteriorly by shallow radial concavity (lacking in few specimens), then rounded radial swelling that extends to escutchion margin, elsewhere broadly convex. Right valve hinge with small tooth at anterior extremity of posterodorsal margin, otherwise edentulous, anterior and posterior parts separated by narrow V-shaped space for lithodesma; right valve with grooved shelf along inside of most of antero- and posterodorsal margins to interlock with corresponding sharp, vertically elevated margins of left valve; posterior groove deeper, its anterior end thickened and elevated. Pallial line, pallial sinus and adductor muscle scars well defined, adult pallial sinus depth 41%-43% of shell length. Lithodesma stout (width 1.65 mm in shell 11.2 mm long), butterfly shaped, bilaterally asymmetrical, ligaments rather thick. Resilifers elevated; L-shaped on right valve, long oblique dorsoventral limb narrowly tapered, short anterodorsal limb at approximately right angle to it; anterolateral part of left valve resilifer smoothly curving to dorsoventral part, otherwise similar to right valve resilifer. Exterior sculptured with rounded commarginal ridges, weaker and fewer on left valve than on right. Periostracum set with minute, crowded, commarginally elongate, hexagonal, calcareous prisms that form outermost shell layer.

# Distribution

Late Pleistocene (Castlecliffian) to Recent. Three Kings, North, South and Stewart Islands, 4–549 m; taken alive (rarely) at 12–549 m (Fig. 11*B*).

# Remarks

Specimens from carbonate-rich substrata tend to attain larger size and have weaker commarginal ridges than specimens from mud, but otherwise appear indistinguishable in shell morphology. After repeated comparison of several hundred specimens, I have concluded that the two variants are ecophenotypes.

As indicated by the synonymy, *Hunkydora novozelandica* has been thoroughly confused with *Asthenothaerus maxwelli* n. sp., and *Thracia vitrea* (Hutton, 1873). Although similar in shape, *H. novozelandica* differs markedly from both these species in numerous details, principally: left valve far more weakly convex than right valve and seated well within perimeter of right valve in paired individuals, posterior of right valve strongly arched right, posterior of left valve flexed right to match right valve; prominent commarginal sculpture on right valve at an early stage of growth and on rostrum to maturity, both valves externally covered with a thin layer of minute, crowded, commarginally elongate, hexagonal prisms. Similarities in gross shell morphology between *Hunkydora* and *Thracia* species are probably due to convergence (see Harper *et al.* 2000).

Following Fleming (1951), *H. novozelandica* has consistently been interpreted as a subspecies of *H. australica* (Reeve, 1859), but the two are distinct species of *Hunkydora*, the eastern Australian one differing principally in being higher, in having the umbones situated further posteriorly and in that the anterior end of the inner posterodorsal margin of the right valve is thicker, higher and more tooth like. *Hunkydora australica* (three syntypes BMNH 1962695 examined), which is clearly referable to *Hunkydora*, was erroneously returned to *Thracia* by Lamprell and Healy (1998) without explanation.



**Fig. 9.** Shells of *Hunkydora* species. *A–F*, *Hunkydora novozelandica* (Reeve, 1859): *A,B*, right valve, length 21.8 mm, W of Motuwheteke Island, Bay of Islands, 42–43 m (M.96057); *C,D*, left valve, length 20.5 mm, Ranfurly Bank, East Cape, 103–106 m (M.71784); *E,F*, juvenile, length 4.85 mm, off Cape Brett, 80 m (M.35731). *G–I*, *Hunkydora rakiura* n. sp.: holotype, left (*I*) and right valve, length 5.20 mm, channel between Ulva Island and Bradshaw Peninsula, Paterson Inlet, Stewart Island, 33 m (M.152724).

# Hunkydora rakiura n. sp.

Figs 4L, 9G–I, 11C

Hunkydora sp. Spencer et al. (in press).

# Material examined

*Holotype.* Channel between Ulva Island and Bradshaw Peninsula, Paterson Inlet, Stewart Island, New Zealand, 46°56.5'S, 168°07.8'E, alive, 33 m, 12 Jan. 1952, MV *Alert* (pr, M.152724).

*Paratypes.* Off East Otago, Saunders Canyon, 45°56'S, 170°54'E, 360 m (3v, M.48858). Off Ruapuke Island, Foveaux Strait, oyster beds, 46°43'S, 168°30'E, 33–35 m (1v, M.71863).

# Description

Shell up to 5.20 mm long, translucent white, inequilateral, right valve more inflated and higher than left,  $1.4 \times$  longer than high; umbones opisthogyrous, at approximately posterior one-third; large, concave escutchion bounded by angulation. Prodissoconch approximately 300 µm wide, almost circular, smooth; PI approximately 270 µm wide, globular. Dissoconch anterodorsal margin very broadly rounded or almost flat, posterodorsal margin concave, anterior margin tightly rounded, posterior margin obliquely subtruncate, ventral margin very broadly rounded. Antero- and posterodorsal margins of valves sharp, upright, margins of left valve seated against ventral side of margins of right valve on well-defined corresponding ledges. Right valve hinge with small tooth at anterior extremity of posterodorsal margin, otherwise edentulous, anterior and posterior parts of hinge area separated by triangular ligamental space; large resilifers sharply delineated, slightly elevated. Lithodesma unknown. Interior nacreous white, pallial sinus depth 40% of shell length. Adductor and anterior pedal retractor muscle scars well defined. Periostracum set with minute, crowded, commarginally elongate, hexagonal calcareous prisms that form outermost shell layer.

#### Distribution

South-eastern South Island and Stewart Island, 33–360 m; taken alive at 33 m from bryozoan/shell substratum (Fig. 11*C*)

# Remarks

*Hunkydora rakiura* differs markedly from *H. novozelandica* in attaining considerably smaller size (length up to 5.20 v. 42 mm) and in having a larger prodissoconch (width 270–300 v. 220–250  $\mu$ m). Comparing adults of *H. rakiura* with juveniles of *H. novozelandica* of the same shell size, *H. rakiura* differs further in being lower (length/height ratio right valve 1.40 v. 1.26) and in that the beaks are smaller, sharper and more strongly posterior, the anterior end is more broadly rounded, the posterodorsal margin is more deeply concave and the rostrum is dorsoventrally shorter and more weakly arched.

# Etymology

After rakiura, the Maori name for Stewart Island, the type locality.

# Superfamily CUSPIDARIOIDEA Dall, 1886

# Family CUSPIDARIIDAE Dall, 1886

# Genus *Pseudogrippina* n. gen.

Type species: Pseudogrippina wanganellica n. sp.; Recent, Wanganella Bank, southern Norfolk Ridge.

#### Diagnosis

Shell up to 2.80 mm long, roundly trigonal, posteriorly obliquely truncate, right valve with strong anterior tooth and no posterior tooth, left valve edentulous, resilifers sunken, lithodesma present, interior surface shell layer a chaotic mass of elongate prisms, thin periostracum minutely pitted.

# Remarks

The type species of *Pseudogrippina* has striking similarity to *Grippina* species in shape and size, including the presence of a lithodesma set on sunken resilifers, but it differs in that the

# 254



Fig. 10. Prodissoconchs of Cuspidariidae and Spheniopsidae (prodissoconch I and II boundaries or prodissoconch boundary indicated by arrows). A, Plectodon regalis n. sp., holotype, off Three Kings Islands, New Zealand, 91 m (M.152684). B, Rhinoclama (Austroneaera) brevirostris Powell, 1937, SE of Great Island, 173-178 m (M.144426). C, Rhinoclama (Austronearea) finlayi Powell, 1937, N of Poor Knights Islands, 146 m (M.144424). D, Rhinoclama (Austroneaera) brooki n. sp., holotype, SE of Nugent Island, Raoul Island, Kermadec Islands, 165-146 m (M.152685). E, Rhinoclama (Austroneaera) raoulensis Powell, 1958, between Dayrell and Chanter Islets, Raoul Island, Kermadec Islands, 31-45 m (M.226999). F, Rhinoclama (Austroneaera) tangaroa n. sp., paratype, Wanganella Bank, Norfolk Ridge, 437-422 m (M.234116). G, Rhinoclama (s. str.) aupouria (Dell, 1950), SE of Great Island, 173-178 m (M.144427). H, Grippina aupouria (Powell, 1937), 11 km NW of Great Island, Three Kings Islands, 310 m (M.93848). I, Grippina rex n. sp., holotype, off Three Kings Islands, 91 m (M.33807). J, Grippina punctata n. sp., holotype, off Spirits Bay, 63 m (M.152679). K, L, Grippina globosa n. sp., paratype, Middlesex Bank, NW of Three Kings Islands, 246-291 m (M.149490). M, Grippina spirata n. sp., holotype, off Spirits Bay, 63 m (M.149519). N, Grippina acherontis n. sp., paratype, off Patea, 40 m (M.53594). O, Grippina pumila n. sp., paratype, off Puysegur Point, 183 m (M.143883). Scale bars: 50 µm.

right valve has only one relatively larger anterior hinge tooth, which stands obliquely instead of at a right angle to the valve plane, the inner surface shell layer is a chaotic mass of elongate prisms instead of smooth and the periostracum is minutely pitted throughout.

The presence of an anterior tooth, lack of a posterior tooth on the right valve and lack of teeth on the left valve are diagnostic characteristics of *Luzonia* Dall & E. A. Smith in Dall, 1890 (type species *Neaera philippinensis* Hinds, 1843; Recent, Philippine Islands; ICZN (1986)). Checking of the type material of *N. philippinensis* (BMNH 1961122) reveals that it comprises three specimens rather than a 'type specimen' as stated by Allen and Morgan (1981). Moreover, two species are represented, one by a complete right valve (length 5.65 mm), the other (smaller) by a live-taken closed pair and a complete right valve and a left valve hinge fragment, evidently from a single individual. Both show the hinge morphology diagnostic of *Luzonia* and both are posteriorly rostrate with small resilifers set in the hinge line. *Pseudoneaera wanganellica* differs from these syntypes of *L. philippinensis* in numerous details, including a much shorter, obliquely truncate posterior end and correspondingly shallower pallial sinus, much stronger tooth on the right valve and in that the resilifers are sunken rather than within the hinge line. Placement of *Pseudogrippina* in Cuspidariidae is provisional pending knowledge of the anatomy.

The specimen selected as lectotype for *N. philippinensis* should, ideally, be the paired individual, but this formality is left to some future worker because I am reluctant to try and separate the valves.

# Etymology

From the Greek pseudos (lie) + genus name Grippina.

# Pseudogrippina wanganellica n. sp.

Figs 3H,I, 4M, 11D, 12A,C, 21I

#### Material examined

Holotype. Wanganella Bank, Norfolk Ridge, summit, 32°34.4'S, 167°31.0'E, alive, 113 m, 29 Jan. 1981, RV Tangaroa (pr, M.152681).

*Paratypes.* Wanganella Bank: summit, 32°31.8′S, 167°29.5′E, 113–118 m (1v, M.257207); Norfolk Ridge, summit, 32°34.4′S, 167°31.0′E, alive, 113 m, (2pr, M.149523); E slope, 32°35.3′S, 167°41.8′E, 437–422 m (4v, M.234118); summit, 32°40.0′S, 167°33.6′E, 133 m (1pr, 4v, M.272623); summit, 32°39.2′S, 167°31.7′E, 133 m (6v, M.272622).

#### Description

Shell up to 2.80 mm long, ovate trigonal, of moderate thickness, moderately inflated, approximately  $1.27\times$  longer than high; beaks almost central, anterior end elongating slightly more rapidly than posterior in large specimens so that posterior end occupies 50%-54% of shell length; fresh specimens translucent white. Prodissoconch approximately 240 µm wide, prominent, oval, smooth; finely wrinkled, elevated, convex central area with shallow depression at summit, otherwise smooth; no PII discernable. Dissoconch anterior margin well rounded, smoothly merging with straight anterodorsal margin and broadly rounded ventral margin, posterodorsal margin very broadly rounded; posterior truncation short, broadly rounded or almost straight, oblique, extremities roundly angulate. Right valve with one strong, well-defined anterior hinge tooth, no posterior tooth, inner anterodorsal margin shelved to seat corresponding elevated narrow dorsal margin of left valve. Left valve edentulous. Lithodesma stout (width 230 µm in adult), longer than wide,



Fig. 11. Maps of the New Zealand region showing distributions of *Myochama*, *Hunkydora*, *Pseudogrippina* and *Plectodon* species (200 and 1000 m isobaths indicated). A, *Myochama tasmanica* (Tenison Woods 1877). B, *Hunkydora novozelandica* (Reeve, 1859). C, *Hunkydora rakiura* n. sp. D, *Pseudogrippina wanganellica* n. sp. E, *Plectodon lepidus* n. sp. F, *Plectodon pruinosus* n. sp.

subtriangular, rounded anteriorly, concave posteriorly, ligaments thin, sunken resilifers shallowly concave, slightly but distinctly elevated. Interior surface a layer of chaotically oriented, narrowly elongate prisms, pallial line and large adductor scars well defined; pallial sinus concave, depth approximately 30% of shell length. Exterior convex, posterior end flattened; sculptured throughout with weak, rounded commarginal ridges and faint radial lines. Periostracum thin, colourless, minutely pitted.

# Distribution

Wanganella Bank, southern Norfolk Ridge, 133–437 m; taken alive at 113–133 m from substratum of rhodolith gravel and white biogenic sand (Fig. 11*D*).

# Remarks

*Pseudogrippina wanganellica* resembles *Luzonia philippensis* (Hinds, 1843) (Philippines, depth unknown) and *L. simplex* (Allen & Morgan, 1981) (western Africa, 619–2357 m), but differs in shape (more ovate than trigonal, with shorter and higher rostrum), stronger commarginal sculpture, smaller size (length 2.8 v. 3.4 mm) and in that the resilifers are sunken rather than set in the hinge line. For additional remarks, see under *Pseudogrippina*.

# Etymology

After Wanganella Bank, the type locality.

### Genus Plectodon Carpenter, 1864

Plectodon Carpenter, 1864: 611. Type species (by monotypy): Plectodon scaber Carpenter, 1864; Recent, California.

# Remarks

The four new species attributed to *Plectodon* described below resemble the holotype of *Plectodon scaber* (Palmer 1958: pl. 6, figs 6–8, subsequently fragmented) and other Californian specimens examined in gross shell morphology, including the presence of crisp calcareous granules set in the periostracum, and in having a substantial lithodesma set on oblique sunken resilifers (Coan *et al.* 2000: 559, pl. 121). However, they differ principally in attaining much smaller size (length up to 6.25 v. 19 mm), in that the sunken resilifers are simply shallow pits rather than distinctly elevated, the hinge teeth are smooth instead of pustulated dorsally and in being more weakly rostrate. *Plectodon scaber* differs further in that the posterior extremity of the anterodorsal margin of the right valve is spirally twisted immediately below the umbo.

The only other strictly similar species known to me is *P. granulatus* (Dall, 1881) from the western central Atlantic. Poutiers and Bernard (1995) referred *Neaera brazieri* E. A. Smith, 1885 (Australia) and *Cuspidaria ligula* Yokoyama, 1922 (Japan) to *Plectodon*, but, according to original descriptions, they lack granulate microsculpture.

# Plectodon lepidus n. sp.

# Figs 3J,K, 4N, 11E, 12B,D, 21J

Plectodon sp. 2 Spencer et al. (in press).

# Material examined

*Holotype.* Off E side of Mayor Island, New Zealand, 37°18.9'S, 176°16.2'E, alive, 59–74 m, 22 Jan. 1979, RV *Tangaroa* (pr, M.152682).

*Paratypes.* Off Rangaunu Bay,  $34^{\circ}42.8'S$ ,  $173^{\circ}14.5'E$ , 63 m (8v, M.149594). NW of Whangaroa Harbour,  $34^{\circ}54.0'S$ ,  $173^{\circ}42.6'E$ , 83 m (1v, M.149596). Bay of Islands: off Twin Rocks,  $35^{\circ}10'S$ ,  $174^{\circ}18'E$ , 46-73 m (3v, M.42188);  $35^{\circ}11.2'S$ ,  $174^{\circ}16.7'E$ , 58-64 m (1v, M.96122); off island at entrance to Deep Water Cove,  $35^{\circ}11.9'S$ ,  $174^{\circ}17.1'E$ , 47-49 m (1v, M.95725); off Deep Water Cove, fish gut (1v, M.112094). Whangaruru Harbour, dredged (2v, M.111956). Off E side of Mayor Island,  $37^{\circ}18.9'S$ ,  $176^{\circ}16.2'E$ , alive, 59-74 m, (3 pr 17v, M.65656). Off White Island,  $37^{\circ}30.5'S$ ,  $177^{\circ}09.7'E$ , 64-69 m, two stations (2v, M.149595; 2v, M.96298);  $37^{\circ}30.6'S$ ,  $177^{\circ}09.7'E$ , 73-59 m (7v, M.94488).

# Description

Shell up to 6.10 mm long, ovate-trigonal, of moderate thickness, moderately inflated,  $1.52-1.60 \times$  longer than high; beaks almost central, fresh specimens translucent white with

narrow, opaque white radial rays. Prodissoconch approximately 230  $\mu$ m wide, prominent, smooth; PI approximately 200  $\mu$ m wide, globular. Dissoconch anterior margin well rounded, smoothly merging with straight anterodorsal margin and broadly rounded ventral margin; posterodorsal margin longer than anterodorsal, straight; posterior truncation more or less flat, slightly oblique, corners roundly angulate. Right valve with two strong, well-defined anterior and posterior hinge teeth, length of inner dorsal margins each with concave shelf to seat corresponding elevated narrow dorsal margins of left valve. Left valve with prominent, posteriorly inclined boss at posterior end of anterodorsal margin. Lithodesma stout (width 500  $\mu$ m in adult), subtriangular, rounded anteriorly, concave posteriorly; ligaments thin, sunken resilifers shallowly concave. Interior with faint, shallow radial grooves that approximately coincide with white lines, pallial line and large adductor scars well defined; pallial sinus shallow, concave. Exterior convex, posterior area flattened, separated by low, broadly rounded angulation; crisply sculptured throughout with crowded, roughly hemispherical, calcareous granules set in thin periostracum, stronger on posterior area.

#### Distribution

North-eastern North Island (34°42.8'-37°30.6'S), 46–83 m; taken alive at 59–74 m from substratum of clean shell and stones (Fig. 11*E*).

# Remarks

Two similar species occur in the New Zealand region and are described below. Another species occurs off Norfolk Island (1 left v, M.225171).

# Etymology

From the Latin *elegans* (elegant).

# Plectodon pruinosus n. sp.

Figs 3L, 4O, 11F, 12E-G

Plectodon sp. 3 Spencer et al. (in press).

#### Material examined

*Holotype.* Middlesex Bank, NW of Three Kings Islands, New Zealand, 33°57.0'S, 171°45.4'E, 98–103 m, 31 Jan. 1981, RV *Tangaroa* (right v, M.152683).

*Paratypes.* Off Three Kings Islands: Middlesex Bank, 33°57.0'S, 171°45.4'E, 98–103 m (2v, M.149573); Middlesex Bank, 33°59.9'S, 171°45.3'E, 186–196 m (3v, M.112807); Middlesex Bank, 34°00.9'S, 171°44.7'E, 201–216 m (2v, M.93232); 34°01'S, 172°07'E, 622 m (2v, M.34952); 13 km N of Great Island, 34°01.8'S, 172°12.0'E, 508 m (1v, M.149574); Middlesex Bank, 34°02.0'S, 171°44.0'E, 246–291 m (3v, M.149571); Middlesex Bank, 34°02.1'S, 171°45.8'E, 221–206 m (1v, M.149570); 24 km NW of Great Island, 34°05.9'S, 171°55.1'E, 710 m (2v, M.94367); S of Great Island, 34°14.1'S, 172°09.0'E, 192–202 m (1v, M.149575); 34°14.8'S, 172°13.6'E, SE of Great Island, 173–178 m (1v, M.149572). NW of Cape Reinga, 34°14.2'S, 172°32.4'E, 100 m (3v, M.149576). Off Spirits Bay, 34°18.36'S, 172°49.39'E, 68 m (1v, M.152487).

# Description

Shell up to 6.25 mm long, ovate–trigonal, of moderate thickness, moderately inflated,  $1.60-1.79 \times$  longer than high; beaks almost central, fresh specimens translucent white. Prodissoconch approximately 230 µm wide, prominent, convex, central area elevated, finely wrinkled dorsally, minutely and densely maleate; no PII. Dissoconch anterior margin well rounded, smoothly merging with broadly rounded anterodorsal and ventral margins;



**Fig. 12.** Shells of *Pseudogrippina* and *Plectodon* species. *A,C, Pseudogrippina wanganellica* n. sp.: holotype, left (*A* upper) and right valve, length 2.30 mm, Wanganella Bank summit, Norfolk Ridge, 113 m (M.152681). *B,D, Plectodon lepidus* n. sp.: holotype, left (*B* upper) and right valve, length 5.25 mm, off E side of Mayor Island, 59–74 m (M.152682). *E,G, Plectodon pruinosus* n. sp.: holotype, right valve (*E,G* lower), length 5.72 mm (M.152683), and paratype, left valve (*G* lower), length 4.63 mm (M.149573), Middlesex Bank, NW of Three Kings Islands, 98–103 m. *F,H, Plectodon regalis* n. sp.: holotype, left (*H* upper) and right valve, length 5.20 mm, off Three Kings Islands, 91 m (M.152684).

posterodorsal margin straight; posterior truncation broadly rounded or flat, slightly oblique, corners tightly rounded. Right valve with two strong, well-defined anterior and posterior hinge teeth, length of inner dorsal margins each with concave shelf to seat corresponding elevated narrow dorsal margins of left valve. Left valve with prominent, posteriorly inclined boss at posterior end of anterodorsal margin. Resilifers sunken, oblique, elongate–oval, concave, ventral margin slightly elevated (lithodesma almost certainly present although missing in isolated valves). Interior with faint, shallow radial grooves, pallial line and large adductor scars well defined, pallial sinus broad and shallow. Exterior convex, posterior area weakly convex, separated by low, broadly rounded angulation; crisply sculptured

throughout with crowded, roughly hemispherical, calcareous granules set in thin periostracum, stronger on posterior area.

#### Distribution

Off Three Kings Islands and off Spirits Bay, 98–710 m (shells only), from comminuted bryozoan/shell substrata (Fig. 11*F*).

#### Remarks

*Plectodon pruinosus* differs from *P. lepidus* in having shallowly curved rather than flat dorsal margins, finer granulate sculpture, a larger prodissoconch with a more elevated central area, a dorsoventrally shorter posterior truncation and in that the pallial sinus is broader. The two species are allopatric. *Plectodon pruinosus* occurs sympatrically with the species described below.

# Etymology

From the Latin pruinosus (frosty).

#### Plectodon regalis n. sp.

# Figs 3M, 10A, 12F,H, 15A, 21K

*Cuspidaria aupouria.* – Crozier, 1966: 46, figs 16, 17; Powell, 1979: 436. In part + three undescribed species of *Cuspidaria*. Not Dell, 1950. *Plectodon* sp. 1 Spencer *et al.* (in press).

#### Material examined

Holotype. Off Three Kings Islands, New Zealand, 34°11′S, 172°10′E, alive, 91 m, 19 Feb. 1974, RV Acheron (pr, M.152684).

*Paratypes.* Off Three Kings Islands: Middlesex Bank,  $33^{\circ}57.0'S$ ,  $171^{\circ}45.4'E$ , 98-103 m (2v, M.149582); King Bank,  $33^{\circ}57.4'S$ ,  $172^{\circ}19.4'E$ , alive, 128-123 m (1pr, 1v, M.149581); Middlesex Bank,  $33^{\circ}59.8'S$ ,  $171^{\circ}46.8'E$ , 143-163 m (1v, M.149597); Middlesex Bank,  $34^{\circ}01.2'S$ ,  $171^{\circ}44.4'E$ , 206-211 m (1v, M.149579); 13 km N of Great Island,  $34^{\circ}01.8'S$ ,  $172^{\circ}12.0'E$ , 508 m (3v, M.149583); 37 km NW of Great Island,  $34^{\circ}02.0'S$ ,  $171^{\circ}48.4'E$ , 188 m (3v, M.64914); Middlesex Bank,  $34^{\circ}02.0'S$ ,  $171^{\circ}44.4'E$ , 246-291 m (3v, M.149586); 11 km NW of Great Island,  $34^{\circ}06.5'S$ ,  $172^{\circ}04.7'E$ , 310 m (1v, M.149577); off North-east Island,  $34^{\circ}08.5'S$ ,  $172^{\circ}11'E$ , 102 m (1v, M.34534);  $34^{\circ}11'S$ ,  $172^{\circ}10'E$ , alive, 91 m (1v, M.33801); S of Great Island, alive,  $34^{\circ}14.1'S$ ,  $172^{\circ}09.0'E$ , 192-202 m (2pr, 3v, M.149580);  $34^{\circ}20.2'S$ ,  $172^{\circ}21.8'E$ , 121 m (1v, M.149578); 28 km S of Great Island,  $34^{\circ}24.0'S$ ,  $172^{\circ}16.8'E$ , 120 m (2v, M.149584). NW of Cape Reinga,  $34^{\circ}14.2'S$ ,  $172^{\circ}32.4'E$ , 100 m (1v, M.94571);  $34^{\circ}22.8'S$ ,  $172^{\circ}24.6'E$ , alive, 121 m (5pr, 4v, M.149585). N of Cape Reinga,  $34^{\circ}14'S$ ,  $172^{\circ}37'E$ , alive, 88 m (1pr, 2v, M.36002).

*Other material examined.* Three Kings Trough, NW of Three Kings Islands, 34°00'S, 171°55'E, 805 m (1v, M.149809).

#### Description

Shell up to 6.10 mm long, ovate–trigonal, of moderate thickness, moderately inflated, approximately  $1.7 \times$  longer than high; anterior end slightly longer than posterior, fresh specimens translucent white with irregular, opaque white radial rays. Prodissoconch approximately 220 µm wide, prominent, convex, smooth; PI approximately 200 µm wide, globular. Dissoconch anterior margin well rounded, smoothly merging with straight anterodorsal and broadly rounded ventral margin; posterodorsal margin straight; posterior truncation more or less flat, slightly oblique, corners tightly rounded. Right valve with two strong, well-defined anterior and posterior hinge teeth, length of inner dorsal margins each with concave shelf to seat corresponding elevated narrow dorsal margins of left valve. Left

valve with prominent, posteriorly inclined boss at posterior end of anterodorsal margin. Lithodesma stout (width 570  $\mu$ m in adult), about as long as broad, rounded anteriorly, concave posteriorly; ligaments thin, sunken resilifers shallowly concave. Interior with faint radial lines, pallial line and large adductor scars well defined; pallial sinus broad and shallow, angulate dorsally. Exterior convex, posterior area weakly convex, separated by low, broadly rounded angulation; crisply sculptured throughout with crowded, columnar, calcareous granules set in thin periostracum, stronger on posterior area.

#### Distribution

Off Three Kings Islands, 88–805 m; taken alive at 88–202 m from comminuted bryozoan/shell substrata (Fig. 15*A*).

# Remarks

*Plectodon regalis* differs from the allopatric species *P. lepidus* in having considerably finer granulate sculpture, in being slightly more elongate and in that the posterior truncation is slightly shorter dorsoventrally. It differs from the sympatric species *P. pruinosus* in being less inflated and having flat, instead of broadly curved, dorsal margins, taller sculptural granules and a weaker swelling on the central area of the prodissoconch.

# Etymology

From the Latin *regalis* (royal).

# Plectodon sp.

# Material examined

N of Norfolk Island, 28°59'S, 167°58'E, 38 m (1v, M.225171).

# Distribution

Off Norfolk Island, 38 m (1v only).

# Remarks

A single left valve (length 3.95 mm) from off Norfolk Island has flat dorsal margins, as in *P. lepidus* and *P. regalis*, but has finer sculpture than *P. lepidus*, coarser sculpture that *P. regalis* and is more inequilateral than both. It clearly represents a distinct, undescribed species, although I am reluctant to base a new taxon on a single valve, particularly a left one.

# Genus Rhinoclama Dall & E. A. Smith in Dall, 1886

Rhinoclama Dall & Smith in Dall, 1886: 300. Type species (ICZN Opinion 1376 (1986)): Cuspidaria (Rhinoclama) adamsi Morgan & Heppell in Allen & Morgan, 1981; Recent, Philippines.

# Rhinoclama (s. str.) aupouria (Dell, 1950) n. comb.

# Figs 10G, 14B,C, 15B, 16B,D, 21N

*Cuspidaria aupouria* Dell, 1950: 21, figs 3, 4; Powell, 1979: 436 (in part = three undescribed species of *Cuspidaria* + *Plectodon regalis*).

Cuspidaria (Rhinoclama) aupouria. - Poutiers & Bernard, 1995: 156.

NOT *Cuspidaria aupouria*. – Crozier, 1966: 46, figs 16, 17 (= three undescribed species of *Cuspidaria* + *Plectodon regalis*).

# Material examined

*Holotype.* N of North Cape, New Zealand, 34°18′S, 173°02′E, 137 m, J. A. Bollons, SS *Hinemoa* (rv, M.4726).

*Other material examined.* Off Three Kings Islands: 34°00'S, 171°55'E, 805 m (3v, M.144430); submarine cave, S side Rosemary Rock, 34°10.8'S, 172°03.00'E, 20 m (1v, M.117166); 11 km NW of Great Island, 34°06.5'S, 172°04.7'E, 310 m (1v, M.149587); off West Island, *Elingamite* wreck, 34°11'S, 172°03'E, 37 m (1v, M.154154); 34°14.8'S, 172°13.6'E, alive, 173–178 m (7pr, 4v, M.144427); 39 km SW of Great Island, 34°17.6S, 171°45.3'E, 427 m (4v, M.94252); 28 km S of Great Island, 34°24.0'S, 172°16.8'E, 120 m (2v, M.144428). Off Mayor Island, 37°16.7'S, 176°17.5'E, 104–109 m (1v, M.144429). Ranfurly Bank, East Cape: 37°32.8'S, 178°48.7'E, alive, 94 m (3pr, 24v, M.144433); 37°33.1'S, 178°49.5'E, 94–89 m (3v, M.144431); 37°33.2'S, 178°50.3'E, 76–71 m (2v, M.71542); 37°33.4'S, 178°48.3'E, alive, 106–103 m (2pr, 7v, M.71792); 37°37.8'S, 178°52.4'E, 50–72 m (1v, M.96179); 37°38.4'S, 178°51.7'E, 79–83 m (5pr, 3v, M.96349).

# Description

Shell up to 5.40 mm long, ovate-trigonal, stout, inflated, approximately 1.4× longer than high, anterior end 37%–42% of shell length, antero- and posterodorsal margins of right valve overlapping those of left; translucent white with narrow, opaque white radial rays; periostracum thin, buff. Prodissoconch approximately 230 µm wide, subcircular, very finely maleate, wrinkled, PI approximately 200 µm wide, globular. Dissoconch anterior margin well rounded, smoothly merging with flat or very weakly convex anterodorsal margin and broadly convex ventral margin; posterodorsal margin more or less straight; posterior truncation weakly convex, oblique, corners tightly rounded. Right valve with two strong, well-defined anterior and posterior hinge teeth, posterior tooth considerably larger, length of inner dorsal margins shelved, strongly so posteriorly, to seat corresponding elevated narrow dorsal margins of left valve. Left valve edentulous. Resilifers sunken, narrowly ovate concave, lithodesma trigonal. Pallial line and large adductor scars well defined; pallial sinus broad, concave. Exterior convex, posterior area strongly flattened and bounded by tightly rounded angulation; posterodorsal margin of both valves margined by sharp angulation; anterodorsal margin bounded by rounded angulation. Early dissoconch covered with minute punctae; thereafter flattened rostral area with coarse, crowded granules, elsewhere matt.

# Distribution

Off Three Kings Islands, Mayor Island and Ranfurly Bank, East Cape, 20–805 m; taken alive at 71–178 m from comminuted bryozoan/shell substrata (Fig. 15*B*).

#### Remarks

*Cuspidaria aupouria* Dell, 1950, resembles the type species of *Austroneaera* and congeners from the New Zealand region in gross shell morphology, but differs significantly by having much stronger, more crowded granules on the rostral area, a distinct rostral angulation, microscopic punctae on the early dissoconch, a distinctly thicker shell, angulations bounding both the antero- and posterodorsal margins, a pattern of opaque white rays and a stronger area on the right valve. Shell thickness and rostral sculpture approach the conditions in New Zealand *Plectodon* species, all of which have, however, strong exterior sculpture throughout and a prominent boss at the posterior end of the anterodorsal margin. Because *C. aupouria* clearly does not belong in *Rhinoclama (Austroneaera*), it is tentatively referred to *Rhinoclama (s. str.*), although the rostrum is shorter and the commarginal sculpture considerably weaker than in the type species.

The six valves identified as *Cuspidaria aupouria* by Crozier (1966; off Three Kings Islands, 805 m) prove to comprise a mixture of three undescribed species of *Cuspidaria* (M.20826, 149807, 149808) and a specimen of *Plectodon regalis* n. sp. (M.149809).

# Subgenus Austroneaera Powell, 1937

Austroneaera Powell, 1937a: 174. Type species (by original designation): Austroneaera brevirostris Powell, 1937; Recent, New Zealand.

#### Remarks

Allen and Morgan (1981) interpreted *Austroneaera* as a synonym of *Rhinoclama* because of similarity between their type species in gross shell morphology. They are also similar in the presence of a lithodesma set on sunken resilifers, although Morgan and Heppell (in Allen and Morgan 1981) failed to mention these characteristics in their description of the holotype of *R. adamsi*. However, the type species of *Austroneaera* differs in that the posterior area is defined by a low, rounded angulation rather than an elevated, well-defined ridge and in lacking prominent commarginal ridges (misleadingly described as 'striations' by Morgan and Heppell (in Allen and Morgan 1981) in their description of *R. adamsi*). Regrettably, the holotype of *R. adamsi* has been severely damaged by Byne's disease, so it is impossible to tell whether it has the minute, thinly distributed, calcareous periostracal granules characteristic of the type species of *Austroneaera* and most other congeners from the New Zealand region. Whatever the case, relatively coarse, crowded granules, such as those characteristic of *Plectodon* species, are certainly absent.

Poutiers and Bernard (1995) interpreted *Austroneaera* as a synonym of *Rhinoclama*, which they grouped as a subgenus of *Cuspidaria* Nardo, 1840, stating (p. 148) that the '... general arrangement of the genus depends on the type and presence of hinge teeth, which seem rather mutable'. I prefer a more conservative approach and follow Krylova (1994), who interpreted *Austroneaera* as a subgenus of *Rhinoclama*.

#### Rhinoclama (Austroneaera) brevirostris Powell, 1937

# Figs 3N, 10B, 13A,C, 15C, 21L

Austroneaera brevirostris Powell, 1937a: 174, pl. 48, fig. 11; Powell, 1979: 438, pl. 79, fig. 18. *Cuspidaria (Rhinoclama) brevirostris.* – Poutiers & Bernard, 1995: 157.

#### Material examined

Holotype. 34°13.3'S, 172°12.0'E, off Three Kings Islands, New Zealand, 260 m, 17 Aug. 1932, RRS Discovery II (1v, BMNH 1962950).

*Other material examined.* Off Three Kings Islands: Three Kings Trough, 34°00'S, 171°55'E, 805 m (1v, M.17522); Middlesex Bank, 34°00.9'S, 171°44.7'E, alive, 201–216 m (1pr, 10v, M.93234); Middlesex Bank, 34°01.2'S, 171°44.4'E, alive, 206–211 m (1pr, 4v, M.144420); Middlesex Bank, 34°02.0'S, 171°44.0'E, 246–291 m (11v, M.144421); 11 km NW of Great Island, 34°06.5'S, 172°04.7'E, alive, 310 m (5v, 1pr, M.93861); S of Great Island, 34°14.1'S, 172°09.0'E, 192–202 m (2v, M.149801); SE of Great Island, 34°14.8'S, 172°13.6'E, alive, 173–178 m (6pr, 16v, M.144426); 37 km SW of Great Island, 34°20.4'S, 171°48.2'E, 440 m (1v, M.144423). Ranfurly Bank, East Cape: 37°32.8'S, 178°48.7'E, alive, 94 m (5pr, 11v, M.60800); 37°33.1'S, 178°49.5'E, alive, 94–89 m (10pr, 20v, M.74703); 37°33.2'S, 178°50.3'E, alive, 76–71 m (1pr, 1v, M.70542); 37°33.4'S, 178°48.3'E, alive, 106–103 m (9pr, 64v, M.70791). Rungapapa Knoll, WNW of White Island, 37°33.8'S, 176°59.0'E, alive, 188–228 m (13pr, 9v, M.113550). Bradshaw Basin floor, Bradshaw Sound, 45°17.3'S, 167°02.6'E, 415 m (1pr, M.138711). Utah Basin floor, Doubtful Sound, 45°17.9'S, 166°55.5'E, 400 m (1v, M.144425).



**Fig. 13.** Shells of *Rhinoclama (Austroneaera)* species. *A,C, Rhinoclama (Austroneaera) brevirostris* Powell, 1937: left (*A* upper) and right valves, length 3.70 mm, SE of Great Island, 173–178 m (M.144426). *B,D, Rhinoclama (Austroneaera) finlayi* Powell, 1937: left (*B* upper) and right valves, length 2.90 mm, N of Poor Knights Islands, 146 m (M.144424). *E,G, Rhinoclama (Austroneaera) raoulensis* Powell, 1958: left (*G* upper) and right valves, length 3.80 mm, between Dayrell and Chanter Islands, 31–45 m (M.226999). *F,H, Rhinoclama (Austroneaera) brooki* n. sp.: holotype, left (*H* upper) and right valve, length 5.00 mm, SE of Nugent Island, Raoul Island, Kermadec Islands, 165–146 m (M.152685).

#### Description

Shell up to 4.40 mm long, ovate–trigonal, thin, inflated,  $1.33-1.43 \times$  longer than high, translucent white, beaks distinctly opisthogyrous, anterior end 47%–51% of shell length, antero- and posterodorsal margins of right valve overlapping those of left. Prodissoconch approximately 200 µm wide, subcircular, surface exceedingly finely maleate, PI approximately 180 µm wide, globular. Dissoconch anterior margin well rounded, smoothly

merging with flat or very weakly convex anterodorsal margin and broadly convex ventral margin; posterodorsal margin more or less straight; posterior truncation weakly convex, slightly oblique, corners tightly rounded. Right valve with two strong, well-defined anterior and posterior hinge teeth, posterior tooth considerably larger, length of inner dorsal margins weakly shelved to seat corresponding elevated narrow dorsal margins of left valve. Left valve edentulous. Lithodesma stout (width 400 µm in adult), about as long as wide, subtriangular, concave posteriorly; ligaments thin, narrowly ovate sunken resilifers concave. Interior with weak radial grooves, pallial line and large adductor scars well defined; pallial sinus broad, shallow, concave. Exterior convex, posterior area flattened and separated by broadly rounded angulation, posterodorsal margin of both valves bounded by sharp angulation; sculptured with fine commarginal growth lines and extremely small, thinly distributed calcareous granules set in thin, shiny, pale buff periostracum.

# Distribution

Three Kings Islands, NE North Island, and Fiordland, 71–805 m; taken alive at 71–310 m from comminuted bryozoan/shell substrata (Fig. 15*C*).

#### Remarks

Three specimens from Fiordland resemble R. (A.) brevirostris rather than R. (A.) finlayi in shape. Assuming they are R. (A.) brevirostris, the lack of intermediate populations would suggest a relictual distribution.

#### Rhinoclama (Austroneaera) finlayi Powell, 1937

Figs 10C, 13B,D, 15D

Austroneaera finlayi Powell, 1937a: 175, pl. 48, fig. 12; Powell, 1979: 438, pl. 79, fig. 17. Cuspidaria (Rhinoclama) finlayi. – Poutiers & Bernard, 1995: 159. NOT Austroneaera finlayi. – Dell, 1956: 168 (= Cuspidaria (Pseudoneaera) wellmani (Fleming, 1948)).

# Material examined

Holotype. Off Poor Knights Islands, New Zealand, 110 m (1v, AIM AK70071).

*Other material examined.* N of North Cape, 34°20.0'S, 173°06.6'E, 163–168 m (1v, M.144422). Off Poor Knights Islands: 35°22'S, 174°43'E, alive, 146 m (6pr, 20v, M.144424); 35°29'S, 174°43.5'E, 110 m (7v, M.67870). Off Mayor Island, 37°15.2'S, 176°14.5'E, 188–193 m (1v, M.60990); 37°16.7'S, 176°17.5'E, alive, 104–109 m (4pr, 12v, M.67497). Off White Island, 37°31'S, 177°08'E, 213 m (1pr, 2v, M.10363).

# Description

Shell up to 3.50 mm long, ovate-trigonal, thin, inflated,  $1.20-1.32 \times$  longer than high, translucent white, beaks distinctly opisthogyrous, anterior end 42%-43% of shell length, antero- and posterodorsal margins of right valve overlapping those of left. Prodissoconch approximately 200 µm wide, subcircular, smooth, PII not apparent. Dissoconch anterior margin well rounded, smoothly merging with flat or very weakly convex anterodorsal margin; posterodorsal margin more or less straight; posterior truncation weakly convex, slightly oblique, corners tightly rounded. Right valve with two strong, well-defined anterior and posterior hinge teeth, posterior tooth considerably larger, length of inner dorsal margins weakly shelved to seat corresponding elevated narrow dorsal margins of left valve. Left valve edentulous. Lithodesma stout, narrowly ovate sunken resilifers concave. Interior with weak radial grooves, pallial line and large adductor scars well defined; pallial sinus broad, shallow, concave. Exterior convex,



**Fig. 14.** Dissoconch microsculpture of Cuspidariidae and Spheniopsidae (exterior, right valve and adult unless indicated). *A*, *Rhinoclama (Austroneaera) brooki* n. sp., holotype, SE of Nugent Island, Raoul Island, Kermadec Islands, 165–146 m (M.152685). *B*,*C*, *Rhinoclama (s. str.) aupouria* (Dell, 1950), prodissoconch and early dissoconch (*B*, boundary arrowed) and adult posterodorsal area (*C*), SE of Great Island, Three Kings Islands, 173–178 m (M.144427). *D*, *Grippina aupouria* (Powell, 1937), view showing free edge of periostracum folded around ventral margin onto interior surface of adult right valve (demonstrating that calcareous beads are periostracal), Middlesex Bank, 201–216 m (M.93236). *E*, *Grippina rex* n. sp., holotype, off Three Kings Islands, 91 m (M.33807). *F*, *Grippina punctata* n. sp., holotype, off Spirits Bay, 63 m (M.152679). *G*, *Grippina spirata* n. sp., holotype, off Spirits Bay, 63 m (M.149519). *H*, *Grippina acherontis* n. sp., holotype, off Patea, 40 m (M.152680). *I*, *Grippina punila* n. sp., holotype, off Puysegur Point, 183 m (M.152677). Scale bars: *B*, 25 μm; others, 50 μm.

posterior area flattened, posterodorsal margin of both valves bounded by sharp angulation; sculptured with fine commarginal growth lines and minute, thinly distributed calcareous granules set in thin, shiny, pale buff periostracum.

# Distribution

Off North Cape and Poor Knights, Mayor and White Islands, North-eastern North Island, 110–213 m; taken alive at 104–146 m from muddy substrata (Fig. 15*D*).

# Remarks

Compared with *R*. (*A*.) *brevirostris*, *R*. (*A*.) *finlayi* differs in being more narrowly trigonal (length/width ratio right valves 1.20-1.32, mean 1.259, SD 0.037, n = 10: as against 1.33-1.48, mean 1.418, SD 0.043, n = 10; and 1.32-1.43, mean 1.371, SD 0.040, n = 10 for *R*. (*A*.) *brevirostris* from off the Three Kings Islands and Ranfurly Bank, East Cape respectively), in attaining a smaller size (length up to 3.5 v. 4.40 mm), in that the shelf inside the posterodorsal margin of the right valve extends further posteriorly and in that the shell wall is distinctly thicker. Approximately 10% of specimens of *R*. (*A*.) *brevirostris* are as narrowly trigonal as *R*. (*A*.) *finlayi*, although all seen are larger than the few broadly trigonal

specimens of R. (A.) finlayi. Most specimens of R. (A.) finlayi and are also more inflated that R. (A.) brevirostris and have flatter area on rostrum. Rhinoclama (A.) finlayi has its distributional range within that of R. (A.) brevirostris and lives in muddy substrata rather than clean bryozoan/shell hash. Rhinoclama (A.) finlayi may ultimately prove to be an ecophenotypic variant of R. (A.) brevirostris, although supportive comparative anatomical data are wanting.

Juveniles of R. (A.) finlayi and R. (A.) brevirostris are superficially similar to Cuspidaria (Pseudoneaera) wellmani (Fleming, 1948), but are immediately separable by the sunken resilifers (in hinge line in *P. wellmani*), the broader lithodesma and the stronger hinge teeth. Specimens of *C. wellmani* from off the Chatham Islands (M.10849–51) were misidentified as *R*. (A.) finlayi by Dell (1956), the latter species being unknown from the Chatham Islands.

#### Rhinoclama (Austroneaera) raoulensis Powell, 1958

Figs 30, 10E, 13E–G, 15E, 21M

Austroneaera raoulensis Powell, 1958: 78, pl. 9, fig. 9. Cuspidaria (Rhinoclama) raoulensis. – Poutiers & Bernard, 1995: 165.

#### Material examined

Holotype. 29°15'S, 177°51'W, off Raoul Island, Kermadec Islands, New Zealand, 75–85 m, RV Galathea (rv, Zoological Museum, Copenhagen).

*Other material examined.* Off Meyer Island, Raoul Island, Kermadec Islands, 29°14.7'S, 177°52.7'W, alive, 27–22 m (3pr, 1v, M.225814). Between Dayrell and Chanter Islets, Raoul Island, 29°15.00'S, 177°50.90'W, alive, 31–45 m (1pr, M.226999).

# Description

Shell up to 3.80 mm beaks distinctly opisthogyrous, anterior end 45%-52% of shell length, antero- and posterodorsal margins of right valve overlapping those of left. Prodissoconch approximately 170 µm wide, subcircular, smooth, long, elongate-trigonal, thin, inflated, 1.54-1.64× longer than high, translucent white, PII not apparent. Dissoconch anterior margin well rounded, smoothly merging with flat or very weakly convex anterodorsal margin and broadly convex ventral margin; posterodorsal margin more or less straight; posterior truncation weakly convex, slightly oblique, corners tightly rounded. Right valve with two strong, well-defined anterior and posterior hinge teeth, posterior tooth considerably larger, length of inner dorsal margins weakly shelved to seat corresponding elevated narrow dorsal margins of left valve. Left valve edentulous. Lithodesma stout (width 300 µm in adult), ligaments thin, narrowly ovate sunken resilifers concave. Pallial line and large adductor scars well defined; pallial sinus broad, shallow. Exterior convex, posterior area flattened, posterodorsal margin of both valves bounded by low, rounded angulation; sculptured with fine commarginal growth lines and extremely small, thinly distributed calcareous granules set in thin, shiny, translucent, pale buff periostracum.

#### Distribution

Raoul Island, Kermadec Islands, 22–85 m; taken alive at 22–45 m from clean basaltic sand (Fig. 15*E*).



Fig. 15. Maps of the New Zealand region showing distributions of *Plectodon*, *Rhinoclama* (*s. str.*) and *Rhinoclama* (*Austroneaera*) species (200 and 1000 m isobaths indicated). *A*, *Plectodon regalis* n. sp. *B*, *Rhinoclama* (*s. str.*) *aupouria* (Dell, 1950). *C*, *Rhinoclama* (*Austroneara*) *brevirostris* Powell, 1937. *D*, *Rhinoclama* (*Austroneara*) *finlayi* Powell, 1937. *E*, *Rhinoclama* (*Austroneara*) *raoulensis* Powell, 1958. *F*, *Rhinoclama* (*Austroneara*) *brooki* n. sp.

# Remarks

*Rhinoclama (Austroneaera) raoulensis* differs from *R. (A.) brevirostris* and *R. (A.) finlayi* principally in being shorter dorsoventrally and in having a less sharply defined angulation outside the posterodorsal margin. The gut of one specimen contained an ostracod and remains of two cumaceans (det. W. R. Webber, NMNZ). A second species from off Raoul Island is described below.

# Rhinoclama (Austroneaera) brooki n. sp.

Figs 10D, 13F,H, 14A, 15F

*Austroneaera raoulensis.* – Brook & Marshall in Brook, 1998: 214 (in part not Powell, 1958). *Cuspidaria (Rhinoclama)* sp. 1 Spencer *et al.* (in press).

#### Material examined

Holotype. SE of Nugent Island, Raoul Island, Kermadec Islands, New Zealand, 29°14.7'S, 177°49.4'W, 165–146 m, 28 Oct. 1975, RV Acheron (pr, M.152685).

*Paratypes.* Kermadec Islands. SE of Nugent Island, Raoul Island, 29°14.7'S, 177°49.4'W, 165–146 m, 28 Oct. 1975, RV *Acheron* (4v, M.225684). Off Raoul Island: NW of Fleetwood Bluff, 29°12.7'S, 177°56.1'W, 135 m (2v, M.225440); 29°13.0'S, 177°59.8'W, 201–146 m (1v, M.226915); E of Dayrell Islet, 29°14.73'S, 177°50.34'W, 135–146 m (1v, M.226672); SE of D'Arcy Point, 29°18.8'S, 177°54.2'W, 274–219 m (1v, M.226815); SE of Smith Bluff, 29°18.90'S, 177°56.40'W, 82–100 m (1v, M.227116).

#### Description

Shell up to 6.60 mm long, ovate-trigonal, thin, inflated, approximately 1.3× longer than high, translucent white, beaks distinctly opisthogyrous, anterior end 45%-50% of shell length, antero- and posterodorsal margins of right valve overlapping those of left. Prodissoconch approximately 170 µm wide, low, convex, smooth, no PII. Dissoconch anterior margin well rounded, smoothly merging with flat or very weakly convex anterodorsal margin and broadly convex ventral margin; posterodorsal margin more or less straight; posterior truncation weakly convex, oblique, corners tightly rounded. Right valve with two strong, well-defined anterior and posterior hinge teeth, posterior tooth considerably larger, length of inner dorsal margins weakly shelved to seat corresponding elevated narrow dorsal margins of left valve. Left valve edentulous. Resilifers sunken, narrowly ovate concave (lithodesma almost certainly present although absent from single valves). Interior with weak radial grooves, pallial line and large adductor scars well defined; pallial sinus broad, shallow, concave. Exterior convex, posterior area weakly flattened; posterodorsal margin of both valves margined by sharp angulation; sculptured with fine commarginal growth lines and extremely small, thinly distributed calcareous granules set in thin periostracum.

#### Distribution

Off Raoul Island, Kermadec Islands, 82–274 m (shells only; Fig. 15F).

#### Remarks

*Rhinoclama (Austroneaera) brooki* differs from *R. (A.) raoulensis* principally in attaining nearly twice the size (length up to 6.60 v. 3.80 mm), in having a more strongly convex ventral margin and in having a sharp exterior angulation bordering the posterodorsal margin. *Rhinoclama (A.) brooki* ranges deeper than *A. raoulensis (*82–174 v. 22–85 m). Specimens of *R. (A.) raoulensis* from 75–85 m, the deepest record for the species, were isolated valves (taken living at 22–45 m) and whether the two species have overlapping bathymetric ranges in life (syntopic?) remains to be established.

# Etymology

After F. J. Brook (Whangarei), in appreciation of his work on Recent and fossil Mollusca of the Kermadec Islands (Brook 1998).

# Rhinoclama (Austroneaera) tangaroa n. sp.

Figs 10F, 16A, C, 18A

#### Material examined

*Holotype.* Wanganella Bank, Norfolk Ridge, E slope, 32°35.3′S, 167°41.8′E, alive, 437–422 m, 29 Jan. 1981, RV *Tangaroa* (live taken: a right valve, left valve crushed; M.273140).

Paratypes. Same data as holotype (5v, M.234116).

Other material examined. SSE of Philip Island, Norfolk Island, 29°33'S, 168°07'E, 143 m (8v, M.225295); Middlesex Bank, NW of Three Kings Islands, 34°02.0'S, 171°44.0'E, 246–291 m (2v, M.151544).

#### Description

Shell up to 2.52 mm long, ovate-trigonal, beaks slightly anterior, thin and brittle, moderately inflated,  $1.35 \times$  longer than high (holotype right valve), right valve slightly higher than left, fresh specimens strongly translucent apart from an opaque white band on interior against anterodorsal and posterodorsal margins. Prodissoconch approximately 200 µm wide, low, convex, smooth apart from fine commarginal wrinkles. Dissoconch anterior margin well rounded, smoothly merging with straight anterodorsal margin and broadly convex ventral margin; posterodorsal margin very broadly convex; posterior truncation weakly convex, more or less vertical, corners tightly rounded. Right valve with two well-defined anterior and posterior hinge teeth, posterior tooth larger, most of length of inner dorsal margins weakly shelved to seat corresponding elevated narrow dorsal margins of left valve. Left valve edentulous. Lithodesma stout (width 200 µm in holotype), about as long as wide, broadly rounded anteriorly, concave posteriorly; ligaments thin, narrowly ovate sunken resilifers concave. Pallial line and large adductor scars well defined; pallial sinus broadly concave. Exterior convex, posterior area flattened; sculptured with fine commarginal growth lines. Periostracum translucent, glossy, microscopic granules not evident.

# Distribution

Norfolk Island, Wanganella Bank, and Middlesex Bank, 143–437 m; taken alive at 422–437 m (Fig. 18*A*).

# Remarks

*Rhinoclama (Austroneaera) tangaroa* is extremely similar to *R. (A.) finlayi* in shape, but differs in attaining slightly smaller size (length up to 2.52 v. 3.50 mm), in lacking granular microsculpture and in having opaque white lines on the interior against the dorsal margins. The two well-preserved valves from Middlesex Bank (evidently a single individual) are indistinguishable from the Norfolk Ridge specimens.

# Etymology

After RV Tangaroa.

#### Family SPHENIOPSIDAE Gardner, 1928

# Diagnosis

Shell small (length up to 4 mm) subtrigonal, inequilateral or almost equilateral, equivalve, posterior end modestly produced or truncate, beaks median or posterior. Right valve with



**Fig. 16.** Shells of *Rhinoclama (Austroneaera), Rhinoclama (s. str.)* and *Grippina* species. *A*,*C*, *Rhinoclama (Austroneaera) tangaroa* n. sp.: holotype, right valve (*A* lower, *C*), length 2.10 mm (M.273140), and paratype, left valve, length 2.20 mm (M.234116), Wanganella Bank, Norfolk Ridge, 437–422 m (M.234116). *B*,*D*, *Rhinoclama (s. str.) aupouria* (Dell, 1950): left (*B* upper) and right valve, length 4.50 mm, right valve (*D*) length 4.40 mm, SE of Great Island, Three Kings Islands, 173–178 m (M.144427). *E*,*G*, *Grippina aupouria* (Powell, 1937): subadult, left (*G* upper), length 2.70 mm, 11 km NW of Great Island, Three Kings Islands, 310 m (M.93848). *F*,*H*, *Grippina aupouria*: adult, left (*H* upper) and right valve, length 3.50 mm, right valve (*H* lower) length 3.30 mm, Middlesex Bank, 201–216 m (M.93236).

anterior and posterior teeth, left valve without teeth; right valve teeth and left valve hinge line separated by wide gap occupied by lithodesma. Left valve without teeth. Pallial sinus shallow. Exterior with or without weak or strong commarginal ridges, some species with addition of minute punctae. Periostracum in some species set with minute calcareous granules. Dimyarian. Mantle margins fused, small pedal gape, two short conjoined siphons.

# Genus Grippina Dall, 1912

*Grippina* Dall, 1912: 128. Type species (by original designation): *Grippina californica* Dall, 1912; Recent, California.

# Remarks

Keen (1969*a*) referred *Grippina* to Spheniopsidae, which, until now, has been grouped in Heterodonta near Corbiculidae (Gardner 1928; Keen 1969*a*; Coan 1990*b*; Cosel 1995; Coan *et al.* 2000). However, *Grippina* species have sunken resilifers and a lithodesma and some have exterior microsculpture extremely similar to that in some thraciids and cuspidariids, so it seems likely that the Spheniopsidae really belongs in Anomalodesmata (there seems no reason to doubt that *Grippina* and *Spheniopsis* are confamilial). Because shell shape and (when present) commarginal sculpture both suggest affinities with Cuspidariidae, I have tentatively grouped Spheniopsidae in Cuspidarioidea rather than Thracioidea pending knowledge of the anatomy.

Apart from some species of *Spheniopsis* Sandberger, 1861 from the Oligocene of Germany (Neuffer 1973), *S. americana* Dall, 1903 (Early Miocene, Florida) and the *Grippina* species reported below, the only other known spheniopsids are the Recent species *G. californica* Dall, 1912 (California), *S. frankbernardi* Coan, 1990 (Baja California and Costa Rica), *S. senegalensis* Cosel, 1995 (West Africa; Cosel 1995) and the three or four western Atlantic *Spheniopsis* species reported to E. V. Coan by the late D. R. Moore (Coan 1990b), one of which is *S. triquetra* (Verrill & Bush, 1898) (North Carolina). An undescribed *Spheniopsis* species is well represented in collections from bioclastic substrata off northern New Caledonia and Lifou, Loyalty Islands (MNHN). All the New Zealand *Grippina* species have extremely narrow geographic ranges, occur very patchily and most are rare relative to the volumes of sediment processed to recover them, which suggests they have unusually narrow ecological requirements. Spheniopsid diet is unknown, but if they are, indeed, cuspidariids, they are probably carnivores.

Coan (1990*b*) reported brooding in *G. californica*, the type species of the genus. The animal of an unspecified *Grippina* species was briefly described by Coan *et al.* (2000: 481): '... Dimyarian ... Mantle margins fused, with small pedal gape and two short conjoined siphons'.

# Grippina aupouria (Powell, 1937)

Figs 10H, 14D, 16E–H, 18B, 21O

*Mysella aupouria* Powell, 1937*a*: 172, pl. 47, fig. 5; Powell, 1979: 395, fig. 98/4. *Grippina aupouria*. – Coan, 1990*b*: 400.

#### Material examined

*Syntypes.* Off Three Kings Islands, New Zealand, 34°13.3'S, 172°12.0'E, 260 m, 17 Aug. 1932, RRS *Discovery II* (left and right v, BMNH 1962944).

*Other material examined.* Off Three Kings Islands: King Bank, 33°57.4'S, 172°19.4'E, 128–123 m (4v, M.148017); 18 km N of Great Island, 33°59.2'S, 172°13.6'E, 155 m (1v, M.93508); Three Kings Trough, 34°00'S, 171°55'E, 805 m (16v, M.17493); Middlesex Bank, 34°00.9'S, 171°44.7'E, 201–216 m (3pr, 4v, M.93236); 34°01'S, 172°07'E, 622 m (3v, M.34959); Middlesex Bank, 34°01.2'S, 171°44.4'E, 206–211 m (1v, M.148108); 13 km N of Great Island, 34°01.8'S, 172°12.0'E, 508 m (2v, M.149721); Middlesex Bank, 34°02.0'S, 171°44.0'E, alive, 246–291 m (7pr, 23v, M.148023); Middlesex Bank, 34°02.1'S, 171°45.8'E, 221–206 m (1pr, 2v, M.148021); 24 km NW of Great Island, 34°05.9'S, 171°55.1'E, alive, 710 m (1pr, 5v, M.94378); 11 km NW of Great Island, 34°06.5'S, 172°04.7'E, alive, 310 m (3pr, 8v, M.93848); off North-east Island, 34°08.5'S, 172°11'E, 102 m (2pr, 3v, M.34536); 34°10'S, 172°12'E, 252

m (1v, M.34199); S of Great Island, 34°14.1'S, 172°09.0'E, alive, 192–202 m (1pr, 15v, M.148019); 39 km SW of Great Island, 34°17.6'S, 171°45.3'E, 427 m (1v, M.94288). NW of Cape Reinga, 34°14.2'S, 172°32.4'E, 100 m (2pr, M.94554).

# Description

Shell up to 4.15 mm long, obliquely ovate-trigonal, of moderate thickness, rather strongly inflated, large individuals up to  $1.26 \times$  longer than high, with increasing shell size anterior end elongating anteroventrally more rapidly than posterior so that beak position shifts from an initial position slightly behind midlength to approximately posterior one-third (increasingly inequilateral); fresh specimens semitranslucent white with small, sparse, irregular, cuneiform, opaque white radial streaks. Prodissoconch approximately 270 µm wide, prominent, globular, smooth; PI approximately 230 µm wide. Dissoconch anterior margin well rounded, smoothly merging with more or less straight anterodorsal margin and broadly rounded ventral margin, posterodorsal margin broadly rounded; posterior truncation short, straight, oblique, extremities roundly angulate. Right valve with two strong, welldefined hinge teeth that gently converge inwards relative to valve margins, separated by very broad space occupied by lithodesma, anterior tooth longer, inner anterodorsal margin shelved to seat corresponding elevated narrow margin of left valve. Left valve with small rounded process on anterior corner of lithodesma space and shallow groove posteriorly that seats dorsal part of posterior right valve tooth, lithodesma space oblique and narrower than in right valve. Lithodesma stout, anteriorly tapered, subtrigonal in dorsal view, ligaments thin, sunken resilifers shallowly concave, 500 µm wide in a shell 3.20 mm long. Interior glossy apart from opaque white patches near anterodorsal and posterodorsal margins; ventrally enlarging groove extending gently posteriorly from below umbones, ending as an oblique pit; pallial line and large adductor scars well defined; pallial sinus shallow, concave. Exterior convex, posterior end flattened, sculptured with low rounded commarginal ridges; and a crisp microsculpture of crowded, flat-topped, calcareous periostracal granules, some of which are connected in groups of two to four that are aligned commarginally.

# Distribution

Off Three Kings Islands, 98–805 m; taken alive at 98–710 m from comminuted bryozoan/shell substrata. Restricted endemic (Fig. 18*B*).

#### Remarks

Originally described in the montacutid genus *Mysella* Angas, 1877, this species was referred to *Grippina* by Coan (1990b) because of similarity to the type species in shape, size, thickness, sculpture, hinge morphology, pallial line and muscle scars. Six additional New Zealand *Grippina* species are described below.

# Grippina rex n. sp.

# Figs 10*I*, 14*E*, 17*A*,*C*, 18*C*

Grippina sp. 7 Spencer et al. (in press).

# Material examined

Holotype. Off Three Kings Islands, New Zealand, 34°11′S, 172°10′E, alive, 91 m, 19 Feb. 1974, RV Acheron (pr, M.33807).

*Paratypes.* Off Three Kings Islands: Middlesex Bank, 33°57.0′S, 171°45.4′E, alive, 98–103 m (3pr, M.148022); King Bank, 33°57.4′S, 172°19.4′E, 128–123 m (2v, M.149514); Three Kings Trough, 34°00′S,

171°55′E, 805 m (5v, M.149518); Middlesex Bank, 34°01.2′S, 171°44.4′E, 206–211 m (1v, M.149517); Middlesex Bank, 34°02.0′S, 171°44.0′E, 246–291 m (9v, M.149512); 11 km NW of Great Island, 34°06.5′S, 172°04.7′E, 310 m (1v, M.149516); off North-east Island, 34°08.5′S, 172°11′E, 102 m (2v, M.149513).

# Description

Shell up to 5.10 mm long, obliquely ovate-trigonal, of moderate thickness, rather strongly inflated, large individuals up to 1.44× longer than high, with increasing shell size anterior end elongating anteroventrally slightly more rapidly than posterior so that beak position shifts from an initial position in front of posterior one-third to behind it; fresh specimens semitranslucent white with small, irregular, cuneiform, opaque white radial streaks. Prodissoconch approximately 230 µm wide, prominent, smooth; PI approximately 200 µm wide, globular. Dissoconch anterior margin well rounded, smoothly merging with more or less straight anterodorsal margin and broadly rounded ventral margin, posterodorsal margin broadly rounded or almost straight; posterior truncation short, straight, oblique, extremities roundly angulate. Right valve with two strong, well defined hinge teeth that converge inwards relative to margins, posterior tooth at a markedly higher angle than anterior; separated by very broad space occupied by lithodesma, anterior tooth longer, inner anterodorsal margin shelved to seat corresponding elevated narrow dorsal margin of left valve. Left valve with increasingly prominent rounded process on anterior corner of lithodesma space and concave space posteriorly that seats posterior right valve tooth, lithodesma space oblique and narrower than in right valve. Lithodesma stout (width 570 µm in adult), anteriorly tapered, subtrigonal in dorsal view, ligaments thin, sunken resilifers shallowly concave. Interior glossy; ventrally enlarging groove extending gently posteriorly from below umbones, ending as an oblique pit; pallial line and large adductor scars well defined, pallial sinus shallow, concave. Exterior convex, posterior end flattened; with a crisp microsculpture of crowded, flat-topped, calcareous periostracal granules, many of which are grouped in pairs aligned commarginally.

# Distribution

Off Three Kings Islands, 91–805 m; taken alive at 91–103 m from comminuted bryozoan/shell substrata. Restricted endemic (Fig. 18*C*).

# Remarks

Compared with the sympatric species *G. aupouria*, which has a similar pattern of white radial streaks, *G. rex* differs in attaining larger size, in that the anterior end increases in length more slowly relative to the posterior with increasing shell size and the posterior tooth on the left valve is inclined at a steeper angle to the valve margin. *Grippina rex* differs further in having a stronger process on the left valve hinge, stronger microsculpture and in lacking commarginal ridges.

# Etymology

From the Latin rex (king).

# Grippina punctata n. sp.

Figs 10J, 14F, 17B,D, 18D

Grippina sp. 2 Spencer et al. (in press).

#### Material examined

*Holotype.* Off Spirits Bay, New Zealand, 34°19.98'S, 172°45.79'E, 63 m, 25 Jan. 1999, RV *Kaharoa* (pr, M.152679).

*Paratypes.* Off Three Kings Islands: 37 km NE of Great Island, 33°58.0'S, 172°30.6'E, 550 m (4v, M.149724); Three Kings Trough, 34°00'S, 171°55'E, 805 m (18v, M.149521); 22 km ENE of Great Island, 34°05.0'S, 172°24.6'E, 200 m (1pr, M.149601); outer South-east Bay, Great Island, 34°10'S, 172°08'E, 55 m (4v, M.33965); S of Great Island, 34°14.1'S, 172°09.0E, alive, 192–202 m (1pr, M.149520). Off Spirits Bay, 34°18.36'S, 172°49.39'E, alive, 68 m (3pr, many v, M.149508); 34°19.98'S, 172°45.79'E, 63 m (20v, M.149509); 34°21.60'S, 172°43.19'E, alive, 48 m (2pr, 3v, M. 153097). NW of Cape Maria van Diemen, 34°20.0'S, 172°30.0'E, 100 m (1v, M.93740).

*Other material examined.* Off Spirits Bay, 34°21.15′S, 172°45.92′E, 54 m (7v, M.154153). NNE of Cape Reinga, 34°22.47′S, 172°42.08′E, 53 m (1pr, 1v, M.152031).

#### Description

Shell up to 2.15 mm long, obliquely ovate-trigonal, thick to very thick, strong, moderately inflated, up to 1.32× longer than high, posterior end occupying approximately 36% of shell length, fresh specimens semitranslucent white. Prodissoconch approximately 220 µm wide, convex, topographically complex, minutely maleate and wrinkled, PI approximately 200 µm wide. Dissoconch anterior margin well rounded, smoothly merging with broadly rounded or almost flat anterodorsal margin and broadly rounded ventral margin, posterodorsal margin broadly rounded or straight; posterior truncation short, typically broadly concave, straight in few specimens, oblique, extremities angulate. Right valve with two strong, well-defined hinge teeth, both gently converging inwards relative to valve margins or posterior tooth more or less parallel to margin, separated by very broad space occupied by lithodesma (present but lost), inner anterodorsal margin shelved almost to anterior extremity to seat corresponding elevated narrow dorsal margin of left valve. Left valve with indentations that seat dorsal parts right valve teeth, lithodesma space smaller than in right valve. Resilifers sunken, narrowly ovate, oblique. Interior glossy, shallow groove extending gently posteriorly from below umbones, pallial line and large adductor scars well defined; pallial sinus shallow, concave. Exterior convex, posterior end flattened, valves up to approximately 700 µm length sculptured with deep circular pits arranged commarginally, thereafter with low, broad, rounded commarginal ridges and a crisp microsculpture of densely crowded, flat-topped, calcareous periostracal granules that are grouped in more or less commarginal rows.

# Distribution

Off Three Kings Islands and of northern tip of Northland, 53–805 m; taken alive at 48–202 m from comminuted bryozoan/shell and sand substratum (Fig. 18*D*).

# Remarks

*Grippina punctata* differs from *G. aupouria* and *G. rex* principally in attaining smaller size, in being much thicker, in having a projecting boss on the prodissoconch, in having a



**Fig. 17.** Shells of *Grippina* species. *A*,*C*, *Grippina rex* n. sp., holotype, left (*A* upper) and right valve, length 4.10 mm, off Three Kings Islands, 91 m (M.33807). *B*,*D*, *Grippina punctata* n. sp.: holotype, left (*B* upper) and right valve, length 1.45 mm, off Spirits Bay, 63 m (M.152679). *E*,*G*, *Grippina globosa* n. sp.: holotype, right valve (*E*,*G* lower), length 1.30 mm (M.152672), and paratype, left valve (M.149490), Middlesex Bank, NW of Three Kings Islands, 246–291 m. *F*,*H*, *Grippina spirata* n. sp.: holotype, left (*F* upper) and right valve, length 2.10 mm, off Spirits Bay, 63 m (M.149519).

(typically) concave instead of straight posterior truncation, in having pits on the early dissoconch and in lacking opaque white radial streaks.

# Etymology

From the Latin *punctatus* (pricked with small holes).

# Grippina globosa n. sp.

Figs 10*K*,*L*, 17*E*,*G*, 18*E*, 21*P* 

Grippina sp. 3 Spencer et al. (in press).

# Material examined

*Holotype.* Middlesex Bank, NW of Three Kings Islands, New Zealand, 34°02.0'S, 171°44.0'E, alive, 246–291 m, 31 Jan. 1981, RV *Tangaroa* (1v, M.152672).

*Paratypes.* 37 km NE of Great Island, Three Kings Islands, 33°58.0'S, 172°30.6'E, 550 m (1v, M.149723). Middlesex Bank, NW of Three Kings Islands, 34°02.0'S, 171°44.0'E, alive, 246–291 m (2pr, 3v, M.149490). 37 km NW of Great Island, 34°02.0'S, 171°48.4'E, 188 m (1pr, M.150078). NW of Cape Reinga, 34°14.2'S, 172°32.4'E, 100 m (1pr, M.94549).

# Description

Shell up to 1.43 mm long, obliquely ovate-trigonal, of moderate thickness, strongly inflated, up to 1.18× longer than high, posterior end occupying approximately 38% of shell length, fresh specimens semitranslucent white. Prodissoconch approximately 400 µm wide, broadly convex, smooth, PI approximately 200 µm wide. Dissoconch anterior margin well rounded, smoothly merging with broadly rounded anterodorsal and ventral margins, posterodorsal margin almost straight; posterior truncation short, very weakly convex, oblique, extremities roundly angulate. Right valve with two strong, well-defined hinge teeth that are separated by a broad space occupied by lithodesma and small boss in front of posterior tooth; inner anterodorsal margin shelved to seat corresponding elevated narrow dorsal margin of left valve. Left valve with rather deep indentations that seat dorsal sides of right valve teeth, depth of indentations effectively forming small anterior and posterior teeth bordering space occupied by lithodesma. Lithodesma stout (width 170 µm in adult holotype), about as long as wide, anteriorly tapered, subtrigonal in dorsal view, ligaments thin, sunken resilifers shallowly concave. Interior glossy, shallow groove extending gently posteriorly from below umbones, pallial line and large adductor scars well defined; pallial sinus shallow, concave. Exterior convex, posterior end flattened, sculptured throughout with fine commarginal growth lines; minute, crowded pits on early dissoconch, no calcareous periostracal granules.

# Distribution

Off Three Kings Islands, 100–550 m from comminuted bryozoan/shell substratum; taken alive at 246–291 m (Fig. 18*E*).

# Remarks

*Grippina globosa* is much smaller than previously described *Grippina* species and is further distinguished by the depressed, relatively large prodissoconch, the strongly convex dissoconch, the short anterior end and the presence of a boss in front of the posterior tooth on the right valve.

*Grippina* sp. 5 of Spencer *et al.* (in press) from off the Three Kings Islands at 805 m, is very similar to *G. globosa* in shape, but larger and with a considerably broader prodissoconch. Regrettably, the only known specimen, a single valve, was lost during preparation for scanning electron microscopy.

# Etymology

From the Latin globosus (spherical).



**Fig. 18.** Maps of the New Zealand region showing distributions of *Rhinoclama (Austroneaera)* and *Grippina* species (200 and 1000 m isobaths indicated). *A*, *Rhinoclama (Austroneaera) tangaroa* n. sp. *B*, *Grippina aupouria* (Powell, 1937). *C*, *Grippina rex* n. sp. *D*, *Grippina punctata* n. sp. *E*, *Grippina globosa* n. sp. *F*, *Grippina spirata* n. sp.

# Grippina spirata n. sp.

# Figs 10M, 14G, 17F,H, 18F

Grippina sp. 6 Spencer et al. (in press).

# Material examined

*Holotype.* Off Spirits Bay, New Zealand, 34°19.98'S, 172°45.79'E, alive, 63 m, 25 Jan. 1999, RV *Kaharoa* (pr, M.149519).

*Paratypes.* Three Kings Islands, 22 km ENE of Great Island, 34°05.0′S, 172°24.6′E, 200 m (1pr, M.149602). Off Spirits Bay, 34°18.36′S, 172°49.39′E, 68 m (4v, M.149522).

#### Description

Shell up to 2.05 mm long, obliquely ovate-trigonal, of moderate thickness, strong, moderately inflated, up to  $1.46 \times$  longer than high, posterior end occupying approximately



**Fig. 19.** Shells of *Grippina* species. *A,C, Grippina acherontis* n. sp.: holotype, left (*A* upper) and right valve, length 2.20 mm, off Patea, 40 m (M.152680). *B,D, Grippina pumila* n. sp.: holotype, left (*B* upper) and right valve, length 1.45 mm, off Puysegur Point, 183 m (M.152677).

34% of shell length, fresh specimens semitranslucent white. Prodissoconch approximately 230 µm wide, prominent, globose, smooth, PI approximately 200 µm wide. Dissoconch anterior margin well rounded, smoothly merging with very broadly rounded anterodorsal and ventral margins, posterodorsal margin almost straight; posterior truncation short, oblique, concave, extremities angulate. Right valve with two strong, well-defined hinge teeth, anterior tooth longer, separated by very broad space occupied by lithodesma, inner anterodorsal margin shelved to seat corresponding elevated narrow dorsal margin of left valve. Left valve with shallow posterior indentation that seats dorsal part of right valve posterior tooth, lithodesma space smaller than in right valve. Lithodesma stout (lost), subrectangular, ligaments thin, sunken resilifers shallowly concave. Interior glossy, smooth, pallial line and large adductor scars well defined; ventrally enlarging groove extending gently posteriorly from below umbones, ending as an oblique pit; pallial sinus shallow, broadly concave. Exterior convex, posterior end shallowly concave, sculptured with low, rounded commarginal ridges, and a microsculpture of densely crowded, flat-topped, mostly elongate, calcareous periostracal granules that are aligned commarginally.

# Distribution

East of Three Kings Islands and off Spirits Bay, 63–200 m; taken alive at 63 m from comminuted shell and sand substratum (Fig. 18*F*).

# Remarks

*Grippina spirata* resembles *G. punctata* in gross facies, but differs markedly in having a smoothly convex prodissoconch and in lacking punctations on the early dissoconch. The two species are sympatric. *Grippina spirata* differs principally from *G. aupouria* and *G. rex* in attaining smaller size, in lacking white radial streaks and in details of exterior microsculpture and differs from *G. globosa* in details of shape, size, hinge morphology and sculpture.

# Etymology

From the type locality, Spirits Bay.

# Grippina acherontis n. sp.

Figs 10N, 14H, 19A,C, 20A, 21Q

Grippina sp. 1 Spencer et al. (in press).

# Material examined

*Holotype.* S of Patea, New Zealand, 39°56'S, 174°26'E, alive, 40 m, 2 Mar. 1976, RV *Acheron* (pr, M.152680).

Paratypes. Same data as holotype (22pr, M.53594).

# Description

Shell up to 2.45 mm long, obliquely ovate-trigonal, of moderate thickness, moderately inflated, up to 1.32× longer than high, anterior end becoming somewhat longer relative to anterior with increasing shell size, posterior end occupying up to 35% of shell length in adults, fresh specimens semitranslucent white. Prodissoconch approximately 270 µm wide, prominent, broadly convex, smooth, PI approximately 230 µm wide. Dissoconch anterior margin well rounded, smoothly merging with more or less straight anterodorsal margin and broadly rounded ventral margin, posterodorsal margin very broadly rounded; posterior truncation short, straight, oblique, extremities roundly angulate. Right valve with two strong, well-defined hinge teeth, anterior tooth longer, posterior tooth more prominent, separated by very broad space occupied by lithodesma, inner anterodorsal margin shelved to seat corresponding elevated narrow dorsal margin of left valve. Left valve with small boss at posterior end of elevated dorsal margin, lithodesma space smaller than in right valve. Lithodesma stout (width 320 µm in adult), subrectangular in dorsal view, ligaments thin, sunken resilifers shallowly concave. Interior glossy, smooth, pallial line and large adductor scars well defined, pallial sinus shallow, concave. Exterior convex, posterior end flattened, with low, very weak, rounded commarginal ridges; and a crisp microsculpture of crowded, flat-topped, calcareous periostracal granules that are grouped in commarginal rows.

# Distribution

Off Patea, south-western North Island, 40 m, living in clean sand (Fig. 20A).

#### Remarks

*Grippina acherontis* most closely resembles *G. aupouria* in gross facies, but differs in attaining smaller size, in lacking opaque white radial streaks and in that the beaks in adults are more posterior than in subadults of *G. aupouria* of the same size.



**Fig. 20.** Maps of the New Zealand region showing distributions of *Grippina* species (200 and 1000 m isobaths indicated). *A*, *Grippina acherontis* n. sp. *B*, *Grippina pumila* n. sp.

# Etymology

After RV Acheron from which the type material was dredged.

# Grippina pumila n. sp.

Figs 10O, 14I, 19B,D, 20B, 21R

Grippina sp. 4 Spencer et al. (in press).

# Material examined

*Holotype.* Off Puysegur Point, New Zealand, 46°11′S, 166°30′E, 183 m, Dec. 1908, SS *Rakiura* (pr, M.152677).

*Paratypes.* Off Puysegur Point, New Zealand,  $46^{\circ}11'S$ ,  $166^{\circ}30'E$ , 183 m, Dec. 1908, SS *Rakiura* (16v, M.143883). Off Poutama Island, Big South Cape Island, Stewart Island,  $47^{\circ}16'S$ ,  $167^{\circ}23'E$ , 55 m (1pr, 36v, M.63606).

#### Description

Shell up to 2.80 mm long, obliquely ovate-trigonal, of moderate thickness, moderately inflated, approximately 1.2× longer than high; umbones at approximately mid-shell length in specimens less than approximately 1.60 mm long, anterior end elongating to varying degrees in larger specimens so that umbones are at approximately the posterior one-third of shell length in adults, fresh specimens semitranslucent white. Prodissoconch approximately 270 µm wide, prominent, broadly convex, topographically complex, PI approximately 230 µm wide. Dissoconch anterior margin well rounded, smoothly merging with more or less straight anterodorsal margin and broadly rounded ventral margin, posterodorsal margin broadly rounded; posterior truncation short, straight, oblique, extremities roundly angulate. Right valve with two strong, well-defined hinge teeth, anterior tooth longer, posterior tooth more prominent, separated by very broad space occupied by lithodesma, inner anterodorsal margin shelved to seat corresponding elevated narrow dorsal margin of left valve. Left valve with small boss at posterior end of elevated anterodorsal margin (pseudocardinal), lithodesma space smaller than in right valve. Lithodesma stout (width 170 µm in holotype), about as long as broad, ligaments thin, sunken resilifers shallowly concave. Interior glossy, smooth, pallial line and large adductor scars well defined, pallial sinus shallow, concave. Exterior convex, posterior area flattened. Sculptured with low, rounded irregular commarginal ridges and incised growth lines; no calcareous periostracal granules.



Fig. 21. Lithodesmas of Thraciidae, Myochamidae, Cuspidariidae and Spheniopsidae. A, Thracia (Odoncineta) vitrea (Hutton, 1873), width 1.00 mm, ex shell 15.0 mm long, Manaroa Bay, Pelorus Sound, 5-18 m (M.144402). B, Asthenothaerus maxwelli n. sp., width 6.45 mm, ex holotype 30.5 mm long, Horseshoe Bay, Stewart Island (M.152678). C, Barythaerus biconvexus (Powell, 1927), width 0.23 mm, ex shell 2.00 mm long, off Poutama Island, Big South Cape Island, Stewart Island, 55 m (M.19878). D, Barythaerus cuneatus (Powell, 1937), width 0.48 mm, ex shell 3.20 mm long, off Spirits Bay, 68 m (M.149504). E, Parvithracia (s. str.) suteri (Finlay, 1927), width 0.40 mm, ex shell 3.42 mm long, North Arm, Port Pegasus, Stewart Island, 37-44 m (M.44790). F, Parvithracia (s. str.) ampla n. sp., width 0.90 mm, ex holotype 6.60 mm long, Saunders Canyon, off Otago Peninsula, 457 m (M.9125). G, Parvithracia (Pseudoasthenothaerus) fragilissima n. sp., width 0.87 mm (face down), ex holotype 6.70 mm long, Challenger Plateau, 1005-1009 m (NZOI H-808). H, Hunkydora novozelandica (Reeve, 1859), width 1.65 mm, ex shell 11.2 mm long, off point SW of Munro Bay, Whangarei Harbour, 7-10 m (M.152673). I, Pseudogrippina wanganellica n. sp., width 0.23 mm ex holotype 2.30 mm long, Wanganella Bank summit, Norfolk Ridge, 113 m (M.152681). J, Plectodon lepidus n. sp., width 0.50 mm, ex holotype 5.25 mm long, off E side of Mayor Island, 59-74 m (M.152682). K, Plectodon regalis n. sp., width 0.57 mm, ex holotype 5.20 mm long, off Three Kings Islands, 91 m (M.152684). L, Austroneaera brevirostris Powell, 1937, width 0.40 mm, ex shell 3.70 mm long, SE of Great Island, 173-178 m (M.144426). M, Austroneaera raoulensis Powell, 1958, width 0.30 mm, ex shell 3.20 mm long, off Meyer Island, Raoul Island, Kermadec Islands, 27-22 m (M.225814). N, Rhinoclama aupouria (Dell, 1950), width 0.48 mm, ex shell 4.40 mm long, SE of Great Island, Three Kings Islands, 173-178 m (M.144427). O, Grippina aupouria (Powell, 1937), width 0.50 mm, ex shell 3.20 mm long, Middlesex Bank, NW of Three Kings Islands, 246-291 m (M.148023). P, Grippina globosa n. sp., width 0.17 mm, ex holotype 1.30 mm long, Middlesex Bank, 246-291 m (M.152672). Q, Grippina acherontis n. sp., width 0.32 mm, ex holotype 2.20 mm long, off Patea, 40 m (M.152680). R, Grippina pumila n. sp. width 0.17 mm, ex holotype 1.45 mm long, off Puysegur Point, 183 m (M.152677).

# Distribution

Off Puysegur Point, south-western South Island, and southern Stewart Island, 55–183 m; taken alive at 183 m from comminuted bryozoan/shell substratum (Fig. 20*B*).

# Remarks

*Grippina pumila* resembles *G. punctata* in having a topographically complex prodissoconch, but differs from that and all other *Grippina* species from the New Zealand region other than *G. globosa* in lacking granulate microsculpture. *Grippina pumila* differs from the allopatric species *G. globosa* in attaining larger size and in various details of shape and sculpture.

# Etymology

From the Latin *pumilus* (small).

# Acknowledgments

I am grateful to M. G. Harasewych (National Museum of Natural History, Washington D.C.), V. Héros (Muséum National d'Histoire Naturelle, Paris) and J. Pickering and K. M. Way (The Natural History Museum, London) for loan of type material and to T. Landers (Auckland Institute and Museum, Auckland), I. Loch (Australian Museum, Sydney), and J. H. McLean (Los Angeles County Museum of Natural History, Los Angeles) for loan of additional specimens. Special thanks to to R. Coory and N. Heke (Museum of New Zealand Te Papa Tongarewa, Wellington) for assistance with digital technology (notably the distribution maps) and photography, respectively, to J. Patterson and W. Wallace (Victoria University of Wellington) and K. Card (Industrial Research Limited, Lower Hutt) for assistance with scanning electron microscopy.

#### References

- Allen, J. A., and Morgan, R. E. (1981). The functional morphology of Atlantic deep water species of the families Cuspidariidae and Poromyidae (Bivalvia): an analysis of the evolution of the septibranch condition. *Philosophical Transactions of the Royal Society of London* B294, 413–546.
- Bernard, F. R. (1989). Living Periplomatidae of the Pacific and Indo-Pacific region. Venus 48, 1-11.
- Beu, A. G., and Maxwell, P. A. (1990). Cenozoic Mollusca of New Zealand. New Zealand Geological Survey Paleontological Bulletin 58, 1–518.
- Blainville, H. M. D. (1824). 'Dictionaire des Sciences Naturelles, Dans Lequel on Trait Méthodiquement des Différens Êtres de la Nature, Considérés Soit en eux-Mêmes, d'Après l'État Actuel de nos Connoissances, Soit Relativement a l'Utilité qu'en Peuvent Retirer la Médecine, l'Agriculture, le Commerce et les Arts.' Vol. 32. (Levrault et Le Normant: Strasbourg and Paris.)
- Blainville, H. M. D. (1825–27). 'Manuel de Malacologie et de Conchyliologie.' (pp. 1–647. [1825].) 'Nouvelles Additions et Corrections au Genera.' (pp. 649–664. [1827].) (Levrault: Strasbourg.)
- Boreham, A. (1959). Biological type specimens in the New Zealand Geological Survey. 1. Recent Mollusca. *New Zealand Geological Survey Paleontological Bulletin* **30**, 1–87.
- Branch, M. L., Arnaud, P. M., Cantera, J., and Gianakouras, D. (1991). The benthic Mollusca and Brachiopoda of subantarctic Marion and Prince Edward Islands: (1) illustrated keys to the species; (2) records of the 1982–1989 University of Cape Town surveys. *South African Journal of Antarctic Research* 21, 45–64.
- Bronn, H. G. (1862). 'Die Klassen und Ordnungen des Thier-Reichs wissenschaftlich dargestellt in Wort und Bild.' 3 (1). Malacozoa Acephala. (Leipzig.)
- Brook, F. J. (1998). The coastal molluscan fauna of northern Kermadec Islands, southwest Pacific Ocean. *Journal of the Royal Society of New Zealand* 28, 185–233.
- Brook, F. J., and Marshall, B. A. (1998). Checklist of benthic coastal marine Mollusca from the northern Kermadec Islands. Appendix. pp. 210–233. In Brook, F. J. 'The Coastal Molluscan Fauna of Northern Kermadec Islands, Southwest Pacific Ocean'. *Journal of the Royal Society of New Zealand* 28, 185–233.

#### Some Recent Anomalodesmata from New Zealand

- Carpenter, P. P. (1864). Supplementary report on the present state of our knowledge with regard to the Mollusca of the west coast of North America. *Report of the British Association for the Advancement of Science* 1863, 517–686.
- Cattaneo-Vietti, R., Chiantore, M., Schiaparelli, S., and Albertelli, G. (2000). Shallow and deep-water mollusc distribution at Terra Nova Bay (Ross Sea, Antarctica). *Polar Biology* **23**, 173–182.
- Coan, E. V. (1990*a*). The Recent eastern Pacific species of the bivalve family Thraciidae. *The Veliger* **33**, 20–55.
- Coan, E. V. (1990b). The eastern Pacific species of the bivalve family Spheniopsidae. *The Veliger* 33, 394–401.
- Coan, E. V., Valentich Scott, P., and Bernard, F. R. (2000). Bivalve seashells of western North America. Marine bivalve mollusks from Arctic Alaska to Baja California. Santa Barbara Museum of Natural History Monograph 2 Studies in Biodiversity 2, 1–764.
- Cosel, R. (1995). Fifty-one new species of marine bivalves from tropical West Africa. Iberus 13, 1-115.
- Costa, O. G. (1829). 'Catalogo Sistematico e Regionato de'Testacei delle Due Sicilie.' (Minerva: Napoli.) Cotton, B. C. (1961). 'South Australian Mollusca. Pelecypoda.' (Government Printer: Adelaide.)
- Cotton, B. C., and Godfrey, F. K. (1938). 'The molluscs of South Australia. Part 1: Pelecypoda.' (Government Printer: Adelaide.)
- Cox, L. R. (1969). General features of Bivalvia. In 'Treatise on Invertebrate Paleontology. Part N2. Mollusca 6. Bivalvia'. (ed. R. C. Moore.) pp. 2–129. (Geological Society of America and University of Kansas Press: Lawrence.)
- Crozier, M. A. (1966). New species and records of Mollusca from off the Three Kings Islands. *Transactions of the Royal Society of New Zealand*, Zoology 8, 39–49.
- Dall, W. H. (1886). Report on the Mollusca. Part 1. Brachiopoda and Pelecypoda. Reports on the results of dredging, under the supervision of Alexander Agassiz, in the Gulf of Mexico (1877–78) and in the Caribbean Sea (1879–80), by the U.S. Coast Survey Steamer 'Blake', Lieut.-Commander C. D. Sigsbee, U.S.N., and Commander J. R. Bartlett, U.S.N., commanding. 6. *Bulletin of the Museum of Comparative Zoology, Harvard* 12, 171–318.
- Dall, W. H. (1890). Preliminary report on the collection of Mollusca and Brachiopoda obtained in 1887–'88. Scientific results of explorations by the US Fish Commission Steamer Albatross. *Proceedings of the United States National Museum* **12**, 219–362.
- Dall, W. H. (1903). Contributions to the Tertiary fauna of Florida with special reference to the Silex beds of Tampa and the Pliocene beds of the Caloosahatchie River, including in many cases a complete revision of the generic groups treated and their American Tertiary species. 6. *Transactions of the Wagner Free Institute of Science* 3, 1219–1654.
- Dall, W. H. (1912). New Californian Mollusca. The Nautilus 25, 127-129.
- Dell, R. K. (1950). Notes on the taxonomy and distribution of some New Zealand Mollusca with descriptions of four new species. *Dominion Museum Records in Zoology* **3**, 21–28.
- Dell, R. K. (1956). The archibenthal Mollusca of New Zealand. Dominion Museum Bulletin 18. 1-235.
- Dell, R. K. (1990). Antarctic Mollusca with special reference to the fauna of the Ross Sea. *Royal Society* of New Zealand Bulletin **27**, 1–311.
- Dell, R. K. (1995). New species and records of deep-water Mollusca from off New Zealand. Tuhinga: Records of the Museum of New Zealand Te Papa Tongarewa 2, 1–26.
- Finlay, H. J. (1926). A further commentary on New Zealand molluscan systematics. Transactions of the New Zealand Institute 57, 320–485.
- Finlay, H. J. (1927). New specific names for austral Mollusca. *Transactions of the New Zealand Institute* **57**, 488–533.
- Fleming, C. A. (1948). New species and genera of marine Mollusca from the Southland fiords. *Transactions of the Royal Society of New Zealand* **77**, 72–92.
- Fleming, C. A. (1951). Some Australasian Mollusca in the British Museum (Natural History). *Transactions of the Royal Society of New Zealand* **79**, 126–139.
- Gardner, J. (1928). The molluscan fauna of the Alum Bluff Group of Florida. 5. Tellinacea, Solenacea, Mactracea, Myacea, Molluscoidea. *United States Geological Survey Professional Paper* **142E**, 185–249.
- Gardner, N. W. (1976). *Myochama tasmanica* Ten. Woods from Northland. *Poirieria. Conchology Section* Auckland Institute and Museum **8**, 40–41.
- Habe, T. (1960). New species of molluscs from the Amakusa Marine Biological Laboratory, Reihoku-cho, Amakusa, Kumamoto Pref., Japan. Publications of the Seto Marine Biological Laboratory 8, 289–298.

Hain, S. (1990). The benthic seashells (Gastropoda and Bivalvia) of the Weddel Sea, Antarctica. Berichte zur Polarforschung 70, 1–180.

- Harper, E. M., and Morton, B. (2000). The biology and functional morphology of *Myochama anomioides* Stutchbury, 1830 (Bivalvia: Anomalodesmata: Pandoroidea), with reference to cementation. *Journal of Molluscan Studies* 66, 403–416.
- Harper, E. M., Hide, E. A., and Morton, B. (2000). Relationships between extant Anomalodesmata: a cladistic test. In 'The Evolutionary Biology of the Bivalvia'. (Eds E. M. Harper, J. D. Taylor and J. A. Crame.) pp. 129–143. (Geological Society of London, Special Publication: London.)
- Hedley, C. (1901). Some new or unfigured Australian shells. Records of the Australian Museum 4, 22-27.
- Higo, S., Callomon, P., and Goto, Y. (2001). 'Catalogue and Bibliography of the Marine Shell-Bearing Mollusca of Japan. Type figures.' (Elle: Osaka.)
- Hutton, F. W. (1873a). 'Catalogue of the Marine Mollusca of New Zealand, With Diagnoses of the Species.' (Didsbury, Government Printer: Wellington.)
- Hutton, F. W. (1873b). 'Catalogue of the Tertiary Mollusca and Echinodermata of New Zealand, in the Collection of the Colonial Museum.' (Didsbury, Government Printer: Wellington.)
- Hutton, F. W. (1878). Révision des Coquilles de la Novelle-Zélande et des Îles Chatham. Journal de Conchyliologie 26, 5–57.
- Hutton, F. W. (1880). 'Manual of the New Zealand Mollusca.' (Hughes, Government Printer: Wellington.)
- Hutton, F. W. (1884). Revision of the Recent Lamellibranchiata of New Zealand. *Proceedings of the Linnean Society of New South Wales* 9, 512–533.
- Hutton, F.W. (1893). The Pliocene Mollusca of New Zealand. In 'The MacLeay Memorial Volume.' (Ed. J. J. Fletcher.) pp. 35–92. (Linnean Society of New South Wales: Sydney.)
- ICZN (1986). Opinion 1376. Cuspidaria (Rhinoclama) adamsi Morgan & Heppell, 1981 designated as type species of Rhinoclama Dall & Smith, 1886 (Mollusca, Bivalvia). Bulletin of Zoological Nomenclature 43, 30–32.
- Iredale, T. (1930). More notes on the marine Mollusca of New South Wales. Records of the Australian Museum 17, 384–407.
- Kamanev, G. M. (2002). Genus Parvithracia (Bivalvia: Thraciidae) with descriptions of a new subgenus and two new species from the northwestern Pacific. Malacologia 44, 107–134.
- Kamenev, G. M., and Nadtochy, V. A. (1998). Two new species of *Lampeia* (Bivalvia: Thraciidae) from the northwestern Pacific, with notes on *Lampeia adamsi* (MacGinitie, 1959). *The Veliger* 41, 259–273.
- Keen, A. M. (1969*a*). Superfamily Myacea. In 'Treatise on Invertebrate Paleontology. Part N2. Mollusca 6. Bivalvia'. (Ed. R. C. Moore.) pp. 690–699. (Geological Society of America and University of Kansas Press: Lawrence.)
- Keen, A. M. (1969b). Superfamily Pandoracea. In 'Treatise on Invertebrate Paleontology. Part N2. Mollusca 6. Bivalvia'. (Ed. R. C. Moore.) pp. 843–852. (Geological Society of America and University of Kansas Press: Lawrence.)
- Krylova, E. M. (1994). Bivalve molluses of the genus *Rhinoclama* (Septibranchia, Cuspidariidae) from seamounts. *Transactions of the P. P. Shirshov Institute of Oceanology* **129**, 55–64.
- Krylova, E. M. (2001). Septibranchiate molluscs of the family Poromyidae (Bivalvia: Poromyoidea) from the tropical western Pacific Ocean. In 'Tropical Deep-Sea Benthos', 22. (Eds P. Bouchet and B. A. Marshall.) Mémoires du Muséum National d'Histoire Naturelle 185, 165–200.
- Lamprell, K., and Healy, J. (1998). 'Bivalves of Australia.' Vol. 2. (Backhuys: Leiden.)
- Linnaeus, C. (1758). 'Systema naturae per regna tria naturae, secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis, etc.' 1. Editio decima, reformata. (Salvii: Holmiae.)
- Linse, K. (1997). Distribution of epibenthic Mollusca from the Chilean Beagle Channel. Berichte zur Polarforschung 228, 1–131.
- Linse, K., and Brandt, A. (1998). Distribution of epibenthic Mollusca on a transect through the Beagle Channel (southern Chile). *Journal of the Marine Biological Association of the United Kingdom* **78**, 875–889.
- Macpherson, J. H. (1958). 'An illustrated index of Tasmanian shells,' revised edition. (Shea: Tasmania.)
- Marshall, B. A. (1998). Marine Mollusca from Wellington Harbour and approaches. Appendix 5. In 'Te Whanganui a Tara Wellington Harbour. Review of Scientific and Technical Studies of Wellington Harbour, New Zealand, to 1997'. (Ed. J. Heine.) pp. 145–155. (East Harbour Association: Wellington.)
- Martens, E. (1873). 'Critical list of the Mollusca of New Zealand Contained in European Collections, with References to Descriptions and Synonyms.' (Hughes: Wellington.)

Martens, E. (1874). 'Errata and Addenda.' (Colonial Museum: Wellington.)

May, W. L. (1921). 'A Checklist of the Mollusca of Tasmania.' (Government Printer: Hobart.)

- May, W. L. (1923). 'An Illustrated Index of the Mollusca of Tasmania.' (Government Printer: Hobart.)
- Morton, B. (1977). Some aspects of the biology and functional morphology of *Myadora striata* (Quoy and Gaimard) (Bivalvia: Anomalodesmata: Pandoracea). *Journal of Molluscan Studies* **43**, 141–154.
- Morton, B. (1980). Some aspects of the biology and functional morphology (including the presence of a ligamental lithodesma) of *Montacutona compacta* and *M. olivacea* (Bivalvia: Leptonacea) associated

with coelenterates in Hong Kong. Journal of the Zoological Society of London 192, 431-455.

Morton, B. (1981). The Anomalodesmata. *Malacologia* **21**, 35–60.

- Neuffer, F. O. (1973). Die Bivalven des Unteren Meeressandes (Rupelium) im Mainzer Becken. Abhandlungen der Hessischen Landesamtes für Bodenforschung (Wiesbaden) 68, 1–113.
- Ockelmann, K. W. (1965). Redescription, distribution, biology, and dimorphous sperm of *Montacuta tenella* Lovén (Mollusca, Leptonacea). *Ophelia* 2, 211–222.

Odhner, N. H. (1924). New Zealand Mollusca. Papers from Dr. Th. Mortensen's Pacific Expedition 1914–16. Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening; Kjobenhavn 77, 1–90.

Palmer, K. V. W. (1958). Type specimens of marine Mollusca described by P. P. Carpenter from the West Coast (San Diego to British Columbia). *The Geological Society of America Memoir* 76, 1–376.

Petterd, W. F. (1884). Descriptions of new Tasmanian shells. Journal of Conchology 4, 135–145.

- Poutiers, J. -M., and Bernard, F. R. (1995). Carnivorous bivalve molluscs (Anomalodesmata) from the tropical western Pacific Ocean, with a proposed classification and a catalogue of Recent species. Résultats des Campagnes MUSORSTOM 14. Mémoires du Muséum National d'Histoire Naturelle, Zoologie 167, 107–187.
- Powell, A. W. B. (1927). Deep-water Mollusca from south-west Otago. *Records of the Canterbury Museum* **3**, 113–124.
- Powell, A. W. B. (1937*a*). New species of marine Mollusca from New Zealand. *Discovery Reports* 15, 153–222.

Powell, A. W. B. (1937b). 'The Shellfish of New Zealand: An Illustrated Index.' (Unity: Auckland.)

- Powell, A. W. B. (1946). 'The Shellfish of New Zealand: An Illustrated Handbook,' 2nd edn. (Whitcombe and Tombs: Christchurch.)
- Powell, A. W. B. (1957). 'Shells of New Zealand: An Illustrated Handbook,' 3rd end. (Whitcombe and Tombs: Christchurch.)
- Powell, A. W. B. (1958). Mollusca of the Kermadec Islands. Part 1. Records of the Auckland Institute and Museum 5, 65–85.
- Powell, A. W. B. (1962). 'Shells of New Zealand: An Illustrated Handbook,' 4th edn. (Whitcombe and Tombs: Christchurch.)

Powell, A. W. B. (1974). New Zealand molluscan systematics with descriptions of new species, part 8. *Records of the Auckland Institute and Museum* 11, 197–207.

- Powell, A. W. B. (1976). On the considerable influx of warm water molluscs that have invaded northern New Zealand waters within recent years. *Records of the Auckland Institute and Museum* 13, 141–166.
- Powell, A. W. B. (1979). 'New Zealand Mollusca. Marine, Land and Freshwater Shells.' (Collins: Auckland.)
- Prezant, R. S. (1998). Subclass Anomalodesmata. In 'Mollusca: The Southern Synthesis. Fauna of Australia 5. Part A'. (Eds P. L. Beesley, G. J. B. Ross and A. Wells). Chapter 9. (CSIRO Publishing: Melbourne.)
- Rafinesque, C. S. (1815). 'Analyse de la Nature ou Tableau de l'Universe et des Corps organisés, etc.' (Palermo.)

Reeve, L. (1859). 'Monograph of the genus *Thracia*. 'Conchologia iconica' 12.' (Reeve: Brook.)

- Smith, E. A. (1885). Report on the Lamellibranchiata collected by the HMS Challenger during the years 1873–76. Report on the scientific results of the voyage of HMS Challenger during the years 1873–76. *Zoology* 13, 1–341.
- Soot-Ryen, T. (1941). Northern pelecypods in the collection of Tromsø Museum. 1. Order Anomalodesmacea: families Pholadomyidae, Thraciidae and Periplomatidae. *Tromsø Museums Årshefter* **61**, 1–41.
- Spencer, H. G., Marshall, B. A., and Willan, R. C. (in press). Phylum Mollusca. In 'The New Zealand Inventory of Biodiversity: A Species 2000 Symposium Review'. (Ed. D. P. Gordon.) (Canterbury Press: Christchurch.)
- Stoliczka, F. (1870–71). 'The Pelecypoda, with a review of all known genera of this class, fossil and Recent. Cretaceous fauna of Southern India. 3. Palaeontologica Indica'. Memoirs of the Geological Survey of India. (Geological Survey of India: Calcutta.)

Stutchbury, S. (1830). On two new genera of testaceous Mollusca, and five new species of the genus *Anatina*, lately discovered at Port Jackson, New South Wales. *Zoological Journal* **5**, 95–101.

Suter, H. (1913). 'A Manual of the New Zealand Mollusca, with an Atlas of Quarto Plates.' (1915) (Government Printer: Wellington.)

Suter, H. (1914). Revision of the Tertiary Mollusca of New Zealand, based on type material. Part 1. New Zealand Geological Survey Paleontological Bulletin 2, 1–64.

Tate, R., and May, W. L. (1901). A revised census of the marine Mollusca of Tasmania. Proceedings of the Linnean Society of New South Wales 26, 344–471.

- Taylor, J. D., Kennedy, W. J., and Hall, A. (1973). The shell structure and mineralogy of the Bivalvia. 2. Lucinacea–Clavagellacea: conclusions. *Bulletin of the British Museum (Natural History) Zoology* 22, 255–294.
- Tenison Woods, J. T. (1877). On some new Tasmanian marine shells. Proceedings of the Royal Society of Tasmania 1876, 131–159.
- Thiele, J. (1912). Die antarktischen Schnecken und Muscheln. *Deutsche Südpolar-Expedition 1901–1903* 13, 183–285.
- Troncoso, N., van Goethem, J. L., and Troncoso, J. S. (2001). Contribution to the marine molluscan fauna of Kerguelen Islands, South Indian Ocean. *Iberus* **19**, 83–114.
- Yamamoto, G., and Habe, T. (1959). Fauna of shell-bearing mollusks in Mutsu Bay. Lamellibranchia (2). Bulletin of the Marine Biological Station of Asamushi, Tohoku University 9, 85–122.
- Yonge, C. M., and Morton, B. (1980). Ligament and lithodesma in the Pandoracea and the Poromyacea with a discussion on evolutionary history in the Anomalodesmata (Mollusca: Bivalvia). *Journal of the Zoological Society of London* **191**, 263–292.

http://www.publish.csiro.au/journals/mr