

# ANIMAL PRODUCTION SCIENCE

## CONTENTS Volume 56, Issue 5, 2016, 797–940

Starch utilisation in chicken-meat production: the foremost influential factors

*Ha H. Truong, Sonia Y. Liu and Peter H. Selle* 797–814

Effect of dietary ALA on growth rate, feed conversion ratio, mortality rate and breast meat omega-3 LCPUFA content in broiler chickens

*John F. Carragher, Beverly S. Mühlhäusler, Mark S. Geier, James D. House, Robert J. Hughes and Robert A. Gibson* 815–823

Increased proportion of female lambs by feeding Border Leicester × Merino ewes a diet high in omega-6 fatty acids around mating

*E. H. Clayton, J. F. Wilkins and M. A. Friend* 824–833

The effect of maternal nutrition level during mid-gestation on postnatal muscle fibre composition and meat quality in lambs

*U. Sen, E. Sirin, U. Ensoy, Y. Aksoy, Z. Ulutas and M. Kuran* 834–843

Birthweight has no influence on chemical body composition and muscle energy stores in suckling piglets

*Maartje De Vos, Veronique Huygelen, Myriam Hesta, Sofie A. Willemsen, Erik Fransen, Christophe Casteleyn, Steven Van Cruchten and Chris Van Ginneken* 844–849

Carob pulp inclusion in lamb diets: effect on intake, performance, feeding behaviour and blood metabolites

*M. N. Noor-Ehsan Gobindram, M. Bognanno, G. Luciano, M. Lanza and L. Biondi* 850–858

Bayesian analysis of direct and maternal effects for birthweight in Iranian buffaloes using Gibbs sampling

*Navid Ghavi Hossein-Zadeh* 859–865

Two-step weaning in beef calves: permanence of nose flaps for 7 or 21 days does not influence the behaviour response

*P. Alvez, G. Quintans, M. J. Hötzel and R. Ungerfeld* 866–870

Effects of increasing rates of molasses supplementation and forage quality on the productivity of steers

*R. A. Hunter and P. M. Kennedy* 871–881

Resource use and environmental impacts from beef production in eastern Australia investigated using life cycle assessment

*Stephen Wiedemann, Eugene McGahan, Caoilinn Murphy and Mingjia Yan* 882–894

Endophyte metabolites associated with perennial ryegrass toxicosis

*K. F. M. Reed, W. J. Mace, L. V. Walker and L. R. Fletcher* 895–907

Eggshell quality: a comparison between Fayoumi, Gimieizah and Brown Hy-Line strains for mechanical properties and ultrastructure of their eggshells

*Lamiaa M. Radwan* 908–912

Correlated response to selection for some egg performance traits in egg line of Japanese quail

*B. Y. F. Mahmoud, G. Abou Khadiga and E. A. El-Full* 913–917

Heat-induced apoptosis and gene expression in bovine mammary epithelial cells

*Han Hu, Jiaqi Wang, Haina Gao, Songli Li, Yangdong Zhang and Nan Zheng* 918–926

Genetic parameters for calving difficulty using complex genetic models in five beef breeds in Australia

*M. G. Jeyaruban, D. J. Johnston, B. Tier and H.-U. Graser* 927–933

Estimation of genetic parameters for lambing ease, birthweight and gestation length in Australian sheep

*L. Li and D. J. Brown* 934–940