

The Magellanic System's Interactive Formations

M. E. Putman

Research School of Astronomy & Astrophysics, Australian National University,
Private Bag, Weston Creek PO, ACT 2611, Australia
putman@mso.anu.edu.au

Received 1999 November 17, accepted 2000 January 7

Abstract: The interaction between the Galaxy and the Magellanic Clouds has resulted in several high-velocity complexes which are connected to the Clouds. The complexes are known as the Magellanic Bridge, an HI connection between the Large and Small Magellanic Clouds, the Magellanic Stream, a $10^\circ \times 100^\circ$ HI filament which trails the Clouds, and the Leading Arm, a diffuse HI filament which leads the Clouds. The mechanism responsible for these features formation remains under some debate, with the lack of detailed HI observations being one of the limiting factors in resolving the issue. Here I present several large mosaics of HI Parkes All-Sky Survey (HIPASS) data which show the full extent of the three Magellanic complexes at almost twice the resolution of previous observations. These interactive features are connected, but unique in their spatial and velocity distribution. The differences may shed light on their origin and present environment. Dense clumps of HI along the sightline to the Sculptor Group, which may or may not be associated with the Magellanic complexes, are also discussed.

Keywords: intergalactic medium — Local Group — Magellanic Clouds — galaxies: HI

1 Introduction

Almost every galaxy in the Universe is either currently, or was at some time, part of an interacting system. This includes our Galaxy, but until the recent discovery of the Sagittarius dwarf, HI features about the Magellanic Clouds were the only clear sign of our interactive past. Giant neutral hydrogen structures, known as the Magellanic Stream, the Magellanic Bridge and the Leading Arm, depict the Clouds' interaction with each other and the Galaxy. These Magellanic HI complexes make up a large fraction of the high-velocity HI concentrations in the southern sky which are known as high-velocity clouds (HVCs). High-velocity clouds are objects which do not fit into the classical picture of Galactic HI rotation and fall in the velocity range of $|v_{lsr}| = 90 - 400 \text{ km s}^{-1}$ (see Wakker & van Woerden 1997 for a review). The overall HVC population has had a number of possibilities put forth for its origin: from superbubbles in the Galactic disk, to tidal debris, to building blocks of the Local Group. To distinguish between the range of possibilities and determine if more HVCs are related to the interaction of our Galaxy with the Magellanic Clouds, we must better understand the overall HI distribution of HVCs. Maps from the HI Parkes All Sky Survey (HIPASS) have enabled

this by providing a marked improvement in spatial resolution compared to previous HVC surveys (see Putman & Gibson 1999), and revealing previously hidden structure and small high-velocity clumps about the Magellanic System.

2 Magellanic Complexes

The three HVCs which are classified as Magellanic debris are the Magellanic Bridge, the Magellanic Stream and the Leading Arm (see Putman 2000 for a review). The Magellanic Bridge (see Figure 1) is a $\sim 10^8 M_\odot$ filament of neutral hydrogen which joins the two Magellanic Clouds and has a velocity gradient which proceeds from 125 km s^{-1} at the tail of the SMC or Shapley's Wing, to $\sim 300 \text{ km s}^{-1}$ at the extended arm of the LMC. Tidal models predict that the Bridge was pulled from the SMC during a close encounter between the two Clouds 0.2 Gyr ago (Gardiner & Noguchi 1996). The HIPASS data reveal new structure along the Bridge, as well as an extension from the LMC which suggests that the Bridge is made of both SMC and LMC material. The Bridge is the only HVC known to contain stars, and the age of many of the stellar concentrations (10–25 Myr) indicates that the Bridge is a star forming region (Grondin, Demers & Kunkel 1992).




Figure 1—Neutral hydrogen column density map of the Magellanic Bridge which joins the Large and Small Magellanic Clouds. The projection was chosen based on previous maps of the system and the main features are labelled. The contours are 1, 2, 4, 8, 16 and $32 \times 10^{20} \text{ cm}^{-2}$.

The HIPASS map of the Magellanic Stream is shown in Figure 2. The Stream trails the Magellanic Clouds for over 100° and is not a confined filament as previous maps depict (Mathewson, Cleary & Murray 1974), but a complex network of filaments and clumps. The beginning of the Stream consists of multiple filaments and head-tail structures as it spews from the northern side of the SMC and Bridge at $v_{lsr} = 90 - 240 \text{ km s}^{-1}$. Most of these filaments end at declination $\sim 60^\circ$, but the main filament of the Stream continues to march northward, its bifurcated structure diminishing towards the northern tip.¹ Dense clumps follow the Stream in position and velocity, except for the region of the Sculptor Group which will be discussed in the next section. There are no stars in the Stream (e.g. Guhathakurta & Reitzel 1998), but $\text{H}\alpha$ appears to be detectable at every location along the Stream with a column density greater than 10^{19} M_\odot (Weiner & Williams 1996; Bland-Hawthorn & Maloney 1999b). These detections, together with the fact that no [OIII] has been detected from the Stream, indicates that the Stream is being ionised by photons escaping from the Galaxy (Bland-Hawthorn & Maloney 1999a).

The third Magellanic complex is a natural tidal counterpart to the Stream, and is called the Leading Arm (see Figure 3). This feature is more diffuse than the Stream and its leading position indicates that

tidal forces are the dominant mechanism responsible for forming the Magellanic Stream. This complex was only recently confidently defined as Magellanic debris (Putman et al. 1998), as the coarse resolution and observing grid used in the previous surveys missed the filaments between the clumps and the connection to the Magellanic Clouds. Further evidence that this feature is of Magellanic origin is the metallicity determination of Lu et al. (1998), using the background galaxy NGC 3783.

3 Is it all Magellanic Debris?

The ability of HIPASS to uncover the Leading Arm leads to speculation about the origin of other southern HI clouds. An important class of HVC that HIPASS is successful in revealing are the compact HVCs (CHVCs) which surround the Magellanic Stream and Leading Arm (see Figures 2 and 3). Are these objects additional interactive debris or possibly distant remnants of the Local Group (Blitz et al. 1999; Braun & Burton 1999)? The majority of the CHVCs follow the Magellanic complexes in position and velocity, suggesting they are related to the Stream or Leading Arm; however, between declinations -30° to -40° the CHVCs have a much larger velocity range ($\sim 500 \text{ km s}^{-1}$; from $v_{lsr} \approx -250$ to 250 km s^{-1}) than

¹ Figure 2 does not show the last $\approx 15^\circ$ of the Stream—see Wayte (1989) for detailed maps of this region.




Figure 2—Integrated intensity map of the Magellanic Stream ($v_{lsr} = -400$ to $+400$ km s $^{-1}$), which starts at the SMC and Bridge and extends to declination $+02^\circ$. The location of the clumps MS I to MS IV, which were originally classified by Mathewson et al. (1979), are labelled for reference. The Stream passes through the velocity of Galactic emission at approximately declination -35° , making it somewhat difficult to trace. At these velocities, Galactic emission has been blanked as much as possible.

the narrow range of the Stream material (~ 80 km s $^{-1}$). This declination region also encompasses the southern and near side (at ~ 2 Mpc) of the Sculptor Group of galaxies. Figure 4 shows the complexity of this region, with the positive velocity CHVCs mixed in amongst the galaxies while the Magellanic Stream continues southwards. These results are particularly interesting when one considers the recent distance determination work of Jerjen, Freeman & Binggeli (1998) which indicates that the Sculptor Group is not actually a separate group, but an extension of the Local Group, forming a 6×2 Mpc filament of galaxies with M31 and the Milky Way at one end.² The CHVCs along this sightline do not fit

into the normal distribution of Magellanic Stream material, and could lie anywhere along this filament of galaxies. At a distance of 2 Mpc most of the clouds would have HI masses of $\sim 10^7$ M $_{\odot}$ and diameters of ~ 20 kpc. The clouds may be leftover galactic building blocks and the local equivalent of the Ly-limit absorber (e.g. Stocke et al. 1995). They also fit nicely into hierarchical structure formation simulations which predict that there should be more satellites associated with the Local Group galaxies than are currently observed (e.g. Moore et al. 1999).

The Sculptor Group sightline may be a nearby tracer of the cosmic web and should be investigated more closely. H α observations are in progress and

² This prolate cloud of galaxies is actually part of an even larger structure known as the Coma–Sculptor cloud which stretches out to the Coma I cluster at 15 Mpc, in the direction of the Virgo cluster.




Figure 3—A HIPASS peak intensity map which shows the full extent of the Leading Arm, as well as the Magellanic Clouds, the Bridge and the beginning of the Stream (as labelled). The position of the background galaxy, NGC 3783, is also noted (see text). To avoid the emission from the Galactic Plane (which extends out to 120 km s^{-1} in this direction), only velocities between 130 and 400 km s^{-1} were used. (Thus the truncated appearance of the SMC which begins at $\approx 80 \text{ km s}^{-1}$.)

will provide significant insight into the origin of the CHVCs. If the CHVCs are within a 100 kpc radius from the Galactic Plane they should be detectable, as ionising photons from the Milky Way can easily reach this radius (Bland-Hawthorn & Maloney 1999a). This is shown through the $\text{H}\alpha$ detections of the Magellanic Stream and other HVCs which have been detected at the $0.2\text{--}0.4$ Rayleigh level (Weiner & Williams 1996; Tufte, Reynolds & Haffner 1998). If the population of CHVCs shown

here are truly extragalactic objects at Mpc distances they should not be detectable in $\text{H}\alpha$.

4 Conclusions

HIPASS provides a monumental amount of information on the distribution of neutral hydrogen in the southern sky. The interaction of the Magellanic Clouds with the Milky Way has created striking features, and the structure revealed here will offer important clues as to the exact mechanism responsible for their formation. The resolution and spatial sampling of HIPASS also allow the first thorough investigation of the population of compact HVCs and their possible extragalactic nature. The sightline to the Sculptor Group is an ideal place to begin this investigation. Further studies in progress of both the Magellanic and compact HVCs include sensitive $\text{H}\alpha$ observations, higher velocity resolution HI observations, and optical and UV measurements to determine metallicities.

Acknowledgments

I thank the multibeam working group for their help in the data collection and development of the instrumentation and software, in particular Lister Staveley-Smith.

References

- Bland-Hawthorn, J., & Maloney, P. 1999a, *ApJ*, 510, L33
- Bland-Hawthorn, J., & Maloney, P. 1999b, in *Stromlo Workshop on High-velocity Clouds*, ASP Conf. Series, Vol. 166 (San Francisco: ASP), p. 212
- Blitz, L., et al. 1999, *ApJ*, 514, 818
- Braun, R., & Burton, W. B. 1999, *A&A*, 341, 437
- Gardiner, L., & Noguchi, M. 1996, *MNRAS*, 278, 191
- Grondin, L., Demers, S., & Kunkel, W. E. 1992, *AJ*, 103, 1234
- Guhathakurta, P., & Reitzel, D. B. 1998, in *Galactic Halos: UC Santa Cruz Workshop*, ed. D. Zaritsky, ASP Conf. Series, Vol. 136 (San Francisco: ASP)
- Jerjen, H., Freeman, K., & Binggeli, B. 1998, *AJ*, 116, 2873
- Lu, L., et al. 1998, *AJ*, 115, 162.
- Mathewson, D. S., et al. 1979, in *The Large-scale Structure of the Galaxy* (Dordrecht: Kluwer), p. 556
- Mathewson, D. S., Cleary, M. N., & Murray, J. D. 1974, *ApJ*, 190, 291.
- Moore, B., et al. 1999, *ApJ*, 524, L19
- Putman, M. E. 2000, in *High-velocity Clouds*, ed. H. van Woerden et al. (Kluwer: Dordrecht) (see astro-ph/9909080)
- Putman, M. E., & Gibson, B. K. 1999, *PASA*, 16, 70
- Putman, M. E., Gibson, B. K., Staveley-Smith, L., et al. 1998, *Nature*, 394, 752
- Stocke, J. T., Shull, J. M., Penton, S., Donahue, M., & Carilli, C. 1995, 451, 24
- Tufte, S. L., Reynolds, R. J., & Haffner, L. M. 1998, *ApJ*, 504, 773
- Wakker, B. P., & van Woerden, H. 1997, *ARA&A*, 35, 21
- Wayte, S. R. 1989, *PASA*, 8, 195
- Weiner, B. J., & Williams, T. B. 1996, *AJ*, 111, 1156




Figure 4—The Magellanic Stream meets the Sculptor Group. A peak HI intensity map including velocities from 73 to 450 km s^{-1} . Several galaxies of the Sculptor Group are labelled, as well as the positions of two nearby dwarf galaxies. The bifurcation of the Magellanic Stream and the clumps which follow the Stream are also depicted. The contours are 0.4 , 0.8 , 1.6 , 3.2 and 6.4 K .