10.1071/CH20109\_AC

©CSIRO 2021

Australian Journal of Chemistry 2021, 74(2), 111-124

## **Supplementary Material**

## Some Products from C=O Condensations of Quinacridones

Anthony S. R. Chesman<sup>A,B,C</sup> and Andris J. Liepa<sup>A</sup>

<sup>A</sup>CSIRO Manufacturing, Jerry Price Laboratory, Clayton, Vic. 3168, Australia.

<sup>B</sup> Melbourne Centre of Nanofabrication, Clayton, Vic. 3168, Australia.

<sup>C</sup>Corresponding author. Email: <u>anthony.chesman@csiro.au</u>



Figure S1. <sup>1</sup>H NMR spectrum of **1b**.



Figure S2. <sup>1</sup>H NMR spectrum of Intermediate A. Peaks at 8.25, 3.24 and 3.11 ppm due to residual DMF.



Figure S3. <sup>13</sup>C NMR spectrum of Intermediate A.



Figure S4. a) <sup>1</sup>H NMR spectrum of 1d.



Figure S5. <sup>13</sup>C NMR spectrum of 1d.



**Figure S6.** <sup>1</sup>H NMR spectrum of **1f**. Peak at 5.35 ppm is residual DCM.



Figure S7. <sup>13</sup>C NMR spectrum of 1f.



Figure S8. <sup>1</sup>H NMR spectrum of 2a.



Figure S9. <sup>13</sup>C NMR spectrum of 2a.



Figure S10. <sup>1</sup>H NMR spectrum of 2b.



Figure S11. <sup>13</sup>C NMR spectrum of 2b.



Figure S12. <sup>1</sup>H NMR spectrum of 2c.



Figure S13.<sup>13</sup>C NMR spectrum of 2c.



**Figure S14.** <sup>1</sup>H NMR spectrum of **2d**. Peaks at 2.03 and 4.10 ppm due to residual ethyl acetate, which also contributes to peak series at 1.2-1.95 ppm.



Figure S15. <sup>13</sup>C NMR spectrum of 2d.



Figure S16. <sup>1</sup>H NMR spectrum of 3a.



Figure S17. <sup>13</sup>C NMR spectrum of 3a.



Figure S18. <sup>1</sup>H NMR spectrum of 3b.



Figure S19. <sup>13</sup>C NMR spectrum of 3b.



Figure S20. <sup>1</sup>H NMR spectrum of 3c.



Figure S21. <sup>13</sup>C NMR spectrum of 3c.



Figure S22. <sup>1</sup>H NMR spectrum of 3d. Peak at 2.63 ppm due to DMSO contamination.



Figure S23. <sup>13</sup>C NMR spectrum of 3d.



Figure S24. <sup>1</sup>H NMR spectrum of 3e.



Figure S25. <sup>13</sup>C NMR spectrum of 3e.



Figure S26. a) <sup>1</sup>H NMR spectrum 4a.



Figure S27. <sup>13</sup>C NMR spectrum 4a.



Figure S28. <sup>1</sup>H NMR spectrum of 4b.



Figure S29. <sup>13</sup>C NMR spectrum of 4b.



Figure S30. <sup>1</sup>H NMR spectrum of 4c.



Figure S31. <sup>13</sup>C NMR spectrum of 4c.



Figure S32. <sup>1</sup>H NMR spectrum of 5a.



Figure S33. <sup>13</sup>C NMR spectrum of 5a.



Figure S34. <sup>1</sup>H NMR spectrum of 5b.



Figure S35. <sup>13</sup>C NMR spectrum of 5b.



Figure S36. a) <sup>1</sup>H NMR spectrum of 5c.



Figure S37. <sup>13</sup>C NMR spectrum of 5c.



Figure S38. <sup>1</sup>H NMR spectrum of 6a.



Figure S39. <sup>13</sup>C NMR spectrum of 6a.



Figure S40. <sup>1</sup>H NMR spectrum of 6b.



Figure S41. <sup>13</sup>C NMR spectrum of 6b.



Figure S42. <sup>1</sup>H NMR spectrum of 6c. Peak at 3.49 ppm due to residual methanol.



Figure S43. <sup>13</sup>C NMR spectrum of 6c.



Figure S44. <sup>1</sup>H NMR spectrum of 6d.



Figure S45. <sup>13</sup>C NMR spectrum of 6d.



Figure S46. <sup>1</sup>H NMR spectrum of 6e.



Figure S47. <sup>13</sup>C NMR spectrum of 6e.