

10.1071/CP17204\_AC  
© CSIRO 2018

**Supplementary Material:** *Crop & Pasture Science*, 2018, 69, 131–141.

**Bacterial artificial chromosome clones randomly selected for sequencing reveal genomic differences between soybean cultivars**

*Tingting He<sup>A</sup>, Longshu Yang<sup>A</sup>, Xianlong Ding<sup>A</sup>, Linfeng Chen<sup>A</sup>, Yanwei Li<sup>A</sup>, Tanliu Wang<sup>A</sup>, Hao Zhang<sup>A</sup>, Junyi Gai<sup>A</sup> and Shouping Yang<sup>A,B</sup>*

<sup>A</sup>Soybean Research Institute, National Center for Soybean Improvement, Key Laboratory of Biology and Genetic Improvement of Soybean (General Ministry of Agriculture), State Key Laboratory of Crop Genetics and Germplasm Enhancement, Jiangsu Collaborative Innovation Center for Modern Crop Production, Nanjing Agricultural University, Nanjing 210095, China.

<sup>B</sup>Corresponding author. Email: [spyung@126.com](mailto:spyung@126.com)

**Supplementary Fig. 1.** Comparative analysis of the ten BAC clones in N23601 and their homologous regions in the reference genome.

