

10.1071/CP18569_AC

© CSIRO 2019

Supplementary Material: *Crop & Pasture Science*, 2019, 70, 555–565.

Elevated CO₂ causes large changes to morphology of perennial ryegrass (*Lolium perenne*)

Rose Brinkhoff^{A,B}, Meagan Porter^A and Mark J. Hovenden^A

^ABiological Sciences, School of Natural Sciences, University of Tasmania, Locked Bag 55, Hobart, Tas. 7001, Australia.

^BCorresponding author. Email: roseb2@utas.edu.au

Supplementary Table

Table S1. The effect of CO₂ treatment on mean soil volumetric water content as defined by the daily averages of the hourly values with *F* and *P* values from ANOVA.

Means in a single row with the same superscript are not significantly different ($P = 0.05$). Cells shaded light grey were receiving the limited water treatment in that month, and cells shaded dark grey received the excess water treatment

Month	Mean soil volumetric water content			<i>F</i>	<i>P</i>
	Adequate	Limited	Excess		
August*	25.6b	28.1a	25.6b	15.56	<0.001
September	26.9b	29.4a	28.8b	8.12	<0.001
October	30.8a	29.1b	30.5a	6.91	<0.001
November	25.1a	18.3a	21.8b	65.44	<0.001
December	24.7a	20.4b	21.5b	21.21	<0.001
January	20.5b	18.1c	27.0a	177.2	<0.001
February	25.1b	18.5c	28.5a	90.63	<0.001

*During winter all sectors received the same irrigation.