

10.1071/CP19017_AC

© CSIRO 2019

Supplementary Material: *Crop & Pasture Science*, 2019, 70, 463–472.

Nutritional and anti-nutritional seed-quality traits of faba bean (*Vicia faba*) grown in South Australia

Daniel J. Skylas^{A,B,F}, Jeffrey G. Paull^C, David G. D. Hughes^D, Beverley Gogel^D, Hao Long^E, Brett Williams^E, Sagadevan Mundree^E, Christopher L. Blanchard^B and Ken J. Quail^A

^AAustralian Export Grains Innovation Centre, North Ryde, NSW 2113, Australia.

^BARC Industrial Transformation Training Centre for Functional Grains, Charles Sturt University, Wagga Wagga, NSW 2650, Australia.

^CSchool of Agriculture, Food and Wine, Waite Campus, University of Adelaide, Glen Osmond, SA 5064, Australia.

^DCentre for Bioinformatics and Biometrics, National Institute for Applied Statistics Research Australia, Faculty of Engineering and Information Sciences, The University of Wollongong, NSW 2522, Australia.

^ECentre for Tropical Crops and Biocommodities, Queensland University of Technology, Brisbane, Qld 4001, Australia.

^FCorresponding author. Email: daniel.skylas@aegic.org.au

Supplementary Table

Table 1. Trial yields (t ha^{-1}) expressed as percentage of the site mean (2016 and 2017), sourced from the NVT program (<http://www.grdc-nvt.com.au>).

Variety	Freeling				Charlick			
	2016		2017		2016		2017	
	t ha^{-1}	%	t ha^{-1}	%	t ha^{-1}	%	t ha^{-1}	%
Doza	3.28	85	2.01	88	4.92	98	3.37	101
Farah	2.71	70	2.24	98	4.15	82	3.27	98
Fiesta VF	2.72	71	2.23	98	4.5	89	3.25	97
Fiord	2.86	74	1.87	82	4.7	93	3.36	101
Nura	3.59	93	2.19	96	4.23	84	3.08	92
PBA Nasma	3.02	78	2.07	90	5.22	104	3.3	99
PBA Rana	3.04	79	2.17	95	4.31	86	2.94	88
PBA Samira	4.92	127	2.67	117	5.22	104	3.42	103
PBA Warda	3.44	89	2.16	95	5.6	111	3.21	96
PBA Zahra	4.57	119	2.84	124	4.48	89	3.44	103
Site Mean	3.86		2.29		5.04		3.34	
CV (%)	9.76		8.21		8.85		7.46	
Probability	<0.001		<0.001		<0.001		<0.001	
LSD	0.6	15	0.31	13	0.6	12	0.34	10