

[10.1071/CP21265](https://doi.org/10.1071/CP21265)

Crop & Pasture Science

Supplementary Material

Quantification of endophyte *Serendipita indica* in *Brassica napus* roots by qPCR

Nastaran Abin^A, Nader Rokni^{B,}, Ali Reza Shafeinia^A, and M. Hossein Borhan^C*

^AFaculty of Agriculture, Department of Biotechnology, Ramin Agricultural and Natural Resources University, PO Box 73637-63417, Ahwaz, Iran.

^BFaculty of Science, Jundi-Shapur University of Technology, PO Box 64615-334, Dezful, Iran.

^CSaskatoon Research Centre, Agriculture and Agri-Food Canada, Saskatoon, SK, Canada.

*Correspondence to: Nader Rokni Faculty of Science, Jundi-Shapur University of Technology, PO Box 64615-334, Dezful, Iran Email: naderrokni@jsu.ac.ir

Supplementary Table S1. ANOVA analysis for “PRCs” after 25 d interaction between *Serendipta indica* and *Brassica napus* following the initial inoculation of seeds (g) with different weight ratios of fungal biomass (1, 3, 5, 7, and 10%)

Source	DF	Sum of Squares	Mean square	F value	Pr> F**
Treatment	4	0.728	0.182	4553.08	0.0001 >
Error	10	0.000	0.000		
Total	14	0.729			

Significant at $P = 0.0001$ **
PRC: percent root colonization

Supplementary Table S2. ANOVA analysis for the comparison of inoculum densities (Log CFU/g root) in the colonized *Brassica napus* roots resulting from inoculation of the seeds (g) with different weight ratios (1, 3, 5, 7, and 10%) of *Serendipita indica*

Source	DF	Sum of Squares	Mean square	F value	Pr> F**
Tre	5	55.955	11.191	466.217	< 0.0001
Error	25	0.600	0.024		
Total	30	56.555			

** Significant at $P = 0.0001$