

Experimental Agriculture

Contents Volume 47, Issue 5, 2007, 479–644

Animal Production

Comparing irrigated biodynamic and conventionally managed dairy farms. 1. Soil and pasture properties.
L. L. Burkitt, D. R. Small, J. W. McDonald, W. J. Wales and M. L. Jenkin 479–488

Comparing irrigated biodynamic and conventionally managed dairy farms. 2. Milk production and composition and animal health.

L. L. Burkitt, W. J. Wales, J. W. McDonald, D. R. Small and M. L. Jenkin 489–494

Effects of body condition and diet in late gestation on the subsequent health and performance of dairy cows.
C. R. Stockdale 495–501

Feeding concentrates based on individual cow requirements improves the yield of milk solids in dairy cows grazing restricted pasture.

S. C. García, M. Pedernera, W. J. Fulkerson, A. Horadagoda and K. Nandra 502–508

Seasonal variation in milk production and cheese yield from commercial dairy farms located in northern Victoria is associated with pasture and grazing management and supplementary feeding practices.

G. P. Walker, R. Williams, P. T. Doyle and F. R. Dunshea 509–524

Optimal use of on-farm fibre diameter measurement and its impact on reproduction in commercial Merino flocks.
M. J. Kelly, A. A. Swan and K. D. Atkins 525–534

Vitamin E and selenium status of sheep during autumn in Western Australia and its relationship to the incidence of apparent white muscle disease.

C. L. White and L. Rewell 535–543

Relationships between metabolic endocrine systems and voluntary feed intake in Merino sheep fed a high salt diet.
Dominique Blache, Micaela J. Grandison, David G. Masters, Robyn A. Dynes, Margaret A. Blackberry and Graeme B. Martin 544–550

Pasture, Fodder Crops

Evaluation of annual pasture legumes in northern New South Wales. 1. *Trifolium subterraneum* cultivars and lines.
G. M. Lodge and S. Harden 551–562

Evaluation of annual pasture legumes in northern New South Wales. 2. *Trifolium* and *Medicago* spp. and other legumes.

G. M. Lodge and S. Harden 563–574

Seasonal variation in the quality of a lucerne-based pasture and its relationship with morphological and maturity estimates.

C. F. Machado, S. T. Morris, J. Hodgson, C. Matthew and N. Auza 575–582

Field Crops

Yield components of nodulated cowpea (*Vigna unguiculata*) and maize (*Zea mays*) plants grown with exogenous phosphorus in different cropping systems.

Patrick A. Ndakidemi and Felix D. Dakora 583–589

Root penetration rate – a benchmark to identify soil and plant limitations to rooting depth in wheat.
J. A. Kirkegaard and J. M. Lilley 590–602

The effect of 3BS locus of Sumai 3 on Fusarium head blight resistance in Australian wheats.
G. Q. Xie, M. C. Zhang, S. Chakraborty and C. J. Liu 603–607

Residual effects from lime application on soil pH, rhizobial population and crop productivity in dryland farming systems of central New South Wales.

N. A. Fettell, C. M. Evans, D. J. Carpenter and J. Brockwell 608–619

The role of plant-parasitic nematodes in reducing yield of sugarcane in fine textured soils in Queensland, Australia.
B. L. Blair and G. R. Stirling 620–634

Horticulture

Ammonium thiosulfate and 6-benzyladenine improve the crop load and fruit quality of 'Delicious' apples.
S. A. Bound and S. J. Wilson 635–644

Front cover: Healthy Merino sheep used to investigate the interaction between grazing ruminants and their environment in the south-west of Western Australia. Photograph supplied Dr Hayley Norman, CSIRO Livestock Industries, Perth.