

Geophysical Signature of the Ellendale Lamproite Pipes, Western Australia

Graham Jenke¹ & Duncan R. Cowan²

1. Western Mining Corporation Ltd, P.O. Box 71, Kalgoorlie, W.A. 6430.
2. Cowan Geodata Services, 12 Edna Road, Dalkeith, W.A. 6009

Abstract

The Ellendale lamproite diatremes are located in the West Kimberley 125 km east-southeast of Derby. They are of Miocene age and occur in Lennard Shelf sedimentary rocks which overlie the King Leopold intracratonic Mobile Zone. The Ellendale province was discovered by the Ashton Joint Venture during stream sampling of the West Kimberley in 1976. Follow-up of indicator minerals led to the discovery of the diamondiferous vent, Ellendale 4, and nearby vents were then rapidly delineated by an aeromagnetic survey. Two pipes, Ellendale 4 and Ellendale 9, have significant diamond content but are subeconomic at present diamond prices.

The Ellendale province contains 48 lamproite intrusions in an elongate cluster, 40 km long by 10 km wide, oriented west-northwest, parallel to the major faults in the area. The lamproites are intruded into flat-lying Permian sandstones and Devonian to Carboniferous shales and limestones. The terrain is fairly flat with low hills bordering some vents.

A range of airborne and ground geophysical techniques has been used over the Ellendale lamproites.

Aeromagnetic, helicopter magnetic and ground magnetic surveys proved to be very effective, since the response of the weakly magnetic lamproites is quite clear against a background devoid of other shallow magnetic features. Airborne radiometric surveys proved effective in mapping areas covered by black soils and gave clear responses over a number of diatremes. Electromagnetic surveys were used to explore for lamproites which may not have a clear magnetic response. INPUT, DIGHEM, Turam and SIROTEM produced good responses in the more resistive areas but pipe responses were ambiguous in areas covered by conductive black soil.

PLACER DEPOSITS

Geophysical Characteristics of the Tertiary Palaeochannels in the Yilgarn Block, Western Australia

Erica L. Smyth¹ & Douglas M. Barrett²

1. Mineral Deposits Limited, BHP Minerals, P.O. Box 6062, East Perth, W.A. 6892. 2. Doug Barrett and Associates, 6 Yolande Place, City Beach, W.A. 6015

Abstract

The precise geometric definition of the Tertiary palaeochannels in the Yilgarn Block of Western Australia is important in the exploration for uranium and for secondary and placer gold which occur in the channel sediments. The positions of the palaeochannels normally show considerable displacement from the positions of the present-day drainages. Physical property contrasts which exist between the channel sediments and the underlying Archaean bedrock can be differentiated by geophysical methods to locate the best parts of the channels for follow-up drilling. The low densities of the channel-fill sediments in many areas give rise to 1 to 2 mGal gravity lows

and the gravity technique can be used to define the broad shape and depth of the channel. The deepest parts of the channel, which correspond to the zones of highest salinity, can then be delineated using time-domain electromagnetics. Downhole gamma logs are useful in identifying and quantifying prospective uranium zones.

Geophysical Surveys of the Eneabba Heavy-Mineral Sand Field, Eneabba, Western Australia

Stephen T. Mudge

RGC Exploration Pty Limited, P.O. Box 590, Fyshwick, A.C.T. 2609

Abstract

Magnetic petrophysical studies of ilmenite concentrates from the Eneabba area have shown that ilmenite-rich placer deposits can be detected with high-resolution magnetic surveys and that some of them can be remanently magnetised. Airborne magnetic surveys have been of limited use as the magnetic response of widespread surface laterite and cultural sources such as roads and buildings can mask the weak target anomalies. However, routine use of low-cost, high-resolution ground magnetics on east-west survey lines of up to 20 km length and spaced 1 km apart has proved successful in detecting the very weak anomalies which are typically less than 10 nT in amplitude. The results have been used to target exploratory drilling.

Airborne radiometrics has failed to detect buried thorium-rich monazite placers except where exposed by mining. Although induced polarisation and resistivity surveys have been used to detect ilmenite elsewhere, relatively high costs have precluded their use at Eneabba. Ground-penetrating radar failed to detect the buried mineralised placers.

Rutile-, zircon- and monazite-rich placers, which have minor concentrations of ilmenite, do not produce an observable magnetic response so extensive drilling is a necessary exploration technique. Because drilling is necessary and inexpensive, the routine application of geophysical techniques is restricted to ground magnetics, which is also inexpensive and able to quickly locate ilmenite-rich placers.

MULTISPECTRAL SCANNERS

Geoscan Airborne Multi-Spectral Scanners as Exploration Tools for Western Australian Diamond and Gold Deposits

Robert A. Agar

Geoscan Pty Ltd, 39 Hay Street, Subiaco, W.A. 6008

Abstract

The use of remote sensing in mineral exploration has evolved from basic photo-geology to interpretation of more sophisticated satellite and airborne multi-spectral data sets. Although the mineral mapping capabilities of Geoscan airborne multi-spectral scanners have been demonstrated for well-exposed and arid terrains, the question remains as to their effectiveness in deeply weathered regimes such as Western Australia.