

Supplementary material

Human impacts on connectivity in marine and freshwater ecosystems assessed using graph theory: a review

*Megan I. Saunders^{A,J}, Christopher J. Brown^A, Melissa M. Foley^{B,H}, Catherine M. Febria^C,
Rebecca Albright^{D,I}, Molly G. Mehling^E, Maria T. Kavanaugh^F, Dana D. Burfeind^G*

^AThe Global Change Institute, The University of Queensland,
St Lucia, Qld 4072, Australia;

^BCenter for Ocean Solutions, Stanford University, Monterey, CA 93940, USA.

^CFreshwater Ecology Research Group, School of Biological Sciences,
Christchurch, 8400, New Zealand.

^DAustralian Institute of Marine Science, Townsville MC, Townsville, Qld 4810, Australia.

^EFalk School of Sustainability, 1 Woodland Road, Chatham University,
Pittsburgh, PA 15232, USA.

^FMarine Chemistry and Geochemistry, Woods Hole Oceanographic Institution,
Woods Hole, MA 02540, USA.

^GThe School of Biological Sciences, The University of Queensland,
St Lucia, Qld 4072, Australia.

^HPresent address: United States Geological Survey, Pacific Coastal and Marine Science Center,
Santa Cruz, CA 95060, USA.

^IPresent address: Department of Global Ecology, Carnegie Institution for Science,
260 Panama Street, Stanford, CA 94305, USA.

^JCorresponding author: Email: m.saunders1@uq.edu.au

Table S1. Studies that have applied graph theory to spatial connectivity in aquatic ecosystems

Text in parentheses for ‘Human influence on connectivity’ are other influencing factors

Citation	Year	Aquatic realm	Quantitative or review	Ecosystem type	Ecosystem	Taxa	Organism	Life stage	Region	Human influence on connectivity
Adams <i>et al.</i> 2012	2012	Marine	Quantitative	Pelagic	Coastal pelagic	Invertebrates	Sea lice	Larvae	Scotland	Habitat addition (climate, salinity)
Anadón <i>et al.</i> 2013	2013	Marine	Quantitative	Benthic or pelagic	Temperate coastal	Fish	Fish (64 species)	Larvae	Gulf of California	Fishing – MPAs (spacing)
Andreollo <i>et al.</i> 2013	2013	Marine	Quantitative	Pelagic	Pelagic	Fish	Fish	Larvae	Mediterranean	Fishing – MPA; (location, season, PLD, Biological characteristics, vertical migration, spawning month)
Auerbach and Poff 2011	2011	Freshwater	Quantitative	Lotic	Dendritic	Not specified	Not specified	Unspecified	Theoretical	None (biological characteristics)
Barták <i>et al.</i> 2013	2013	Freshwater	Quantitative	Lotic	Rivers	Mammal	Beavers	Unspecified	Czech Republic	Habitat linkages – dams
Bunn <i>et al.</i> 2000	2000	Freshwater	Quantitative	Lotic	Coastal plain	Mammal, bird	Mink; Warblers	Adult	North Carolina	None (species)
Cantwell and Forman 1993	1993	Freshwater	Quantitative	Lotic	Rivers	Not specified	NA	Not specified		Habitat – landscape variability
Carranza <i>et al.</i> 2012	2012	Freshwater	Quantitative	Lotic	Rivers	Mammal	River otters	Adult	Italy	Habitat – loss and degradation
Decout <i>et al.</i> 2012	2012	Freshwater	Quantitative	Lentic	Ponds, forest	Amphibian	Common frog	Adult	French Alps	Habitat – land degradation (anthropogenic barriers, landscape permeability, habitat suitability)
Erős <i>et al.</i> 2011	2011	Freshwater	Quantitative	Lotic	Rivers	Fish	Fish	All	Hungary	Habitat – dams
Erős <i>et al.</i> 2012	2012	Freshwater	Review	Lotic	Dendritic					NA
Fortuna <i>et al.</i> 2006	2006	Freshwater	Quantitative	Lentic	Ephemeral ponds	Amphibian	Amphibians	Adult	Spain	Climate – drought
Fox and Bellwood 2014	2014	Marine	Quantitative	Demersal	Coral reefs	Fish	Reef fish	Adult	Great Barrier Reef	NA
Fullerton <i>et al.</i> 2011	2011	Freshwater	Quantitative	Lotic	Rivers	Fish	Pacific salmon	All	Oregon and Washington (lower quality, dams, fish)	Habitat – reduced habitat quality, dams, fish

Citation	Year	Aquatic realm	Quantitative or review	Ecosystem type	Ecosystem	Taxa	Organism	Life stage	Region	Human influence on connectivity
Campbell Grant <i>et al.</i> 2007	2007	Freshwater	Review	Lotic	Dendritic Rivers	Migratory fish	Fish	All	Columbia and Willamette Rivers)	hatcheries
Grill <i>et al.</i> 2014	2014	Freshwater	Quantitative	Lotic	Invertebrate	Crown of thorns starfish	Larvae	All	Mekong River Basin	NA
Hock <i>et al.</i> 2014	2014	Marine	Quantitative	Benthic or pelagic	Coral reefs	Fish, invertebrates	Fish, freshwater Invertebrates	Adult	Great Barrier Reef	Habitat – dams
Ishiyama <i>et al.</i> 2014	2014	Freshwater	Quantitative	Lotic	Wetlands	Bird	Shorebirds – 10 species	All	Japan	None (pest management, inferred)
Iwamura <i>et al.</i> 2013	2013	Marine	Quantitative	Benthic	Intertidal mudflats	Bird	Migratory shorebirds	Adult	East Asian– Australasian Flyway	Climate change – sea-level rise and habitat loss
Iwamura <i>et al.</i> 2014	2014	Marine	Quantitative	Benthic	Intertidal mudflats	Invertebrates	Mussels	All	East Asian– Australasian Flyway	Climate change – sea-level rise and habitat loss
Jacobi and Jonsson 2011	2011	Marine	Quantitative	Benthic or pelagic	Coastal temperate	Fish	Sharks (2 species)	Larvae	Baltic Sea	Fishing – MPAs
Jacoby <i>et al.</i> 2012	2012	Marine	Quantitative	Pelagic	Coastal pelagic	Fish	Fish	Adult	Bahamas; UK	Habitat removal
Jaeger <i>et al.</i> 2014	2014	Freshwater	Quantitative	Lentic	Streams	Invertebrates	Coral	All	American south-west	Climate change – drought
Kininmonth <i>et al.</i> 2010a	2010	Marine	Quantitative	Benthic or pelagic	Coral reefs	Wildlife – general	General or coral reef	Larvae	Great Barrier Reef	NA
Kininmonth <i>et al.</i> 2010b	2010	Marine	Quantitative	Benthic or pelagic	Theoretical – coral	Invertebrates	Theoretical or coral reef	Larvae	Great Barrier Reef	None (biological characteristics)
Kininmonth <i>et al.</i> 2011	2011	Marine	Quantitative	Benthic or pelagic	Wetlands	Fish	Wildlife – general	Larvae	Theoretical	Fishing – MPA network type
McIntyre and Strauss 2013	2013	Freshwater	Quantitative	Lentic	Dendritic Rivers	Fish	Wildlife – general		Texas	None (spatial scale)
McKay <i>et al.</i> 2013	2013	Freshwater	Quantitative	Lotic	Coral reefs	Invertebrates	Fish	Unspecified	Theoretical or Nevada, USA	Habitat – dams
Mumby <i>et al.</i> 2011	2011	Marine	Quantitative	Benthic or pelagic	Dentritic Ponds	Fish Reptile	Fish Turtles	Larvae	Bahamas	Climate change – warming temperatures
Padgham and Webb 2010	2010	Freshwater	Quantitative	Lotic	Streams	Fish Amphibians	Fish Frogs	Unspecified All	Theoretical Coastal plain, Portugal	Habitat quantity or quality
Pereira <i>et al.</i> 2011	2011	Freshwater	Quantitative	Lentic	Ponds				Habitat loss	
Perkin <i>et al.</i> 2013	2013	Freshwater	Quantitative	Lotic				Adult	Theoretical	Habitat – road crossings
Peterman <i>et al.</i> 2013	2013	Freshwater	Quantitative	Lotic				All	Missouri	(None – rainfall, population size, habitat type)

Citation	Year	Aquatic realm	Quantitative or review	Ecosystem type	Ecosystem	Taxa	Organism	Life stage	Region	Human influence on connectivity
Ribeiro <i>et al.</i> 2011	2011	Freshwater	Quantitative	Lentic	Ponds	Amphibian	Amphibians (9 species)	All	Central western Spain NA	None, although implications for MPA design discussed
Rossi <i>et al.</i> 2014	2014	Marine	Quantitative	Pelagic	Ocean basin	Not specified	n.a.	Larvae	Mediterranean	Habitat – dams
Schick and Lindley 2007	2007	Freshwater	Quantitative	Lotic	Dendritic	Fish	Fish	All	California	Habitat – dams
Segurado <i>et al.</i> 2013	2013	Freshwater	Quantitative	Lotic	Rivers	Fish	Fish	Adult	Portugal	Habitat – dams
Søndergaard and Jeppesen 2007	2007	Freshwater	Review	Lotic	Dendritic				NA	
Thomas <i>et al.</i> 2014	2014	Marine	Quantitative	Benthic or pelagic	Coral reefs	Invertebrates	Coral	Larvae	Great Barrier Reef	None – (species)
Tremel <i>et al.</i> 2008	2008	Marine	Quantitative	Benthic or pelagic	Coral reefs	Invertebrates	General – based on PLD – giant clam, coral	Larvae	Tropical Pacific	None (season; PLD)
Van Looy <i>et al.</i> 2013	2013	Freshwater	Quantitative	Lotic	Rivers	Mammal	River otters	Adult	France	Habitat – river straightening, urbanisation, forest fragmentation
Watson <i>et al.</i> 2011	2011	Marine	Quantitative	Benthic or pelagic	Coastal pelagic	Fish, invertebrates	Fish, sea urchin	Larvae	California	Fishing – MPAs

References

- Adams, T., Black, K., MacIntyre, C., MacIntyre, I., and Dean, R. (2012). Connectivity modelling and network analysis of sea lice infection in Loch Fyne, west coast of Scotland. *Aquaculture Environment Interactions* **3**, 51–63 [doi:10.3354/aei00052](https://doi.org/10.3354/aei00052).
- Anadón, J. D., del Mar Mancha-Cisneros, M., Best, B. D., and Gerber, L. R. (2013). Habitat-specific larval dispersal and marine connectivity: implications for spatial conservation planning. *Ecosphere* **4**, art82 [doi:10.1890/ES13-00119.1](https://doi.org/10.1890/ES13-00119.1).
- Andrello, M., Mouillot, D., Beuvier, J., Albouy, C., Thuiller, W., and Manel, S. (2013). Low connectivity between Mediterranean marine protected areas: a biophysical modeling approach for the dusky grouper *Epinephelus marginatus*. *PLoS One* **8**(7), e68564 [doi:10.1371/journal.pone.0068564](https://doi.org/10.1371/journal.pone.0068564).
- Auerbach, D. A., and Poff, N. L. (2011). Spatiotemporal controls of simulated metacommunity dynamics in dendritic networks. *Journal of the North American Benthological Society* **30**, 235–251. [doi:10.1899/09-126.1](https://doi.org/10.1899/09-126.1)

- Barták, V., Vorel, A., Šimová, P., and Puš, V. (2013). Spatial spread of Eurasian beavers in river networks: a comparison of range expansion rates. *Journal of Animal Ecology* **82**, 587–597. [doi:10.1111/j.1365-2656.12040](https://doi.org/10.1111/j.1365-2656.12040)
- Bunn, A. G., Urban, D. L., and Keitt, T. H. (2000). Landscape connectivity: A conservation application of graph theory. *Journal of Environmental Management* **59**, 265–278. [doi:10.1006/jema.2000.0373](https://doi.org/10.1006/jema.2000.0373)
- Campbell Grant, E. H., Lowe, W. H., and Fagan, W. F. (2007). Living in the branches: population dynamics and ecological processes in dendritic networks. *Ecology Letters* **10**, 165–175 [doi:10.1111/j.1461-0248.2006.01007.x](https://doi.org/10.1111/j.1461-0248.2006.01007.x).
- Cantwell, M. D., and Forman, R. T. T. (1993). Landscape graphs – ecological modeling with graph-theory to detect configurations common to diverse landscapes. *Landscape Ecology* **8**, 239–255 [doi:10.1007/BF00125131](https://doi.org/10.1007/BF00125131).
- Carranza, M. L., D'Alessandro, E., Saura, S., and Loy, A. (2012). Connectivity providers for semi-aquatic vertebrates: the case of the endangered otter in Italy. *Landscape Ecology* **27**, 281–290 [doi:10.1007/s10980-011-9682-3](https://doi.org/10.1007/s10980-011-9682-3).
- Decout, S., Manel, S., Miaud, C., and Luque, S. (2012). Integrative approach for landscape-based graph connectivity analysis: a case study with the common frog (*Rana temporaria*) in human-dominated landscapes. *Landscape Ecology* **27**, 267–279 [doi:10.1007/s10980-011-9694-z](https://doi.org/10.1007/s10980-011-9694-z).
- Erős, T., Schmera, D., and Schick, R. S. (2011). Network thinking in riverscape conservation – a graph-based approach. *Biological Conservation* **144**, 184–192 [doi:10.1016/j.biocon.2010.08.013](https://doi.org/10.1016/j.biocon.2010.08.013).
- Erős, T., Olden, J. D., Schick, R. S., Schmera, D., and Fortin, M. J. (2012). Characterizing connectivity relationships in freshwaters using patch-based graphs. *Landscape Ecology* **27**, 303–317 [doi:10.1007/s10980-011-9659-2](https://doi.org/10.1007/s10980-011-9659-2).
- Fortuna, M. A., Gómez-Rodríguez, C., and Bascompte, J. (2006). Spatial network structure and amphibian persistence in stochastic environments. *Proceedings. Biological Sciences* **273**, 1429–1434 [doi:10.1098/rspb.2005.3448](https://doi.org/10.1098/rspb.2005.3448).
- Fox, R. J., and Bellwood, D. R. (2014). Herbivores in a small world: network theory highlights vulnerability in the function of herbivory on coral reefs. *Functional Ecology* **28**, 642–651 [doi:10.1111/j.1365-2435.12190](https://doi.org/10.1111/j.1365-2435.12190).
- Fullerton, A. H., Lindley, S. T., Pess, G. R., Feist, B. E., Steel, E. A., and McElhany, P. (2011). Human influence on the spatial structure of threatened Pacific salmon metapopulations. *Conservation Biology* **25**, 932–944 [doi:10.1111/j.1523-1739.2011.01718.x](https://doi.org/10.1111/j.1523-1739.2011.01718.x).

- Grill, G., Ouellet Dallaire, C., Fluet Chouinard, E., Sindorf, N., and Lehner, B. (2014). Development of new indicators to evaluate river fragmentation and flow regulation at large scales: a case study for the Mekong River Basin. *Ecological Indicators* **45**, 148–159 [doi:10.1016/j.ecolind.2014.03.026](https://doi.org/10.1016/j.ecolind.2014.03.026).
- Hock, K., Wolff, N. H., Condie, S. A., Anthony, K. R. N., and Mumby, P. J. (2014). Connectivity networks reveal the risks of crown-of-thorns starfish outbreaks on the Great Barrier Reef. *Journal of Applied Ecology* **51**, 1188–1196 [doi:10.1111/1365-2664.12320](https://doi.org/10.1111/1365-2664.12320).
- Ishiyama, N., Akasaka, T., and Nakamura, F. (2014). Mobility-dependent response of aquatic animal species richness to a wetland network in an agricultural landscape. *Aquatic Sciences* **76**(3), 437–449. [doi:10.1007/s00027-014-0345-8](https://doi.org/10.1007/s00027-014-0345-8)
- Iwamura, T., Possingham, H. P., Chades, I., Minton, C., Murray, N. J., Rogers, D. I., Treml, E. A., and Fuller, R. A. (2013). Migratory connectivity magnifies the consequences of habitat loss from sea-level rise for shorebird populations. *Proceedings. Biological Sciences* **280**, 20130325. [doi:10.1098/rspb.2013.0325](https://doi.org/10.1098/rspb.2013.0325)
- Iwamura, T., Fuller, R. A., and Possingham, H. P. (2014). Optimal management of a multispecies shorebird flyway under sea-level rise. *Conservation Biology* **28**, 1710–1720 [doi:10.1111/cobi.12319](https://doi.org/10.1111/cobi.12319).
- Jacobi, M. N., and Jonsson, P. R. (2011). Optimal networks of nature reserves can be found through eigenvalue perturbation theory of the connectivity matrix. *Ecological Applications* **21**, 1861–1870. [doi:10.1890/10-0915.1](https://doi.org/10.1890/10-0915.1)
- Jacoby, D. M. P., Brooks, E. J., Croft, D. P., and Sims, D. W. (2012). Developing a deeper understanding of animal movements and spatial dynamics through novel application of network analyses. *Methods in Ecology and Evolution* **3**, 574–583 [doi:10.1111/j.2041-210X.2012.00187.x](https://doi.org/10.1111/j.2041-210X.2012.00187.x).
- Jaeger, K. L., Olden, J. D., and Pelland, N. A. (2014). Climate change poised to threaten hydrologic connectivity and endemic fishes in dryland streams. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 13894–13899 [doi:10.1073/pnas.1320890111](https://doi.org/10.1073/pnas.1320890111).
- Kininmonth, S. J., De'ath, G., and Possingham, H. P. (2010a). Graph theoretic topology of the Great but small Barrier Reef world. *Theoretical Ecology* **3**, 75–88 [doi:10.1007/s12080-009-0055-3](https://doi.org/10.1007/s12080-009-0055-3).
- Kininmonth, S., van Oppen, M. J. H., and Possingham, H. P. (2010b). Determining the community structure of the coral *Seriatopora hystrix* from hydrodynamic and genetic networks. *Ecological Modelling* **221**, 2870–2880. [doi:10.1016/j.ecolmodel.2010.08.042](https://doi.org/10.1016/j.ecolmodel.2010.08.042)
- Kininmonth, S., Beger, M., Bode, M., Peterson, E., Adams, V. M., Dorfman, D., Brumbaugh, D. R., and Possingham, H. P. (2011). Dispersal connectivity and reserve selection for marine conservation. *Ecological Modelling* **222**, 1272–1282 [doi:10.1016/j.ecolmodel.2011.01.012](https://doi.org/10.1016/j.ecolmodel.2011.01.012).

- McIntyre, N., and Strauss, R. (2013). A new, multi-scaled graph visualization approach: an example within the playa wetland network of the Great Plains. *Landscape Ecology* **28**, 769–782. [doi:10.1007/s10980-013-9862-4](https://doi.org/10.1007/s10980-013-9862-4)
- McKay, S. K., Schramski, J. R., Conyngham, J. N., and Fischenich, J. C. (2013). Assessing upstream fish passage connectivity with network analysis. *Ecological Applications* **23**, 1396–1409 [doi:10.1890/12-1564.1](https://doi.org/10.1890/12-1564.1).
- Mumby, P. J., Elliott, I. A., Eakin, C. M., Skirving, W., Paris, C. B., Edwards, H. J., Enríquez, S., Iglesias-Prieto, R., Cherubin, L. M., and Stevens, J. R. (2011). Reserve design for uncertain responses of coral reefs to climate change. *Ecology Letters* **14**, 132–140 [doi:10.1111/j.1461-0248.2010.01562.x](https://doi.org/10.1111/j.1461-0248.2010.01562.x).
- Padgham, M., and Webb, J. A. (2010). Multiple structural modifications to dendritic ecological networks produce simple responses. *Ecological Modelling* **221**, 2537–2545. [doi:10.1016/j.ecolmodel.2010.04.022](https://doi.org/10.1016/j.ecolmodel.2010.04.022)
- Pereira, M., Segurado, P., and Neves, N. (2011). Using spatial network structure in landscape management and planning: a case study with pond turtles. *Landscape and Urban Planning* **100**, 67–76 [doi:10.1016/j.landurbplan.2010.11.009](https://doi.org/10.1016/j.landurbplan.2010.11.009).
- Perkin, J. S., Gido, K. B., Al-Ta’ani, O., and Scoglio, C. (2013). Simulating fish dispersal in stream networks fragmented by multiple road crossings. *Ecological Modelling* **257**, 44–56 [doi:10.1016/j.ecolmodel.2013.02.021](https://doi.org/10.1016/j.ecolmodel.2013.02.021).
- Peterman, W., Rittenhouse, T. G., Earl, J., and Semlitsch, R. (2013). Demographic network and multi-season occupancy modeling of *Rana sylvatica* reveal spatial and temporal patterns of population connectivity and persistence. *Landscape Ecology* **28**(8), 1601–1613. [doi:10.1007/s10980-013-9906-9](https://doi.org/10.1007/s10980-013-9906-9)
- Ribeiro, R., Carretero, M., Sillero, N., Alarcos, G., Ortiz-Santaliestra, M., Lizana, M., and Llorente, G. (2011). The pond network: can structural connectivity reflect on (amphibian) biodiversity patterns? *Landscape Ecology* **26**, 673–682 [doi:10.1007/s10980-011-9592-4](https://doi.org/10.1007/s10980-011-9592-4).
- Rossi, V., Ser-Giacomi, E., López, C., and Hernández-García, E. (2014). Hydrodynamic provinces and oceanic connectivity from a transport network help designing marine reserves. *Geophysical Research Letters* **41**, 2883–2891 [doi:10.1002/2014GL059540](https://doi.org/10.1002/2014GL059540).
- Schick, R. S., and Lindley, S. T. (2007). Directed connectivity among fish populations in a riverine network. *Journal of Applied Ecology* **44**, 1116–1126 [doi:10.1111/j.1365-2664.2007.01383.x](https://doi.org/10.1111/j.1365-2664.2007.01383.x).
- Segurado, P., Branco, P., and Ferreira, M. T. (2013). Prioritizing restoration of structural connectivity in rivers: a graph based approach. *Landscape Ecology* **28**, 1231–1238 [doi:10.1007/s10980-013-9883-z](https://doi.org/10.1007/s10980-013-9883-z).

- Søndergaard, M., and Jeppesen, E. (2007). Anthropogenic impacts on lake and stream ecosystems, and approaches to restoration. *Journal of Applied Ecology* **44**, 1089–1094 [doi:10.1111/j.1365-2664.2007.01426.x](https://doi.org/10.1111/j.1365-2664.2007.01426.x).
- Thomas, C. J., Lambrechts, J., Wolanski, E., Traag, V. A., Blondel, V. D., Deleersnijder, E., and Hanert, E. (2014). Numerical modelling and graph theory tools to study ecological connectivity in the Great Barrier Reef. *Ecological Modelling* **272**, 160–174. [doi:10.1016/j.ecolmodel.2013.10.002](https://doi.org/10.1016/j.ecolmodel.2013.10.002)
- Treml, E. A., Halpin, P. N., Urban, D. L., and Pratson, L. F. (2008). Modeling population connectivity by ocean currents, a graph-theoretic approach for marine conservation. *Landscape Ecology* **23**, 19–36 [doi:10.1007/s10980-007-9138-y](https://doi.org/10.1007/s10980-007-9138-y).
- Van Looy, K., Cavillon, C., Tormos, T., Piffady, J., Landry, P., and Souchon, Y. (2013). A scale-sensitive connectivity analysis to identify ecological networks and conservation value in river networks. *Landscape Ecology* **28**, 1239–1249 [doi:10.1007/s10980-013-9869-x](https://doi.org/10.1007/s10980-013-9869-x).
- Watson, J. R., Siegel, D. A., Kendall, B. E., Mitarai, S., Rassweiller, A., and Gaines, S. D. (2011). Identifying critical regions in small-world marine metapopulations. *Proceedings of the National Academy of Sciences of the United States of America* **108**, E907–E913 [doi:10.1073/pnas.1111461108](https://doi.org/10.1073/pnas.1111461108).