

Supplementary material

Metabarcoding demonstrates dietary niche partitioning in two coexisting blackfish species

Gavin N. Rees^{A,F}, Michael E. Shackleton^B, Garth O. Watson^A, Georgia K. Dwyer^{C,D}
and Rick J. Stoffels^{A,E}

^ACSIRO Land and Water, Institute for Land, Water and Society, Charles Sturt University, Elizabeth Mitchell Drive, Thurgoona, NSW 2460, Australia.

^BCentre for Freshwater Ecosystems (CFE), La Trobe University, University Drive, Wodonga, Vic. 3690, Australia.

^CDepartment of Ecology, Environment and Evolution (DEEE), La Trobe University, University Drive, Wodonga, Vic. 3690, Australia.

^DPresent address: Centre of Regional and Rural Futures, Deakin University, Locked Bag 20000, Geelong, Vic. 3220, Australia.

^EPresent address: National Institute of Water and Atmospheric Research (NIWA), Riccarton 8011, Christchurch, New Zealand.

^FCorresponding author. Email: gavin.rees@csiro.au




Fig. S1. Frequency of occurrence of the individual operational taxonomic units (OTUs), as generated by the USEARCH pipeline, across the 40 *Gadopsis* specimens.




Fig. S2. Nonmetric multi-dimensional scaling ordination depicting the diets of *Gadopsis* individuals, based on our reduced taxon list (see methods for full description on how the reduced taxon list was derived).