

[10.1071/MF22081](https://doi.org/10.1071/MF22081)

*Marine and Freshwater Research*

### **Supplementary Material**

#### **Characterising fish habitat use of fringing oyster reefs using acoustic imaging**

*Robert P. Dunn<sup>A,B,\*</sup>, Matthew E. Kimball<sup>B</sup>, Caitlin G. O'Brien<sup>B</sup>, and Nathan T. Adams<sup>B</sup>*

<sup>A</sup>North Inlet–Winyah Bay National Estuarine Research Reserve, Georgetown, SC 29440, USA.

<sup>B</sup>Baruch Marine Field Laboratory, University of South Carolina, 2306 Crabhall Road, Georgetown, SC 29440, USA.

\*Correspondence to: Robert P. Dunn North Inlet–Winyah Bay National Estuarine Research Reserve, Georgetown, SC 29440, USA Email: robert@baruch.sc.edu

**Table S1:** Location of oyster reefs ( $n = 10$ ) sampled in Clambank Creek, North Inlet estuary, Georgetown County SC, USA.

Oyster Reef	Latitude (N)	Longitude (W)
1	33.33421	79.19257
2	33.33451	79.19212
3	33.33466	79.19195
4	33.33479	79.19171
5	33.33495	79.19081
6	33.33417	79.19050
7	33.33253	79.18958
8	33.33214	79.18922
9	33.33195	79.18888
10	33.33185	79.18831



**Figure S1:** Top: map of North Inlet Estuary, Georgetown Count, SC, USA. The white box outlines Clambank Creek, the general study area, in the central port portion of the estuary. Bottom: map of Clambank Creek with white rectangle denoting the specific stretch of tidal creek, which includes the locations of the oyster reefs sampled using the ARIS acoustic imaging sonar.