10.1071/MF22260

Marine and Freshwater Research

Supplementary Material

Plasticity of upper thermal limits of Australian *Paratya* spp. (Decapoda, Atyidae) and considerations of climate-change adaptation

Brendan Cox^{A,*}, Amanda Reichelt-Brushett^A, Kathryn Taffs^{A,B}, and Ross Smith^{A,C}

^AFaculty of Science and Engineering, Southern Cross University, Lismore, NSW, Australia.

^BNSW Department of Planning and Environment, Water Science, Ballina, NSW, Australia.

^cHydrobiology Pty Ltd, Brisbane, Qld, Australia.

*Correspondence to: Brendan Cox Faculty of Science and Engineering, Southern Cross University, Lismore, NSW, Australia Email: brendan.cox@scu.edu.au

Stream temperature




Figure S1. Stream water temperature from collection site at Leycester Creek. Three months of water temperature was collected every 15 min using hobo data loggers.

Data

Table S1. Winter acclimatised shrimp replicate PROBIT results sowing estimates and standard deviations.

Replicate	A	В	С	D	Е	F	G	Н	I	Mean	SD
LT ₁₀	31.714	31.714	31.219	31.535	32.633	32.996	32.563	32.559	31.003	32.0	0.667
LT50	34.14	34.14	33.611	33.948	35.12	35.508	35.046	35.041	34.827	34.6	0.661
LT ₉₀	36.565	36.565	36.002	36.361	37.607	38.019	37.529	37.523	38.651	37.2	0.823

Table S2. Warm water laboratory acclimated shrimp results of PROBIT showing estimates and standard deviation.

Replicate	A	В	С	D	Е	F	G	Н	I	Mean	SD
LT ₁₀	35.737	35.61	35.71	35.368	34.89	34.703	35.71	32.392	35.791	35.1	1.027
LT ₅₀	36.363	36.217	36.332	36.147	35.465	35.276	36.332	36.485	36.426	36.1	0.411
LT ₉₀	36.988	36.824	36.954	36.926	36.04	35.849	36.954	40.578	37.061	37.1	1.288