

[10.1071/PC23014](https://doi.org/10.1071/PC23014)

Pacific Conservation Biology

Supplementary Material

The role of grass-tree *Xanthorrhoea semiplana* (Asphodelaceae) canopies in temperature regulation and waterproofing for ground-dwelling wildlife

Sophie Petit^{A,B,}, and Deborah S. Frazer^{A,B}*

^AUniSA STEM, University of South Australia, Mawson Lakes, SA 5095, Australia.

^BKangaroo Island Research Station, Dudley West, Penneshaw, SA 5222, Australia.

*Correspondence to: Sophie Petit UniSA STEM, University of South Australia, Mawson Lakes, SA 5095, Australia Email: sophie.petit@unisa.edu.au

Table S1. Rounded temperature means for the four sites at the times (night in winter – cold, day in summer – hot) of greatest differences between two conditions (grass-tree and ambient or grass-tree and random areas 2 m away). The largest differences over 24-h periods were selected for n = 12 grass-trees at each site, except in winter for Warren (n = 10) and Scott Creek (n = 11). Differences and means of differences are also presented. GT = grass-tree; blue colour indicates winter and orange summer.

	GT means winter	ambient winter	GT - ambient	GT means winter	2 m away winter	GT - 2 m away	GT means summer	ambient summer	GT - ambient	GT summer	2 m away summer	GT - 2 m away summer
Warren	6.9	6.1	0.8	6.9	5.7	1.2	21.1	30.0	-8.9	21.4	41.2	-19.8
Scott Creek	6.0	2.6	3.4	6.0	3.6	2.4	17.3	37.0	-19.7	17.3	28.2	-10.9
Deep Creek	7.6	5.1	2.5	7.1	6.1	1.0	17.9	32.0	-14.1	17.9	33.1	-15.2
Para Wirra	8.2	5.6	2.6	8.2	5.6	2.6	23.3	37.0	-13.7	23.6	42.6	-19.0
means			2.3175			1.785			-14.100			-16.225