

Supplementary Material

Stathmin 1 plays a role in endometrial decidualisation by regulating hypoxia inducible factor-1 α and vascular endothelial growth factor during embryo implantation

Jinhai Gou^A, Jia Jia^A, Juntao Feng^A, Xia Zhao^{A,B}, Tao Yi^B, Tao Cui^A and Zhengyu Li^{A,B,C}

^ADepartment of Gynecology and Obstetrics, West China Second University Hospital, Sichuan University, No. 20, Section 3, Renmin South Road, Chengdu, Sichuan, 610041, PR China.

^BSichuan Key Laboratory of Gynecologic Oncology, West China Second University Hospital, Sichuan University, No.20, Section 3, Renmin South Road, Chengdu, Sichuan, 610041, PR China.

^CCorresponding author. Email: zhengyuli@scu.edu.cn




Fig. S1. Effect of *Stmn1*-siRNA on the phenotype of the surviving embryos. (A) Implanted Embryos from Kunming White strain mice on D15. (B) Placentae from three groups: Blank, *Stmn1*-siRNA, NC-siRNA on D15. (C) Hematoxylin-eosin (HE) staining of uterine placenta on D15 during embryo implantation in mouse. Magnification is indicated by the bars which represent 100um in upper panels and 25um in lower panels, respectively.




Fig. S2. Place of stathmin1 (downstream/upstream) in the various signaling pathways. →

Direct Stimulatory Modification, —| Direct Inhibitory Modification, - - - → Translocation.