

10.1071/RD16056_AC

© CSIRO 2016

Supplementary Material: *Reproduction, Fertility and Development*

Supplementary Material

Evidence of proteinuria, but no other characteristics of pre-eclampsia, in relaxin-deficient mice

Kelly P. O'Sullivan^A, Sarah A. Marshall^A, Scott Cullen^A, Tahnee Saunders^A, Natalie J. Hannan^B, Sevvandi N. Senadheera^A and Laura J. Parry^{A,C}

^ASchool of BioSciences, The University of Melbourne, Parkville, Vic., 3010, Australia.

^BThe Translational Obstetrics Group, Mercy Hospital for Women, Department of Obstetrics and Gynaecology, The University of Melbourne, Vic., 3010, Australia.

^CCorresponding author. Email: ljparry@unimelb.edu.au

Table S1. Primer and probe sequences for the quantitative amplification of murine genes

Gene	Accession number	Oligo type	Primer sequence 5'-3'
<i>18S</i>	NR_003278	Forward	GGGCTACCACATCCAAGGAA
		Reverse	GCTGGAATTAAACCGCGGCT
		Probe	TGCTGGCACCAGACTTGCCCTC
<i>Plgf</i>	NM_008827	Forward	TTCAGTCCGTCCTGTGTCCTT
		Reverse	GCACACAGTGCAGACCTTCA
		Probe	ACCACAGCAGCCACTACAGCGACTCA
<i>Rxfp1</i>	NM_212452	Forward	GCTTCCACTAACTCCTTTGA
		Reverse	CTCGGCACAAACAATGCATG
		Probe	AAACTTCCGAATGCGTGGTTGGCTC
<i>VegfA</i>	NM_001025250	Forward	TGAGACCCTGGTGGACATCT
		Reverse	TTTGACCCTTTCCCTTTCCT
		Probe	CAGGCTGCTGTAACGATGAA