

10.1071/RD16343_AC

© CSIRO 2017

Supplementary Material: *Reproduction, Fertility and Development*, 2017, 29(12), 2336–2344.

Supplementary Material

Andrographolide disrupts meiotic maturation by blocking cytoskeletal reorganisation and decreases the fertilisation potential of mouse oocytes

Hong-xing Liang^A, Sheng-sheng Lu^A, Zheng Yan^{B,C}, Yan-ping Kuang^B, Xiang-xing Zhu^A, Zhi-guang Yan^B, Tong Du^B, Wei-ran Chai^B, Hui Long^{B,C} and Qi-feng Lyu^{B,C}

^AState Key Laboratory for Conservation and Utilisation of Subtropical Agro-bioresources, Guangxi High Education Key Laboratory for Animal Reproduction and Biotechnology, College of Animal Science and Technology, Guangxi University, Nanning, Guangxi 530004, PR China.

^BDepartment of Assisted Reproduction, Shanghai Ninth People's Hospital Affiliated Shanghai Jiao Tong University School of Medicine, Shanghai 200011, PR China.

^CCorresponding authors. Email: lyuqifeng@126.com; yanzheng369@163.com; longhuish@aliyun.com




Fig S1. Effects of AG treatment on mouse *in vivo* matured oocyte fertilisation

(A) Representative phase-contrast images of oocytes treated with or without AG

for 2 h before ICSI. Fertilization was assessed 6 h later. Scale bar: 100 μm . (B) Bar graph represents the oocyte fertilization rate in each of the treatment groups (KOSM control group, $n = 64$; 5 μM AG group, $n = 79$; 10 μM AG group, $n = 66$; $n =$ oocyte number). (C) Representative phase-contrast images of oocytes treated with or without AG for 2 h after ICSI. Fertilisation was assessed 6 h later. Scale bar: 100 μm . (D) Bar graph represents the oocyte fertilization rate in each of the treatment groups (KOSM control group, $n = 60$; 5 μM AG group, $n = 64$; 10 μM AG group, $n = 71$; $n =$ oocyte number). Data obtained from three independent experiments were pooled and were analysed with chi-squared tests.