

Supplementary Material

Localisation of stem cell factor, stanniocalcin-1, connective tissue growth factor and heparin-binding epidermal growth factor in the bovine uterus at the time of blastocyst formation

M. Muñoz^{A,E}, D. Martín^A, S. Carrocera^A, M. Alonso-Guervos^B, M. I. Mora^C, F. J. Corrales^C, N. Peynot^D, C. Giraud-Delville^D, V. Duranthon^D, O. Sandra^D and E. Gómez^A

^ACentro de Biotecnología Animal, Servicio Regional de Investigación y Desarrollo Agroalimentario (SERIDA), Camino de Rioseco 1225, 33394 Gijón, Spain.

^BUnidad de Microscopía Fotónica y Proceso de Imágenes, Servicios Científico Técnicos, Universidad de Oviedo, Instituto Universitario de Oncología de Asturias (IUOPA), 33006, Oviedo, Spain.

^CUnidad de Proteómica, Centro de Investigación Médica Aplicada (CIMA) – Universidad de Navarra, 31008, Pamplona, Navarra, Spain.

^DUMR BDR, INRA, ENVA, Université Paris Saclay, 78350, Jouy en Josas, France.

^ECorresponding author. Email: mmunoz@serida.org




Fig. S1. Control samples were processed by omitting the primary antibodies. Bovine endometrial sections (A, B) or blastocysts, incubated without the first (primary) antibodies showed no signal. Nuclei of endometrial cells and embryonic cells were counter-stained with hematoxylin and Dapi, respectively (blue color). Black scale bars: 15 μ m, green scale bar: 50 μ m.