

10.1071/RD16477\_AC

© CSIRO 2017

Supplementary Material: *Reproduction, Fertility and Development*, 2017, 29(11), 2149–2156.

## Supplementary Material

### **Tissue-specific inhibition of urokinase-type plasminogen activator expression in the testes of mice by inducible lentiviral RNA interference causes male infertility**


*Kai Zhao<sup>A</sup>, Yan Liu<sup>B</sup>, Zhe Xiong<sup>C</sup>, Lian Hu<sup>A,B,D</sup> and Cheng-liang Xiong<sup>A,B</sup>*

<sup>A</sup>Family Planning Research Institute and Center of Reproductive Medicine, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430030, China.

<sup>B</sup>Wuhan Tongji Reproductive Medicine Hospital, Wuhan, 430014, China.

<sup>C</sup>Department of Physiology, Medical School of Jiangnan University, Wuhan, 430056, China.

<sup>D</sup>Corresponding author. Email: hulian02@163.com



**Fig. S1.** Pregnancy rate and litter size of female mice for different weeks. (A) Pregnancy rate of female mice after recovery for 2, 4 and 8 weeks. (B) Litter size of female mice after recovery for 2, 4 and 8 weeks (▲:  $P < 0.05$  compared with blank control).