

10.1071/RD16486_AC

© CSIRO 2018

Supplementary Material: *Reproduction, Fertility and Development*, 2018, 30(3), 519–532.

Supplementary Material

Insulinotropic nucleobindin-2/nesfatin-1 is dynamically expressed in the haemochorial mouse and human placenta

Crystalyn B. Legg-St Pierre^{A,B}, Martina Mackova^C, Ewa I. Miskiewicz^{A,B}, Denise G. Hemmings^C, Suraj Unniappan^{A,B} and Daniel J. MacPhee^{A,B,D,E}

^ADepartment of Veterinary Biomedical Sciences, Western College of Veterinary Medicine, University of Saskatchewan, Saskatoon, SK, S7N 5B4, Canada.

^BOne Reproductive Health Research Group, University of Saskatchewan, Saskatoon, SK, S7N 5B4, Canada.

^CDepartment of Obstetrics and Gynecology, Faculty of Medicine and Dentistry, 227 Heritage Medical Research Centre, University of Alberta, Edmonton, AB, T6G 2S2, Canada.

^DDivision of Biomedical Sciences, Faculty of Medicine, Memorial University of Newfoundland, St. John's, NL, A1B 3V6, Canada.

^ECorresponding author. Email: d.macphee@usask.ca




Fig. S1. Verification of anti-NUCB2/nesfatin-1 antibody specificity for immunohistochemistry experiments. (A–C) Comparison of non-purified and IgG purified NUCB2/nesfatin-1 antisera specificity in immunohistochemistry experiments. Representative experiments with E12.5 mouse placenta are shown. IgG purified

NUCB2/nesfatin-1 antisera (1:500) and non-purified NUCB2/nesfatin-1 (1:500) antisera were utilized for immunohistochemistry experiments (A and B, respectively) and demonstrated consistently similar NUCB2/nesfatin-1 expression patterns in the mouse placenta. (C) A non-specific, matched IgG control used at the same concentration as the primary antisera displayed no significant background staining within the placenta, but some minor background staining in the uterine muscle. Scale bars = 500 μ m. (D, E) Verification of anti-NUCB2/nesfatin-1 specificity by peptide pre-absorption. (D) NUCB2/nesfatin-1 expression in first trimester (wk 8) chorionic villi using a NUCB2/nesfatin-1 specific antibody. (E) For pre-absorption controls, primary antisera was pre-incubated with 20 μ g NUCB2/nesfatin-1 peptide prior to application to tissue sections during immunohistochemistry experiments. Peptide pre-absorption of the antibody resulted in markedly reduced detection of NUCB2/nesfatin-1 in chorionic villi. Scale bars = 20 μ m. Em = embryo. Pl = placenta. YS = yolk sac, CTB = cytotrophoblast, STB = syncytiotrophoblast, EVS = extravillous space, S = villous stroma.