

10.1071/RD17081\_AC

© CSIRO 2017

Supplementary Material: *Reproduction, Fertility and Development*

## Supplementary Material

### **Markers of protein synthesis are increased in fetal membranes and myometrium after human labour and delivery**

*Stella Liong<sup>A,B</sup> and Martha Lappas<sup>A,B</sup>*

<sup>A</sup>Mercy Perinatal Research Centre, Mercy Hospital for Women, 4th Floor, 163 Studley Road, Heidelberg, Vic. 3084, Australia.

<sup>B</sup>Obstetrics, Nutrition and Endocrinology Group, Department of Obstetrics and Gynaecology, The University of Melbourne, Vic. 3084, Australia.

<sup>C</sup>Corresponding author. Email: [mlappas@unimelb.edu.au](mailto:mlappas@unimelb.edu.au)

**Table S1. Clinical characteristics of the patients in the preterm chorioamnionitis group**

Values represent mean ( $\pm$  s.e.m.) unless otherwise specified. \* $P < 0.05$  vs no infection group (Mann-Whitney test)

	<b>Preterm no labour, no infection (<math>n = 8</math>)</b>	<b>Preterm no labour, infected (<math>n = 8</math>)</b>
<b>Maternal age (years)</b>	34.5 (1.5)	33.9 (1.5)
<b>Pre-pregnancy maternal BMI</b>	23.5 (1.0)	24.0 (1.0)
<b>Gravida</b>	2.8 (0.7)	4.6 (0.8)
<b>Parity</b>	2.1 (0.5)	3.1 (0.7)
<b>Gestational age (weeks)</b>	33.6 (0.7)	28.5 (1.5)*
<b>Fetal gender</b>	3 female; 5 male	6 female; 2 male
<b>Birthweight (g)</b>	1963 (160)	1310 (270)
<b>Delivery mode</b>		
<b>Vaginal delivery</b>	0%	0%
<b>Casearean section</b>	100%	100%
<b>Labour</b>		
<b>No labour</b>	100%	100%
<b>Spontaneous</b>	0%	0%
<b>Duration of labour (min)</b>	N/A	N/A
<b>Membrane rupture</b>		
<b>SROM</b>	0%	0%
<b>PROM</b>	75%	87.5%
<b>ARM</b>	25%	12.5%
<b>Antibiotics</b>	50%	100%
<b>Steroids</b>	75%	100%
<b>Doses</b>	2.2 (0.5)	1.8 (0.2)