

Supplementary Material

Angiogenesis and oxidative stress-related gene variants in recurrent pregnancy loss

Marcela Felix Fortis^A, Lucas Rosa Fraga^{A,B,C,G}, Juliano André Boquett^{A,B}, Thayne Woycinck Kowalski^A, Caroline Gross Dutra^A, Rozana Oliveira Gonçalves^D, Fernanda Sales Luiz Vianna^{A,B,C,E}, Lavinia Schüler-Faccini^{A,B,C} and Maria Teresa Vieira Sanseverino^{A,B,C,F,G}

^APostgraduate Program in Genetics and Molecular Biology, Federal University of Rio Grande do Sul (UFRGS), Porto Alegre, 91501-970, Brazil.

^BNational Institute of Population Medical Genetics (INaGeMP), Porto Alegre, 91501-970, Brazil.

^CNational System for Information on Teratogenic Agents (SIAT), Medical Genetics Service, Hospital de Clínicas de Porto Alegre, Porto Alegre, 90035-903, Brazil.

^DObstetrics, Gynaecology and Human Reproduction Department, Federal University of Bahia, Salvador, 40110-100, Brazil.

^EGenomic Medicine Laboratory, Hospital de Clínicas de Porto Alegre, Porto Alegre, 90035-903, Brazil.

^FSchool of Medicine, Pontifical Catholic University of Rio Grande do Sul, Porto Alegre, 90610-000, Brazil.

^GCorresponding authors. Emails: msanseverino@hcpa.edu.br; lrf.fraga@gmail.com

Table S1. Linkage disequilibrium values (D') of polymorphisms in gene VEGFA

	rs699947	rs1570360	rs2010963	rs3025039
rs699947	–			
rs1570360	0.985 ^A	–		
rs2010963	1.0 ^A	0.972 ^A	–	
rs3025039	0.004	0.195	0.128	–

^AHigh statistical significance (LOD>2)