

## Supplementary Material

### ***In vivo* and *in vitro* strategies to support caprine preantral follicle development after ovarian tissue vitrification**

*N. J. Donfack<sup>A</sup>, K. A. Alves<sup>A</sup>, B. G. Alves<sup>A</sup>, R. M. P. Rocha<sup>A</sup>, J. B. Bruno<sup>A</sup>, L. F. Lima<sup>A</sup>, C. H. Lobo<sup>A</sup>, R. R. Santos<sup>B</sup>, S. F. S. Domingues<sup>B</sup>, M. Bertolini<sup>C,D</sup>, J. Smitz<sup>E</sup> and A. P. R. Rodrigues<sup>A,F</sup>*

<sup>A</sup>Faculty of Veterinary Medicine, Laboratory of Manipulation of Oocytes and Preantral Follicles (LAMOFOPA), State University of Ceará, Av. Paranjana, 1700, Campus do Itaperi. Fortaleza, CEP: 60740 903 – CE – Brasil.


<sup>B</sup>Laboratory of Biology and Medicine of Wild Animals of Amazon, Federal University of Pará-Belem, Av. Universitária, CEP: 68745001 – Castanhal, PA – Brasil.

<sup>C</sup>Federal University of Rio Grande do Sul, Veterinay Faculty, Avenida Bento Gonçalves, 9090, CEP: 91540000 – Porto Alegre, RS – Brasil.

<sup>D</sup>Laboratory of Molecular Biology and development, Fortaleza University, Edson Queiroz, 60811905 – Fortaleza, CE – Brasil

<sup>E</sup>Follicle Biology Laboratory, Center for Reproductive Medicine, UZ Brussel, Laarbeeklaan 101, B-1090 Brussels, Belgium.

<sup>F</sup>Corresponding author. Email: [aprrdriguespapers@gmail.com](mailto:aprrdriguespapers@gmail.com)



**Fig. S1.** Experimental design exhibiting the distribution of ovarian fragments in six different treatments. <sup>n</sup> Represent the total number of fragments in each treatment group.