

## Supplementary Material

### **Localisation and function of glucose transporter GLUT1 in chicken (*Gallus gallus domesticus*) spermatozoa: relationship between ATP production pathways and flagellar motility**

*Rangga Setiawan<sup>A</sup>, Chathura Priyadarshana<sup>A</sup>, Atsushi Tajima<sup>B</sup>, Alexander J. Travis<sup>C</sup> and Atsushi Asano<sup>B,D</sup>*

<sup>A</sup>Graduate School of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan.


<sup>B</sup>Faculty of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan.

<sup>C</sup>Baker Institute for Animal Health, Cornell University, Hungerford Hill Road, Ithaca, NY 14853, USA.

<sup>D</sup>Corresponding author. Email: [asano.atsushi.ft@u.tsukuba.ac.jp](mailto:asano.atsushi.ft@u.tsukuba.ac.jp)

Figure S1. Effect of mitochondrial inhibition on glucose-dependent motility was examined with sperm incubated with antimycin A (AA) under glucose supplementation. <sup>a-c</sup> $P < 0.05$ . (n=5).

Supplementary figure 1



Supplementary figure 1. Effect of mitochondrial inhibition on glucose-dependent motility was examined with sperm incubated with antimycin A (AA) under glucose supplementation.  $a-cP < 0.05$ . (n=5).