

10.1071/RD20215_AC

© CSIRO 2020

Supplementary Material: *Reproduction, Fertility and Development*, 2020, 32(17–18), 1350–1356

Supplementary Material

Wild-type p53-induced phosphatase 1 (*WIP1*) regulates the proliferation of swine Sertoli cells through *P53*

Bingyuan Wang^A, Mingrui Zhang^{A,B}, Jingjing Che^A, Kui Li^A, Yulian Mu^{A,C} and Zhiguo Liu^{A,C}

^AInstitute of Animal Sciences, Chinese Academy of Agricultural Sciences, Beijing 100193, China.

^BCollege of Animal Science and Technology, China Agricultural University, Beijing 100193, China.

^CCorresponding authors. Email: mouyulian@caas.cn; liuzhiguo@caas.cn




Fig. S1. Phosphor-P53/total P53 ratios in the protein levels of NC-si-, WIP1-si-, NC-inhibitor-, miR-16 inhibitor-, NC-mimic-, and miR-16-mimic-treated swine Sertoli cells from western blot bands. The protein expression was normalized to endogenous GAPDH.




Fig. S2. The relative mRNA expression of *p38* determined by RT-qPCR in swine Sertoli cells after loss of WIP1 function by siRNA transfection. Statistical significance was calculated from at least 3 independent experiments. * $P < 0.05$ was indicated.