

Supplementary Material

Sexually transmissible infection testing among pregnant women in the US, 2011–15

Jami S. Leichliter^{A,B}, Laura T. Haderxhanaj^A, Thomas L. Gift^A and Patricia J. Dittus^A

^ADivision of STD Prevention, Centers for Disease Control and Prevention, Atlanta, GA 30329, USA.

^BCorresponding author. Email: jleichliter@cdc.gov

Table S1. STD testing by pregnancy-postpartum status at time of interview, 2011–15
Data are weighted and presented as % (95%CI). CI, confidence interval

	Received chlamydia test (n = 1037)			Received other STD test (n = 1041)		
	Yes	No	P value	Yes	No	P value
Total	49.5 (45.42–53.5)	50.5 (46.5–54.6)	0.008	58.0 (53.6–62.2)	42.0 (37.8–46.4)	0.003
Early pregnancy (1 st trimester)	32.1 (23.4–42.3)	67.9 (57.7–76.7)		38.0 (28.3–48.8)	62.0 (51.2–71.7)	
Late pregnancy (2 nd – 3 rd trimester)	49.9 (42.9–56.9)	50.1 (43.1–57.1)		59.7 (51.8–67.1)	40.3 (33.0–48.2)	
Postpartum (<= 6 months from delivery)	53.5 (47.4–59.6)	46.5 (40.4–52.7)		62.0 (56.2–67.5)	38.0 (32.5–43.8)	