

Supplementary material

Wildfire dynamics and impacts on a tropical Andean oak forest

Mauricio Aguilar-Garavito^{A,B,F}, Paola Isaacs-Cubides^A, J. Sebastian Ruiz-Santacruz^{C,D} and Jordi Cortina-Segarra^E

^AAlexander von Humboldt Biological Resources Research Institute, Avenue Paseo Bolívar 16-20 Bogotá, 111311, Colombia.

^BConservation and Restoration of Ecosystem, Department of Ecology, University of Alicante, San Vicente del Raspeig, 03690, Spain.

^CAsian Demographic Research Institute (ADRI), Shanghai 200444, Shanghai University.

^DCentre of Demographic Studies (CED), Autonomous University of Barcelona, Barcelona, 08007, Spain.

^EDepartment of Ecology and IMEM, University of Alicante, San Vicente del Raspeig, 03690, Spain.

^FCorresponding author. Email: mauricioaguil@gmail.com

Table S1. Correlation analysis to compare the relative surface area burned in the Iguaque mountains (Colombia), with El Niño Ocean Index (ONI).

Test name	Iguaque surface area burned vs. ONI				Iguaque surface area burned vs. Niño				Iguaque surface area burned vs. Niña			
	Value	df	p-value	Sample estimates	Value	df	p-value	Sample estimates	Value	df	p-value	Sample estimates
r Pearson	t = 2.8	346	0.006	r = 0.15	t = 2.8	346	0.005	r = 0.15	t = 1.5	346	0.14	r = 0.08
Rho Spearman	S = 6366831		0.08	rho = 0.09	S = 6318989		0.06	rho = 0.1	S = 6469748		0.14	rho = 0.9
Tau Kendall	z = 1.7		0.08	tau = 0.08	z = 1.9		0.06	tau = 0.09	z = 1.5		0.14	tau = 0.07




Fig. S1. Cross-Correlation Function (CCF) plot for relative surface area burned in the Iguaque mountains (Colombia) and the El Niño Ocean Index (ONI) series.