



# Publications of the Astronomical Society of Australia

VOLUME 19, 2002

© ASTRONOMICAL SOCIETY OF AUSTRALIA 2002

*An international journal of  
astronomy and astrophysics*



**For editorial enquiries and manuscripts, please contact:**

The Editor, PASA,  
ATNF, CSIRO,  
PO Box 76,  
Epping, NSW 1710, Australia  
Telephone: +61 2 9372 4590  
Fax: +61 2 9372 4310  
Email: [Michelle.Storey@atnf.csiro.au](mailto:Michelle.Storey@atnf.csiro.au)



**For general enquiries and subscriptions, please contact:**

CSIRO Publishing  
PO Box 1139 (150 Oxford St)  
Collingwood, Vic. 3066, Australia  
Telephone: +61 3 9662 7666  
Fax: +61 3 9662 7555  
Email: [publishing.pasa@csiro.au](mailto:publishing.pasa@csiro.au)

Published by CSIRO Publishing  
for the Astronomical Society of Australia

[www.publish.csiro.au/journals/pasa](http://www.publish.csiro.au/journals/pasa)

# The Parsec-scale Jets of Extragalactic TeV Gamma-ray Sources

B. G. Piner<sup>1</sup>, P. G. Edwards<sup>2</sup>, S. Fodor<sup>1</sup> and T. A. Rector<sup>3</sup>

<sup>1</sup>Department of Physics and Astronomy, Whittier College, 13406 E. Philadelphia St.,  
Whittier, CA 90608, USA  
gpiner@whittier.edu

<sup>2</sup>ISAS, Yoshinodai 3-1-1, Sagami-hara, Kanagawa 229-8510, Japan

<sup>3</sup>NOAO, 950 North Cherry Avenue, Tucson, AZ 85719, USA

Received 2001 July 31, accepted 2002 March 28

**Abstract:** We present results of VLBI monitoring observations of the blazars detected by ground-based TeV gamma-ray telescopes. Preliminary results have shown that the apparent speeds of the parsec-scale jet features in these sources are predominantly subluminal, and significantly slower than those in the GeV-peaked (EGRET) gamma-ray blazars. Here we add results from additional sources and additional epochs that increase the statistical significance of this comparison. These slower speeds may indicate rapid deceleration of the jets in TeV blazars between the TeV gamma-ray emitting scales and VLBI scales.

**Keywords:** BL Lacertae objects: individual (Mrk 501, 1ES 1959+650, PKS 2155–304, 1ES 2344+514) — galaxies: active — galaxies: jets — radio continuum: galaxies — radiation mechanisms: non-thermal

## 1 Introduction

Detectable fluxes of TeV ( $\sim 1 \times 10^{12}$  eV)  $\gamma$ -rays have been reported for eight AGN: Mrk 421 (Punch et al. 1992), Mrk 501 (Quinn et al. 1996), 1ES 2344+514 (Catanese et al. 1998), PKS 2155–304 (Chadwick et al. 1999), 1ES 1959+650 (Nishiyama et al. 1999), 3C 66A (Neshpor et al. 1998), BL Lac (Neshpor et al. 2001), and 1ES 1426+428 (Horan et al. 2001). At present only Mrk 421 and Mrk 501 have been confirmed by independent groups as TeV gamma-ray sources. Short-timescale variability of the  $\gamma$ -ray emission implies high Doppler factors in the  $\gamma$ -ray emitting regions for at least some of these sources (see the review of Catanese & Weekes 1999). We have analysed multi-epoch VLBI observations of five of these TeV blazars, to determine if high Doppler factors are also evident in the parsec-scale jet structure. We have also recently completed a first-epoch observation of 1ES 1426+428, but have not observed 3C66A, whose high redshift makes it an unlikely TeV source.

## 2 Observations

Mrk 421 and Mrk 501 both have a significant number of archival VLBI observations. For these sources, we have combined this archival VLBI data (from the U.S. Naval Observatory Radio Reference Frame Image Database [e.g. Fey & Charlot 1997] and from the VLBA 2 cm survey [e.g. Kellermann et al. 1998]) with our own VLBA and VSOP data. All archival data was independently imaged and/or model fitted by us, starting from the calibrated ( $u, v$ ) data. The other three sources that we studied (1ES 1959+650, PKS 2155–304, and 1ES 2344+514) have fainter VLBI flux densities ( $\sim 100$  mJy) and had not been previously studied with VLBI. We monitored these sources with the NRAO VLBA at 15 GHz during 1999 and 2000. Some of these observations are also discussed

by, e.g. Edwards et al. (2001), who present more information on the kiloparsec-scale structure of these sources. These observations will be described in detail, including complete sets of images and model fits, in forthcoming publications (Edwards & Piner, in preparation, for Mrk 501; Piner et al., in preparation, for 1ES 1959+650, PKS 2155–304, and 1ES 2344+514).

## 3 Results on Individual Sources

### 3.1 Mrk 421


Analysis of 30 VLBI images of Mrk 421 at 15 epochs is presented by Piner et al. (1999). The apparent speeds of three components were measured, and all were found to be subluminal (see Table 1). These measured speeds differed from apparent superluminal speeds obtained from 3 epochs of VLBI data from the early 1980s by Zhang & Bååth (1990). With the benefit of hindsight, it appears the

**Table 1.** Apparent component speeds

Source	Component <sup>a</sup>	Speed <sup>b</sup> ( $c$ )
Mrk 421	C4	$0.2 \pm 0.3$
	C5	$0.3 \pm 0.1$
	C6	$-0.1 \pm 0.1$
Mrk 501	C1	$0.1 \pm 0.2$
	C2	$0.8 \pm 0.1$
	C3	$0.3 \pm 0.1$
	C4	$0.0 \pm 0.1$
1ES 1959+650	C1	$0.1 \pm 1.1$
	C2	$-0.2 \pm 0.8$
PKS 2155–304	C1	$4.6 \pm 3.8$
1ES 2344+514	C1	$1.3 \pm 0.8$
	C2	$0.5 \pm 0.8$
	C3	$-0.2 \pm 0.8$

<sup>a</sup> Components are numbered C1, C2, ... from the outermost component inward.

<sup>b</sup> For  $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$ .



**Figure 1** 15 GHz VLBA images of IES 2344+514, PKS 2155-304, and IES 1959+650. The uniformly weighted restoring beams have FWHM sizes of  $0.6 \times 0.4$  mas,  $1.0 \times 0.4$  mas, and  $0.7 \times 0.3$  mas, respectively. The lowest contour in each image has been set to 3 times the rms noise level in the image, and is equal to 0.35, 0.62, and 0.45 mJy beam $^{-1}$ , respectively. Successive contours are each a factor of 2 higher. The peak flux densities are 85, 194, and 85 mJy beam $^{-1}$  for the three images, respectively. The axes are labelled in milliarcseconds.

uncertainties in component positions assigned by Zhang & Bååth (1990) were too small. With larger assumed errors, these earlier observations are consistent with those of Piner et al. (1999).

### 3.2 Mrk 501

For Mrk 501, we have analysed a total of 11 epochs of VLBI data: four from the USNO RRFID, six from the VLBA 2 cm survey, and one VSOP observation. A total of four jet components can be identified, with apparent speeds ranging up to  $0.8c$  (see Table 1).

### 3.3 IES 1959+650

We observed IES 1959+650 with the VLBA at 3 epochs at 15 GHz between 2000 March and 2000 July. All three epochs have been imaged, an image from the first epoch is shown in Figure 1. Two jet components whose positions are consistent with no apparent motion were identified at all three epochs (see Table 1).

### 3.4 PKS 2155-304

PKS 2155-304 is a well-known X-ray selected BL Lac object. We observed this source with the VLBA at 3 epochs at 15 GHz between 2000 March and 2000 June. All three epochs have been imaged, and an image from the first epoch is shown. We identified one component with an apparent speed of  $4.6 \pm 3.8c$ . Additional epochs would help to reduce the large error on this measurement.

### 3.5 IES 2344+514

We observed IES 2344+514 four times with the VLBA at 15 GHz between 1999 October and 2000 March. All four epochs have been imaged, an image from the latest epoch is shown in Figure 1. A total of three components were identified by model fitting, at each of the four epochs, with apparent speeds ranging up to  $1.3c$  (see Table 1).

## 4 Discussion

The TeV (‘blue’) blazars appear to have substantially slower apparent speeds than the GeV (‘red’) blazars. All but one of the radio components we have measured are subluminal or only slightly superluminal. We have compared the component speeds we have measured in TeV blazars to the apparent speeds measured in the jets of GeV (EGRET) blazars (Jorstad et al. 2001). A Kolmogorov–Smirnov test shows the two distributions are different with a significance of  $2 \times 10^{-5}$ . These slow apparent speeds must somehow be reconciled with the high Doppler factors demanded by rapid TeV variability.

It is possible for low apparent speeds to be caused by a small viewing angle of the parsec-scale jet. However, this requires very small viewing angles (often  $\ll 1$  degree), and in this case other radio properties indicative of a large Doppler factor such as extreme radio variability and very high apparent brightness temperatures should also be observed, and they are not. We consider the most likely explanation to be that the jet is physically slowing down between the region where TeV  $\gamma$ -rays are generated and the scales mapped by VLBI.

A mechanism to decelerate TeV blazar jets is described by Marscher (1999). If the jet is electron–positron with a sufficiently flat electron index, then most of the energy will be carried in the highest energy electrons that lose energy very efficiently to synchrotron and inverse Compton emission, producing the X-rays and TeV  $\gamma$ -rays, and dumping most of the energy and momentum close to the base of the jet. If deceleration of TeV blazar jets is confirmed by additional work, it may be an important argument for light (electron–positron) jets in these sources.

## Acknowledgments

We thank Ken Kellermann and Alan Fey for providing us with VLBI data on Mrk 421 and Mrk 501. This research has made use of the United States Naval Observatory (USNO) Radio Reference Frame Image Database

(RRFID). The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

## References

- Catanese, M., & Weekes, T. C. 1999, *PASP*, 111, 1193  
 Catanese, M., et al. 1998, *ApJ*, 501, 616  
 Chadwick, P. M., et al. 1999, *ApJ*, 513, 161  
 Edwards, P. G., Piner, B. G., Fodor, S., & Rector, T. A. 2001, in *High Energy Gamma-Ray Astronomy*, AIP Conf. Series 558, eds F. A. Aharonian, & H. J. Völk (San Francisco: ASP), 672  
 Fey, A. L., & Charlot, P. 1997, *ApJS*, 111, 95  
 Horan, D., et al. 2001, in *Proc. XXVII Int. Cosmic Ray Conf.* vol. 7, eds M. Simon, E. Lorenz, & M. Pohl (Hamburg, Germany: IUPAP), 2622  
 Jorstad, S. G., Marscher, A. P., Mattox, J. R., Wehrle, A. E., Bloom, S. D., & Yurchenko, A. V. 2001, *ApJS*, 134, 181  
 Kellermann, K. I., Vermeulen, R. C., Zensus, J. A., & Cohen, M. H. 1998, *AJ*, 115, 1295  
 Marscher, A. P. 1999, *APh*, 11, 19  
 Neshpor, Yu. I., et al. 1998, *AstL*, 24, 134  
 Neshpor, Yu. I., et al. 2001, *ARep*, 45, 249  
 Nishiyama, T., et al. 1999, *Proc. XXVI Int. Cosmic Ray Conf.*, Salt Lake City, eds D. Kieda, M. Salamon, & B. Dingus, 3, 370  
 Piner, B. G., Unwin, S. C., Wehrle, A. E., Edwards, P. G., Fey, A. L., & Kingham, K. A. 1999, *ApJ*, 525, 176  
 Punch, M., et al. 1992, *Nature*, 358, 477  
 Quinn, J., et al. 1996, *ApJ*, 456, L83  
 Zhang, F. J., & Bååth, L. B. 1990, *A&A*, 236, 47