

10.1071/CH08554\_AC

© CSIRO 2009

Accessory Publication: Australian Journal of Publishing, 2009, 62(8), 917–920

**Accessory Publication for**

**Polyethylene Glycol: an Alternative Solvent for Synthesis of Cyclic Carbonate from Vicinal Halohydrin and Carbon Dioxide**

*Jing-Lun Wang, Liang-Nian He,\* Xiao-Yong Dou, Fang Wu*



State Key Laboratory and Institute of Elemento-Organic Chemistry, Nankai University,  
Tianjin 300071, P. R. China



Corresponding author. Email: heln@nankai.edu.cn



**Contents**



<b>1.</b> The $^1\text{H}$ NMR and $^{13}\text{C}$ NMR charts for cyclic carbonates	S2-S6
<b>2.</b> The MS charts for cyclic carbonates	S7-S9
<b>3.</b> The IR chart of cyclohexene carbonate	S9



# 1. The $^1\text{H}$ NMR and $^{13}\text{C}$ NMR charts for cyclic carbonates














## 2. The MS charts for cyclic carbonates




T: + c Full ms [ 29.00-300.00]




*Propylene carbonate*

T: + c Full ms [ 29.00-300.00]



*4-Chloromethyl-1,3-dioxolan-2-one*





*Cyclohexene carbonate*

### 3. The IR chart of cyclohexene carbonate

