© CSIRO 2010

Australian Journal of Chemistry, 2010, 63(2), 315-320

## **Accessory Publication**


## Room Temperature Suzuki Reactions in Aqueous Media under Air by Palladium(II) Complexes with Pyrazole Derived Ligands

Kai Xu, Xin-Qi Hao, Jun-Fang Gong, \* Mao-Ping Song,\* and Yang-Jie Wu


Department of Chemistry, Henan Key Laboratory of Chemical Biology and Organic

Chemistry, Zhengzhou University, Zhengzhou 450052, PR China


Corresponding authors. E-mail addresses: gongjf@zzu.edu.cn or mpsong9350@zzu.edu.cn




<sup>1</sup>H NMR spectrum of complex **2a** 




<sup>13</sup>C NMR spectrum of complex **2a** 




<sup>1</sup>H NMR spectrum of ligand **1b** 







<sup>13</sup>C NMR spectrum of ligand **1b** 




<sup>1</sup>H NMR spectrum of complex **2b** 




<sup>13</sup>C NMR spectrum of complex **2b** 



<sup>1</sup>H NMR spectrum of complex **2b** after the sample solution was placed at rt for 3-4 days



 $^{13}\text{C}$  NMR spectrum of complex 2b after the sample solution was placed at rt for 3-4 days



C-H COSY spectrum of complex  ${\bf 2b}$  after the sample solution was placed at rt for 3-4 days