

Accessory Publication: DNA Interstrand Cross-link Formation by the 1,4-Dioxobutane Abasic Lesion.


*Lirui Guan and Marc M. Greenberg**

Department of Chemistry, Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218


mgreenberg@jhu.edu

Contents:


1. **Supporting Information Figure 1.** ^1H NMR spectrum of $^3\text{H-8} + ^3\text{H-9}$. (S2)
2. **Supporting Information Figure 2.** ^1H NMR spectrum of $^3\text{H-7}$. (S3)
3. **Supporting Information Figure 3.** ^1H NMR spectrum of $^3\text{H-5}$. (S4)
4. **Supporting Information Figure 4.** ^{31}P NMR spectrum of $^3\text{H-5}$. (S5)
5. **Supporting Information Figure 5.** Representative HPLC trace showing native nucleosides and adducts **2-4**.




Supporting Information Figure 2. ^1H NMR spectrum of $^3\text{H-7}$.



Supporting Information Figure 3. ^1H NMR spectrum of $^3\text{H-5}$.



Supporting Information Figure 4. ³¹P NMR spectrum of ³H-5.



Supporting Information Figure 5. Representative HPLC trace showing native nucleosides and adducts **2-4**.