

10.1071/CH11041\_AC


© CSIRO 2011

Australian Journal of Chemistry, 2011, 64(3), 273–278


## **Stabilization of triam(m)inechloridoplatinum complexes by oxidation to Pt(IV)**

Helen L. Daly<sup>1</sup>, Matthew D. Hall<sup>1</sup>, Timothy W. Failes<sup>1</sup>, Mei Zhang<sup>2</sup>, Garry J. Foran<sup>3</sup> and Trevor W. Hambley<sup>1,4</sup>

1. School of Chemistry, The University of Sydney, NSW 2006, Australia
2. Department of Radiation Oncology, Royal Prince Alfred Hospital, Camperdown, NSW 2050, Australia
3. Australian Synchrotron Research Program, c/- ANSTO, Private Mail Bag 1, Menai, N.S.W. 2234, Australia
4. Corresponding author. Email: [trevor.hambley@sydney.edu.au](mailto:trevor.hambley@sydney.edu.au)



**Supplementary Figure S1.** XANES spectrum of a control (untreated) A2780 ovarian cancer cell line pellet, demonstrating the lack of an edge due to Pt.



**Supplementary Figure S2.** Example XANES spectra of Pt(II) and Pt(IV) complexes, showing the difference in peak heights for the two oxidation states, and the parameters **a** and **b** used in determining the ratio **a/b**. Note the greater degree of absorption in the white line height of Pt(IV) complexes.