

SUPPLEMENTARY MATERIAL

An efficient synthesis of 2-imidazolines in the presence of molecular iodine under ultrasound irradiation

Guo-Feng Chen,^{A,D} Hong-Yang Li,^A Nan Xiao,^A Bao-Hua Chen,^B Ya-Li Song,^B
Ji-Tai Li,^A and Zhi-Wei Li^C

^A Key Laboratory of Chemical Biology of Hebei Province, College of Chemistry and Environmental Science, Hebei University, Baoding 071002, P.R. China

^B College of Pharmaceutical Sciences, Hebei University, Baoding 071002, P.R. China

^C College of Chemical and Pharmaceutical Engineering, Hebei University of Science and Technology, Shijiazhuang 050018, P.R. China

^D Corresponding author. E-mail: chenguofeng@hbu.cn; Fax: 86-312-5079628




Figure S1. ¹H NMR (600 MHz, CDCl₃) spectrum of 2-(3,4-dichlorophenyl)imidazoline (**3h**).




Figure S2. ¹³C NMR (150 MHz, CDCl₃) spectrum of 2-(3,4-dichlorophenyl)imidazoline (**3h**).




Figure S3. ¹H NMR (600 MHz, CDCl₃) spectrum of 2-(2-furanyl)imidazoline (**3k**)






Figure S4. ¹³C NMR (150 MHz, CDCl₃) spectrum of 2-(2-furanyl)imidazoline (**3k**)







Figure S8. ¹³C NMR (150 MHz, CDCl₃) spectrum of 2-(4-fluorophenyl)imidazoline (**3n**)

