

Supplementary Material

Sequential detection of CN^- and HSO_4^- anions in aqueous environment utilizing a new colorimetric azoimine receptor: mimicking logic gate behavior and a security keypad lock

Khatereh Rezaeian, Hamid Khanmohammadi* and Nafiseh Shabani

Department of Chemistry, Faculty of Science, Arak University, Arak 38156-8-8349, Iran

*Email: h-khanmohammadi@araku.ac.ir

Page No.	Contents
S2	IR spectrum of 1
S3	^1H NMR spectrum of 1
S4	Job's plot of 1 with CN^- in 3:2 DMSO/water
S5	Effect of pH on the UV-Vis spectra of 1 - CN^- complex at 470 nm
S6	Benesi-Hildebrand plot of sensor 1 - CN^- binding with HSO_4^- anion associated with absorbance change at 470 nm
S7	UV-Vis titration of 1 (3×10^{-5} mol L^{-1}) in 3:2 DMSO-water with incremental addition of OH^- (0-5 equiv.) in water.
S8	^1H NMR spectrum of 1 + CN^-
S9	^1H NMR spectrum of 1 + CN^- + HSO_4^-




Figure S1. IR spectrum of 1




Figure S2. ^1H NMR spectrum of **1**




Figure S3. Job's plot of 1 with CN⁻




Figure S4. Effect of pH on the UV-Vis spectra of 1- CN^- complex at 470 nm




Figure S5. Benesi-Hildebrand plot of sensor 1- CN^- binding with HSO_4^- anion




Figure S6. UV-Vis titration of **1** (3×10^{-5} mol L⁻¹) in 3:2 DMSO-water with NaOH

