

Use of productivity-defined indicators to assess exposure of grassland-based livestock systems to climate change and variability

Marion Sautier^{A,B,C}, Michel Duru^A, and Roger Martin-Clouaire^B

^AINRA, UMR 1248 AGIR, BP 52627, F-31326 Castanet Tolosan, France.

^BINRA, UR875 MIAT, BP 52627, F-31326 Castanet Tolosan, France.

^CCorresponding author. Emails: marion.sautier@gmail.com; marion.sautier@toulouse.inra.fr

Table A1: Table of correspondence between rank (by number) of forage-year clusters by decreasing relative frequency in the past and rank (by letter) of forage-year clusters by decreasing average annual herbage balance in the past, at the three study sites.

Cluster rank by decreasing relative frequency in the past	Cluster rank by decreasing average annual herbage balance in the past and relative frequency of the cluster in the past (%)						
	Aulus			Saint Girons		Toulouse	
Rank	Rank	%	Rank	%	Rank	%	
1	H	30	F	37	B	32	
2	B	23	C	23	C	26	
3	F	20	G	17	F	13	
4	D	13	H	13	A	10	
5	A	8	D	10	E	10	
6	C	3	A	0	D	3	
7	E	3	B	0	G	3	
8	G	0	E	0	H	3	