


**Accessory publication****Source analysis of perfluorocarboxylates in Tokyo Bay during dry weather and wet weather using sewage markers**

Michio Murakami,<sup>A</sup> Chiaki Morita,<sup>B</sup> Takuya Morimoto<sup>B</sup> and Hideshige Takada<sup>B,C</sup>

<sup>A</sup>‘Wisdom of Water’ (Suntory), Corporate Sponsored Research Program, Organization for Interdisciplinary Research Projects, The University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-8656, Japan.

<sup>B</sup>Laboratory of Organic Geochemistry (LOG), Institute of Symbiotic Science and Technology, Tokyo University of Agriculture and Technology, Fuchu, Tokyo 183-8509, Japan.

<sup>C</sup>Corresponding author. Email: shige@cc.tuat.ac.jp



**Fig. A1.** Comparison of principal component scores between dry weather (16 May) and wet weather (15 Oct.).

**Table A1. Reproducibility and recovery rates of PFCAs and PPCPs**

Values in parentheses come from results of PFCA standards spiked into the wastewater samples

	Relative standard deviation	Recovery rates
PFHpA	21%	85%
PFOA	14%	103%
PFNA	19%	92%
PFDA	16%	100%
PFUA	5%	87%
PFDoDA	(13%)	114%
PFTriDA	(9%)	95%
PFTeDA	(14%)	87%
Crotamiton	1%	109%
Caffeine	4%	114%