

10.1071/FP15216_AC

© CSIRO 2016

Supplementary Material: *Functional Plant Biology*, 2016, 43(1), 62–74.

Supplementary Material

Root biomass in the upper layer of the soil profile is related to the stomatal response of wheat as the soil dries

Renu Saradadevi^{A,B,F}, *Helen Bramley*^{B,C}, *Jairo A. Palta*^{A,D}, *Everard Edwards*^E and *Kadambot H. M. Siddique*^B

^ASchool of Plant Biology, The University of Western Australia, Perth, WA 6009, Australia.

^BThe UWA Institute of Agriculture, The University of Western Australia, Perth, WA 6009, Australia.

^CPresent address: Plant Breeding Institute, Faculty of Agriculture and Environment, The University of Sydney, 12656 Newell Highway, Narrabri, NSW 2390, Australia.

^DCSIRO Agriculture Flagship, Private Bag No. 5, Wembley, WA 6913, Australia.

^ECSIRO Agriculture Flagship, PMB2, Glen Osmond, SA 5064, Australia.

^FCorresponding author. Email: renusaradadevi@gmail.com




Fig. S1. Changes in leaf water potential (Ψ_{leaf}) with the changes in whole pot soil water content (%) in IGW-3262 (solid circle) and Drysdale (open circle). Each data point represents one pot (replicate). R^2 values are 0.33 (IGW-3262) and 0.40 (Drysdale).