Supplementary Material

Evidence that melatonin promotes soybean seedlings growth from low-temperature stress by mediating plant mineral elements and genes involved in the antioxidant pathway

George Bawa^{A,B,C}, Lingyang Feng^{A,B,C}, Jianyi Shi^{A,B,C}, Guopeng Chen^{A,B,C}, Yajiao Cheng^{A,B,C}, Jie Luo^D, Weishu Wu^D, Bancy Ngoke^D, Ping Cheng^{A,B,C}, Zhongqin Tang^{A,B,C}, Tian Pu^{A,B,C}, Jiang Liu^{A,B,C}, Weiguo Liu^{A,B,C}, Taiwen Yong^{A,B,C}, Junbo Du^{A,B,C}, Wenyu Yang^{A,B,C} and Xiaochun Wang^{A,B,C,E}

^ASichuan Engineering Research Centre for Crop Strip Intercropping System, Sichuan Agricultural University, Chengdu 611130, China.

^BKey Laboratory of Crop Ecophysiology and Farming System in Southwest China (Ministry of Agriculture), Sichuan Agricultural University, Chengdu 611130, China.

^cCollege of Agronomy, Sichuan Agricultural University, 211-Huimin Road, Wenjiang District, Chengdu 611130, China.

^DCollege of Veterinary Medicine, 211-Huimin Road, Wenjiang District, Chengdu 611130, China.

^ECorresponding author. Email: xchwang@sicau.edu.cn



Fig. S1. Effect of MT application at different concentrations 1, 10, and 50 μ mol L⁻¹ on soybean seedlings growth.