

Supplementary Material

The distribution and abundance of electrosensory pores in two benthic sharks: a comparison of the wobbegong shark *Orectolobus maculatus* and the angel shark *Squatina australis*

Channing A. Egeberg^A, Ryan M. Kempster^{A,C}, Susan M. Theiss^B, Nathan S. Hart^A and Shaun P. Collin^A

^AThe UWA Oceans Institute and the School of Animal Biology, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia.

^BFisheries Queensland, Department of Agriculture, Fisheries and Forestry, GPO Box 46, Brisbane, Queensland 4001, Australia.

^CCorresponding author. Email: ryankempster@supportoursharks.com

Table S1. Mean total abundance of electrosensory pores in *Orectolobus maculatus* and *Squatina australis*

Data are presented as the mean number of pores \pm one standard deviation and the range. ‘*’ denotes a statistically significant difference in pore abundance between species (two-sample *t*-test with a *P*-value \leq 0.001)

Genus	Species	<i>n</i>	Ventral*	Dorsal*	Total*
<i>Squatina</i>	<i>australis</i>	8	76.4 \pm 4.9	174.9 \pm 8.8	251.3 \pm 9.8
			66 - 82	164 - 192	237 - 269
<i>Orectolobus</i>	<i>maculatus</i>	4	45.3 \pm 2.9	526.8 \pm 9.4	572.0 \pm 7.7
			41 - 47	515 - 538	561 - 579