

Supplementary material

Historical analysis of blue marlin (*Makaira nigricans* Lacepède, 1802) catches by the pelagic longline fleet in the eastern Pacific Ocean

Emigdio Marín-Enríquez^{A,D}, Leonardo A. Abitia-Cárdenas^B, Xchel G. Moreno-Sánchez^B and Jorge S. Ramírez-Pérez^C

^ACONACYT-Facultad de Ciencias del Mar, Universidad Autónoma de Sinaloa,
Paseo Claussen s/n, Colonia Los Pinos 82000, Mazatlán, Sinaloa, México.

^BInstituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas,
Avenida IPN s/n, 23096, La Paz, Baja California Sur, México.

^CFacultad de Ciencias del Mar, Universidad Autónoma de Sinaloa, Paseo Claussen s/n,
Colonia Los Pinos 82000, Mazatlán, Sinaloa, México.

^DCorresponding author. Email: emarin@conacyt.mx




Fig. S1. Residual (diagnostic) plots of the final generalised additive mixed model fitted to the blue marlin catch-per-unit-effort (CPUE) time series. (a) Normal Q-Q plot, (b) frequency distribution, (c) residuals v. fitted values, (d) residuals v. month, (e) residuals v. year, (f) residuals v. Oceanic Niño index (ONI) and (g) residuals v. Pacific decadal oscillation (PDO).




Fig. S2. Time series of swordfish (*Xiphias gladius*) catch-per-unit-effort (CPUE) by the industrial pelagic longline fleet in the eastern Pacific Ocean, for the period of January 1959–December 2017.