



Cover

Huveneers *et al.* (p. 765) provided the reproductive parameter estimates of wobbegongs necessary for population modelling. They found that the three species of wobbegongs occurring in NSW have a synchronous triennial reproductive cycle and provided maturity, maternity, and fecundity estimates essential for adequate fisheries management and species conservation assessments.



Anger *et al.* (p. 743) studied the early life history of an endemic Jamaican stream crab. As an adaptation to unpredictable planktonic food production in the breeding habitat, *Sesarma meridies* produces large eggs with enhanced lipid reserves, allowing for food-independent larval development. These findings aid the understanding of evolutionary invasions of limnic and terrestrial environments.

CONTENTS

Does adding wood to agricultural streams enhance biodiversity? An experimental approach <i>R. E. Lester, W. Wright and M. Jones-Lennon</i>	687
Influence of upwelling on movement of southern bluefin tuna (<i>Thunnus maccoyii</i>) in the Great Australian Bight <i>J. Willis and A. J. Hobday</i>	699
Processes driving circulation, exchange and flushing within intermittently closing and opening lakes and lagoons <i>E. Gale, C. Pattiaratchi and R. Ranasinghe</i>	709
Investigating the association between pelagic fish and dimethylsulfoniopropionate in a natural coral reef system <i>J. L. DeBose and G. A. Nevitt</i>	720
Contribution of adult aquatic insects to riparian prey availability along tropical forest streams <i>E. K. W. Chan, Y. Zhang and D. Dudgeon</i>	725
Evidence of genetic subdivision among populations of blacklip abalone (<i>Haliotis rubra</i> Leach) in Tasmania <i>N. Temby, K. Miller and C. Mundy</i>	733
Adaptive traits in ecology, reproduction and early life history of <i>Sesarma meridies</i> , an endemic stream crab from Jamaica <i>K. Anger, G. Torres and U. Nettelmann</i>	743
Spatial distribution of commercial dredge fishing effort: application to survey design and the spatial management of a patchily distributed benthic bivalve species <i>J. J. Harrington, J. M. Semmens and M. Haddon</i>	756
Reproductive synchrony of three sympatric species of wobbegong shark (genus <i>Orectolobus</i>) in New South Wales, Australia: reproductive parameter estimates necessary for population modelling <i>C. Huveneers, T. I. Walker, N. M. Otway and R. G. Harcourt</i>	765
Flushing time of solutes and pollutants in the central Great Barrier Reef lagoon, Australia <i>Y. Wang, P. V. Ridd, M. L. Heron, T. C. Stieglitz and A. R. Orpin</i>	778

EARLY ALERT Sign-up at www.publish.csiro.au/journals/mfr for our electronic early alert and receive the next table of contents weeks in advance of the print version