

**Lewin Honeyeater.**—I should be interested to have a few field notes on the Lewin Honeyeater (*Meliphaga lewini*). As I know the bird in coastal Queensland, the colouration is much as in Cayley's illustration (*What Bird is That?*). But at Stanthorpe tableland some while back, and again last year at Ben Bullen in New South Wales, I had what I thought were "Lewin's" under observation for some time; and the birds differed greatly in colour from those on the coast. The inland birds, with bright olive-green bodies, had black heads and throat—the black of throat clearly and sharply defined from the green of body. Wings and tail were also olive-green, and my impression was of a very handsome bird larger than the Lewin Honeyeater is generally, and with a more pleasing, though not so sustained, note. The ears were very yellow. In Queensland orange orchards the Lewin Honeyeater can be very destructive. The bird manages to scoop out oranges and mandarins from below, leaving the empty skin hanging from the tree.—L. M. MAYO, R.A.O.U., Brisbane, Queensland.

---

**Erratum.**—In my paper, "Bird Notes from Rockingham Bay" (*The Emu*, Vol. XXXIII, p. 242), I stated incorrectly that Campbell and Barnard recorded *Drymodes superciliaris* in the locality in question during their 1916 expedition. It was *Pacilodryas superciliosa*, the White-browed Robin, which my predecessors observed, not the Scrub-Robin, as stated.—A. J. MARSHALL, R.A.O.U., Sydney, N.S.W.