NFECTIOUS DISEASES

MEASLES EPIDEMIC, SOUTH COAST NSW

Greg Sam and Paul Van Buynder South East NSW Public Health Unit Shanti Raman National Centre for Epidemiology and Population Health John Skinner North Sydney Public Health Unit

Between August 20 and November 8, 1994, 214 measles notifications were received by the South Eastern Public Health Unit from the South Coast District of NSW. The epidemic was centred in the Bega Valley area, with 43 per cent of known cases occurring in students at Bega High School. The outbreak peaked during the week beginning September 25, which was the first week of the Term 3 school holidays (Figure 2).

Measles cases were defined as people having an illness characterised by:

- a generalised maculopapular rash resembling measles
- a high fever (>38C)
- one or more of the following: cough, coryza, conjunctivitis or Koplik spots.

NSW refugee screening

Continued from page 135

Improved data collection on the proportion of tuberculinpositive children screened who receive preventive therapy is recommended.

The RPR screening test for syphilis used pre-migration has been shown here to miss 80 per cent of refugees with evidence of past treponemal disease, as previously reported². The TPHA test is more sensitive in this situation as it usually remains reactive lifelong. Some positive TPHA results among refugees may be due to other treponemal diseases such as yaws or pinta, which are indistinguishable from syphilis on serological grounds. However, those who do have latent syphilis are at risk of progression to tertiary disease. The detection of these cases, with assessment for therapy, is therefore important in personal and public health terms.

Hepatitis B tests are not performed pre-migration. Because serum is collected for syphilis testing from those 15 years and older, it was decided to limit hepatitis B tests to the same age group. The prevalence of the carrier state is, as expected, high. Testing allows appropriate advice to be given to carriers and their families; further serological testing of adult contacts determines their need for vaccination. Catch-up immunisation against hepatitis B is offered to all children under 15 years old not previously vaccinated. This is done without prior serological testing, based on cost calculations and the known safety of the vaccine⁷.

Immunisation status is well documented for arriving refugees, and coverage for children is high. The screening visit is an excellent opportunity for catch-up vaccination in adults and children. For example, HIB vaccine is not given in the camps, and most adults have never received Sabin.

Case notifications

On September 22, two days before the end of the school term, a South Coast general practitioner notified the South Eastern NSW PHU of two cases of clinically diagnosed measles in Bega High School students. Follow-up with the school identified a high level of measles-related absenteeism in the preceding two weeks.

Potential sources for further case identification were contacted throughout the area, including hospitals, GPs, schools, and preschool and child care centres. The need for prompt notification of cases was stressed.

A high attack rate among high school-aged children was identified and questionnaires were distributed to the two local high schools to ascertain the degree of under-reporting of cases.

Serological confirmation was obtained from 35 cases.

Immunisation campaign

The need for measles vaccination for all unimmunised children as well as a recommendation for a booster dose for children aged 10-17 years who had been previously vaccinated was highlighted in school letters and through the local media.

The current screening program is routinely offered to refugees from South East Asia and Central and South America only. While small numbers from other areas are starting to be seen, routine screening of refugees from Africa, Eastern Europe and the Middle East should be introduced, based on known rates of tuberculosis in the country of origin².

In summary, most of the major recommendations of the 1991 review of refugee screening^a have been implemented, including relocation to a Chest Clinic, increased emphasis on diseases of public health significance, introduction of hepatitis B testing and upgraded data collection. In addition, the program has been streamlined: those with personal health problems are referred to general practitioners as much as possible; there is rapid feedback of results; and routine treatment for intestinal parasites is no longer given, as nearly all South East Asian refugees have received treatment with pyrantel just before their departure for Australia.

^{1.} Reid J, Goldstein GB, Keo L. An evaluation of refugee medical screening in NSW. November 1985. Report to the NSW Department of Health, Western Metropolitan Region. NSW Health Department, Sydney, 1985.

Sydney, 1985. 2. Bek M, Levy M. A review of the NSW refugee medical screening program. June 1991. NSW Health Department, State Health Publication No. (EHSEB) 92-12, Sydney, 1991.

Westley-Wise V, Levy M, Lonie C, McAnulty J, Winks M, Stewart G. Controlling Tuberculosis in New South Wales. March 1993. NSW Health Department, Sydney, 1993.

McAnulty J, Levy M, Rubin G. Eliminating tuberculosis:
McAnulty J, Levy M, Rubin G. Eliminating tuberculosis:
Where is the strategy? NSW Public Health Bulletin 1992; 3(4):39-40.
Westley-Wise V, Levy M, Winks M. Review of tuberculosis services in New South Wales. December 1992. NSW Health Department, Sydney,1992.

Alperstein G, Fett MJ, Reznik R, Thomas M, Senthil M. The prevalence of tuberculosis infection among Year 8 schoolchildren in inner Sydney in 1992. *Med J Aust* 1994; 160:197-201.

^{7.} US Department of Health and Human Services. Protection against viral hepatitis. Recommendations of the Immunization Practices

Community Health immunisation clinics were made available after hours and on weekends during the peak of the epidemic. The clinics were well attended and more than 1,200 children were vaccinated over a four-week period. GPs reported large attendances for measles vaccination.

To date, only two notifications of adverse reactions to MMR vaccine have been notified. Both cases were reported as persistent screaming and high fever, and recovered without hospitalisation.

Epidemic demography

During the outbreak, cases were reported from almost all population centres in the South Coast District (Table 8). The epidemic began in, and was initially confined to, the Bega Valley area, but as time progressed cases were reported further north in the Eurobodalla shire. Overall, 76 per cent (n=163) of cases were reported from the Bega Valley area, and 18 per cent (n=39) from Eurobodalla. Secondary cases were subsequently reported in the Monaro and Southern Tablelands Districts.

For the six weeks from the onset of the outbreak, cases occurred mainly in high school-aged children. As the epidemic progressed, the proportion of cases in primary school-aged children increased. The age and sex





TABLE 8

GEOGRAPHIC DISTRIBUTION OF MEASLES CASES NOTIFIED TO THE SOUTH EASTERN PHU, AUGUST 20-NOVEMBER 8, 1994

0.000	Residential area	Number	Percentage	
00000	Bega	123	57.5	
	Batemans Bay/			
	Moruya/Narooma	36	16.8	
	Eden	22	10.3	
	Pambula/Merimbula	18	8.4	
	Queanbeyan*	6	2.8	
	Bodalla	3	1.4	
	Crookwell*	3	1.4	
	Cooma*	2	0.9	
	Yass*	1	0.5	
	TOTAL	214	100.0	

* not in South Coast District

distribution is shown in Figure 3. There were similar proportions of males and females in each age group.

Hospitalisations

During the epidemic there were 29 hospital admissions. The main reasons for hospital admission were pneumonia/respiratory infection and dehydration. There have been no deaths to date.

Immunisation status of cases

Reporting of the immunisation status of cases during the outbreak was based on parental recall, with 22 per cent of high school-aged children reported as immunised, 36 per cent reported as not immunised and 42 per cent unknown. The large proportion of cases reported as "immunisation status unknown" reflects the inadequacy of parental recall as a measure of immunisation status of children¹. Data based on questionnaires distributed to high school students will be reported in a follow-up report.

Discussion

The characteristics of this epidemic are similar to other recently described measles outbreaks²³ with the highest attack rate occurring in high school students in the Bega Valley area. In the absence of reliable data on the immunisation status of high school students at the time of reporting, the role of vaccine efficacy in determining susceptibility to contracting measles cannot be examined in this report.

Under-reporting was found to occur at all stages of the epidemic despite continued active surveillance. Many GPs were unaware of the requirement to notify measles immediately and some were not aware of the requirement to notify. The importance of prompt notifications for the success of school exclusion policies was also poorly understood. Of eight cases diagnosed in an emergency department, none was notified by the hospital. The delay in notification of early cases until the onset of the school holiday period also made control measures more difficult.

The rapid deployment of immunisation clinics during the early phase of the epidemic by Community Health staff

Continued on page 138 ►

TABLE 9								* First	t diagno	sis	[5. 01/01/9	4-31/06/94					
SURVEILLANCE (JANUARY-NOVE (Diagnoses from	OF NON-NO MBER 1994 sexual hea	TIFIABL	E SEXU tres un!	ALLY T	RANSM erwise	IITTED DISEASES stated in footno	2. 01/0 2. 01/0 3. 01/0 4. 01/0 5. 01/0	1/94-31/ 1/94-31/ 1/94-31/ 1/94-31/ 1/94-31/	04/94 01/94 10/94 03/94 /09/94	8. No SHC in Region 9. Laboratory and SHC data 01/01/94-31/08/94 10. No data yet received for 1994								
AHS ⁺ Infection		CSA'	SSA ²	ESA ³	SWS⁴	WSA4 + WEN	NSA ³	CCA ³	ILL ³	HUN ⁶	NC ⁷	ND ³	WNS ⁵	CW ⁸	SW ⁹	SE ¹⁰	Total	
Chlamydia	Male	1	-	90	2	6	2	1	5	8	-	5	7	_	3	-	130	
trachomatis	Female	1	-	63	5	7	1	3	4	14	1	18	23	_	8	_	148	
	Total	2	-	153	7	13	3	4	9	22	1	23	30	-	11	_	278	
Donovanosis	Male	-	-	-	-	_	_	-	-	_	-	-	_	_	_	_	_	
	Female	-	-	-	-	-	_	-	` -	_	_	_	_	_	_	-	_	
	Total	-	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-	
*Genital herpes	Male	3	1	299	3	12	10	12	-	15	7	3	1	-	5	-	371	
	Female	4	3	186	5	9	9	12	13	15	9	14	6	-	7	-	292	
	Total	7	4	485	8	21	19	24	13	30	16	17	7	-	12	-	663	
*Genital warts	Male	11	6	778	69	74	25	40	75	75	33	8	6	_	8	_	1.208	
	Female	8	6	317	32	37	22	23	28	30	11	27	19	-	10	_	570	
	Total	19	12	1,095	101	111	47	63	103	105	44	35	25	-	18	-	1,778	
Nongonococcal	Male	3	1	584	23	55	17	34	25	43	13	10	7	-	4	-	819	
urethritis	Female	-	-	-	-	3	3	-	-	-	-	-	2	-	2	_	10	
	Total	3	1	584	23	58	20	34	25	43	13	10	9	-	6	-	829	
Lymphogranulor	na Male	-	-	-	-	-	-	_	_	_	-	-	-	-	-	-	-	
venereum	Female	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	_	
	Total	-	-	-	-	-	-	-	-	-	-	-	-	-	S.S	-	-	

† Data from Public Health Unit area of the clinic not the patient.

Infectious diseases

Continued from page 137

enabled large numbers of children throughout a large area to be vaccinated in a relatively short period.

Further clusters of cases continue to occur on the South Coast and a full report will follow.

 Goldstein KP, Kviz FJ, Daum RS. Accuracy of immunization histories provided by adults accompanying preschool children to a pediatric emergency department. JAMA 1993; 270(18):2190-4.
Merianos A, Miller NC, Patel M. Control of a community outbreak of measles which started in a poorly immunised high school population. Aust J Public Health 1993; 17:231-236.
Cheah D, Lane JM, Passaris I. Measles vaccine efficacy study in a Canberra high school: a study following a measles outbreak. J Paed Child Health 1993; 29:455-458.

EDITORIAL COMMENT

In recent years, measles outbreaks have occurred in discrete geographic areas in NSW. The outbreak reported above is an example, and outbreaks have also occurred recently in the Lower and Mid North Coast and Clarence Districts. In 1993, measles outbreaks occurred in the Western Sydney, South Western Sydney and Wentworth Areas. The majority of cases in the outbreaks were in the 10-16 year age group. Immunisation is the only proven control method, and measles immunisation is recommended at the age of 12 months. Surveys by Public Health Units indicate that immunisation rates are in the range 75-85 per cent (short of the 95 per cent required to prevent outbreaks). The NHMRC has recently recommended that rubella immunisation (which previously targeted adolescent girls) should be extended, and that both girls and boys should receive measles-mumps-rubella vaccine in Year 7. Implementation of this has begun in NSW. Parents are encouraged to ensure that children are adequately immunised.

HYDATID DISEASE

John Walker

Department of Parasitology Centre for Infectious Diseases and Microbiology Westmead Hospital

Human hydatid disease results from infection with the larval stages of tapeworms of the genus Echinococcus. In Australia the species involved is E. granulosus, which has a cycle of development usually involving dogs and sheep. The adult worms occur in dogs and the larval stages, hydatid cysts, are found in herbivores which become infected by eating eggs passed in dog faeces. Human infection occurs in the same manner, by the ingestion of eggs, not by eating raw meat or offal containing cysts. In humans the most common sites for development of hydatid cysts are the liver and lungs, but cysts may be found in virtually any organ. Factors such as the site of cysts, their size and number and whether they leak or rupture, determine if infected individuals become symptomatic. Surgery generally is the only reliable method of treatment of hydatid disease but this may not be possible when cysts occur in inaccessible sites. Obviously, prevention of infection is preferable to surgery, especially as there is a significant rate of recurrence of the disease.

Because hydatid disease is a zoonosis usually involving dogs and sheep in Australia, most cases occur in people resident in country regions, especially the relatively cool and moist highlands of south-eastern Australia. This infection was previously highly prevalent in Tasmania, with rates of around 27/100,00, but a control campaign begun in the 1960s has virtually eliminated the parasite from that State. Attempts to mount similar campaigns in mainland States have never been successful and rates of infection as high as 32/100,000 have been reported in some regions, particularly in the Southern Highlands of NSW.

Experience in New Zealand, Tasmania and many other parts of the world has shown that hydatid disease can be controlled with public health measures. A symposium on hydatid disease, its clinical management, epidemiology and

TABLE 10

INFECTIOUS DISEASE NOTIFICATIONS FOR 1994 BY SELECTED MONTH OF ONSET FOR NOTIFICATIONS RECEIVED BY NOVEMBER 30, 1994

Condition	Aug	Sep	Oct	Nov	Total
Adverse event	_				-
after immunisation	3	2	-	-	5
AIDS	32	26	13	3	14
Arboviral infection	4	4	6	5	19
Brucellosis	2	1	-	-	3
Foodborne Illness (NOS)	5	/	5	1	18
Gastroenteritis (instit.)	38	9	11	3	61
Gonorrhoea	21	19	14	4	64
H influenzae epiglottitis	-	2	1	-	3
H influenzae meningitis	2	1	1	1	5
H influenzae septicaemia	1	1	-	-	2
H influenzae infection (NOS)	-	-	1	-	1
Hepatitis A – acute viral	43	31	29	21	124
Hepatitis B – acute viral	10	2	4	-	16
Hepatitis B – chronic/carrier	49	20	12	4	85
Hepatitis B – unspecified	346	348	381	128	1,203
Hepatitis C – acute viral	6	2	-	-	8
Hepatitis C – unspecified	872	740	570	197	2,379
Hepatitis D – unspecified	-	1	1	-	2
Hepatitis, acute viral (NOS)	1	1	-	-	2
HIV infection	35	30	28	21	114
Hydatid disease	2	-	3	-	5
Legionnaires' disease	4	3	1	-	8
Leprosy	1	-	-	-	1
Leptospirosis	2	-	-	-	2
Listeriosis	1	1	1	-	3
Malaria	17	10	8	4	39
Measles	40	209	264	223	736
Meningococcal meningitis	18	4	11	3	36
Meningococcal septicaemia	7	4	5	1	17
Meningococcal infection (NOS)	2	3	4	2	11
Mumps	1	2	4	-	7
Mycobacterial atypical	25	23	7	2	57
Mycobacterial tuberculosis	23	27	13	4	67
Mycobacterial infection (NOS)	8	21	15	5	49
Pertussis	124	115	97	27	363
Q fever	14	15	13	1	43
Rubella	5	6	3	1	15
Salmonella (NOS)	39	45	70	25	179
Salmonella bovis morbificans	1	1	-	-	2
Salmonella typhimurium	22	6	_	_	28
Syphilis	108	76	46	12	242
Tetanus	_	-	1	-	1
Typhoid and paratyphoid	3	3	-	-	6
Tatal	1.012	4.024	1.042	600	C 105
lotal	1,943	1,821	1,643	698	6,105

control will be held at Westmead Hospital, Sydney, on Friday, March 3, 1995. For more information, phone (02) 633 7191 or fax (02)893 8659.

SYPHILIS

In the October 1994 edition of the *NSW Public Health Bulletin* it was noted that the notification rate for syphilis was higher this year than last year. The notification rate for the period to the end of November was 11.2 in 1993 and 14.0 in 1994. Consultations with staff from Public Health Units, sexual health clinics and laboratories confirm the increase is probably the result of improved surveillance. There has been no increase in reported cases of newly acquired syphilis.

PERTUSSIS ON THE NORTH COAST

North Coast PHU reports larger than average numbers of pertussis in the Casino area in recent months. The notification rate for the North Coast in 1994 is 56.8/100,000

TABLE 11

SUMMARY OF NSW INFECTIOUS DISEASE NOTIFICATIONS NOVEMBER 1994

Condition	Num	ber of c	ases notified							
	Peri	iod	Cumul	ative						
	Nov 1993	Nov 1994	Nov 1993	Nov 1994						
Adverse reaction AIDS	23	- 3	23 338	29 323						
Arboviral infection Brucellosis	16	5	643 4	367						
Cholera	-	-	-	-						
Foodborne illness (NOS)	13	- 1	120	149						
Gastroenteritis (instit.)	73	3	406	220						
Gonorrhoea H influenzae epiglottitis	32	4	329	2/5						
H influenzae B – meningitis	1	1	53	15						
H influenzae B – septicaemia	-	-	23	11						
Hinfluenzae Infection (NOS) Hepatitis A	42	21	559	453						
Hepatitis B	433	132	3,762	3,851						
Hepatitis C Hepatitis D	790	197	6,051	7,483						
Hepatitis, acute viral (NOS)	-	-	6	6						
HIV infection	41	21	513	399						
Hydatid disease	3		4 65	15						
Leprosy	-	-	3	3						
Leptospirosis	1	-	15	12						
Malaria	7	4	158	167						
Measles	588	223	2,070	1,088						
Meningococcal meningitis	12	3	90	72						
Meningococcal infection (NOS)	_	2	11	19						
Mumps	3	-	9	10						
Mycobacterial – atypical	32	4	382	357						
Mycobacterial infection (NOS)	8	5	47	93						
Pertussis	315	27	1,304	1,216						
Poliomyelitis	_		_	-						
Q fever	38	1	376	212						
Rubella Salmonella infection (NOS)	129	1 25	782 880	914						
Syphilis	94	12	707	865						
Tetanus Turbaid and pareturbaid	-	-	5	3						
Typhoid and paratyphoid Typhus	1	-	- 25	- 25						
Viral haemorrhagic fevers	-	-	-	-						
I ENOW IEVEL	STATISTICS.			No. of Concession, Name						

compared to 19.6/100,000 for NSW. Eight-six per cent of cases in the Casino area occurred in people aged more than five years, most of whom were fully immunised.

Pertussis-related mortality and morbidity is greatest in infants, while in adults symptoms can be limited to a persistent cough.

An accelerated immunisation program has been advised for infants in the Casino area while the outbreak continues. A study is planned to examine the transmission of pertussis in families during the outbreak.

In 1994 the National Health and Medical Research Council has recommended that a pertussis vaccine booster be added to the combined diphtheria and tetanus toxoid (CDT) vaccine given as a preschool booster. The resulting boosted immunity in primary and secondary school age children should result in a decrease in cases in these age groups in future.

TABLE 12

INFECTIOUS DISEASE NOTIFICATIONS FOR 1994 BY PUBLIC HEALTH UNIT, RECEIVED BY NOVEMBER 30, 1994

Condition	CSA	SSA	ESA	SWS	WSA	WEN	NSA	CCA	ILL	HUN	NC	ND	WNS	CW	SW	SE	U/K	Total
Adverse event after immunisation AIDS Arboviral infection Brucellosis Foodborne illness (NOS) Gastroenteritis (instit) Gonorrhoea H. influenzae epiglottitis H. influenzae epiglottitis H. influenzae infection (NOS) Hepatitis A – acute viral Hepatitis B – acute viral Hepatitis B – acute viral Hepatitis B – acute viral Hepatitis B – unspecified Hepatitis D – unspecified Hepatitis C – acute viral Hepatitis C – acute viral Hepatitis C – acute viral Hepatitis C – acute viral Hepatitis B – acute viral Hepatitis B – acute viral Hepatitis C – acute viral Hepatitis C – acute viral Hepatitis C – acute viral Hepatitis S – acute viral Hepatitis S – acute viral Hepatitis S – acute viral Meatific S – acute viral Hepatitis B – acute viral Mentorico HV infection Hydatid disease Leprosy Leptospirosis Listeriosis Malaria Meningococcal septicaemia Meningococcal infection (NOS) Mumps Mycobacterial tuberculosis Mycobacterial infection (NOS) Other venereal Pertussis Q fever Rubella Salmonella (NOS) Salmonella bovis morbificans Salmonella typhimurium Syphilis		2 18 3 1 12 14 16 3 3 - - 1 2 2 14 4 17 - - 1 2 2 4 3 7 - - - - - - - - - - - - -	2 999 3 - - - - - - - - - - - - - - - - -	2 15 - 31 10 2 15 - - 31 10 8 2 4 1 1,096 19 - 7 3 - 11 35 9 7 2 1 5 3 4 4 1 - - - - - - - - - - - - -	6 37 1 15 13 12 13 10 5 14 13 12 13 10 5 16 11 11 12 13 10 12 13 10 10 10 10 10 10 10 10 10 10	$\begin{array}{c} 4 \\ 16 \\ - \\ 8 \\ 277 \\ 1 \\ 3 \\ - \\ 1 \\ 3 \\ - \\ 1 \\ 3 \\ - \\ 7 \\ 6 \\ - \\ 7 \\ 7 \\ 6 \\ 1 \\ 1 \\ 2 \\ 2 \\ - \\ - \\ 4 \\ 4 \\ 37 \\ 2 \\ - \\ - \\ 17 \\ 6 \\ 1 \\ 2 \\ 2 \\ 2 \\ 3 \\ 14 \\ 7 \\ - \\ 1 \\ 2 \\ 2 \\ 2 \\ 1 \\ 2 \\ 2 \\ 1 \\ 1 \\ 2 \\ 2$	- 23 - 23 - 23 - 11 - 5 - 5 - 1 - 23 - 2 - 2 - 2 - 1 - 13 - 2 - 2 - 2 - 2 - 2 - 1 - 13 - 2 - 2 - 2 - 2 - 2 - 2 - 1 -	$\begin{array}{c} 1 \\ 6 \\ 4 \\ -14 \\ 14 \\ 3 \\ -1 \\ 3 \\ 3 \\ -1 \\ 126 \\ -252 \\ -1 \\ -4 \\ -1 \\ -1 \\ -4 \\ 13 \\ 3 \\ 1 \\ -8 \\ 25 \\ -24 \\ 1 \\ 12 \\ -1 \\ -1$	- 12 - 17 - 2 - 99 - 99 - 395 	- 15 45 - 9 130 - 1 - 23 466 - 17 18 - 31 826 68 - 242 45 - 60 24 - 44 223 9	2 19 195 - 23 10 8 2 2 2 1 4 4 8 33 5 - - 2 2 2 2 1 4 4 5 - - - - - - - - - - - - -	$\begin{array}{c} 1\\ 1\\ 6\\ 58\\ 1\\ 2\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\ -\\$	$\begin{array}{c} - & - \\ - & 26 \\ - & 3 \\ 25 \\ - & - \\ 1 \\ 1 \\ - \\ 5 \\ 6 \\ 6 \\ 10 \\ 6 \\ 45 \\ - \\ - \\ 1 \\ - \\ 1 \\ - \\ 1 \\ - \\ 31 \\ 1 \\ 2 \\ - \\ 32 \\ 4 \\ 4 \\ 25 \\ 55 \\ 105 \\ - \\ 9 \\ 103 \\ - \\ 1 \\ 0 \\ 0$	$\begin{array}{c} -1 \\ -1 \\ 4 \\ -7 \\ 30 \\ 5 \\ -2 \\ -33 \\ 10 \\ 7 \\ -33 \\ 10 \\ 7 \\ -12 \\ -1 \\ 12 \\ -35 \\ 4 \\ 11 \\ -1 \\ 2 \\ -1 \\ 18 \\ 18 \\ -11 \\ -10 \\ 13 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ 10 \\ -1 \\ -1$	2 1 1 10 - 2 1 1 - - 2 2 8 9 9 - - 2 2 1 - - - - - - - 2 - - - - - - - - - - - - -	7 7 2 4 4 2 4 4 2 4 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 2 2 2 2 2 2 2 2 2 2 2 2	93	29 323 367 3 3 149 2275 211 15 5 275 21 15 5 3 9 9 453 69 453 69 453 7 7,468 5 3 399 15 5 5 5 5 3 3 12 2 7 7 7,467 1,088 8 9 10 7 7 7 10 7 7 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
i j priora aria paracj priora					-													

TABLE 13

SELECTED INFECTIOUS DISEASE NOTIFICATIONS FOR 1994 BY PUBLIC HEALTH UNIT, RECEIVED BY NOVEMBER 30, 1994

	Sale							Sulla Sin									1 Maria	
Condition	CSA	SSA	ESA	SWS	WSA	WEN	NSA	CCA	ILL	HUN	NC	ND	WNS	CW	SW	SE	U/K	Total
Adverse event after immunisation H. influenzae epiglottitis H. influenzae infection (NOS) H. influenzae meningitis H. influenzae septicaemia Measles Mumps Pertussis Rubella Tetanus	- 2 - 1 - 34 - 25 13 -	2 3 - 19 79 3 -	2 1 - 14 158 14 -	2 2 4 1 35 1 99 -	6 1 2 1 40 - 115 9 -	4 3 - 1 37 39 2 -	- 2 1 2 2 31 3 69 5 -	1 3 - 1 13 - 24 1 -	- 2 1 1 - 49 1 71 - -	- - 1 62 2 60 -	2 2 1 2 239 1 463 6 1	1 - - 156 - 34 6 -	- - 1 31 - 24 10 -	- - 2 50 - 18 - -	2 - 2 - 5 1 9 3 -	7 - 1 273 29 - 2	1111111111	29 21 9 15 11 1,088 10 1,216 72 3

FOODBORNE INFECTIOUS DISEASE NOTIFICATIONS FOR 1994 BY PUBLIC HEALTH UNIT, RECEIVED BY NOVEMBER 30, 1994																		
Condition	CSA	SSA	ESA	SWS	WSA	WEN	NSA	CCA	ILL	HUN	NC	ND	WNS	CW	SW	SE	U/K	Total
Foodborne illness (NOS)	3	12	10	31	15	8	5	14	1	9	23	2	3	7	2	4	-	149
Hepatitis A – acute viral	25	21	45	42	25	6	30	3	7	23	46	50	5	33	89	3	-	453
Salmonella (NOS)	27	41	33	49	51	26	65	23	16	44	72	63	25	11	29	13	-	588
Salmonella bovis morbificans Salmonella typhimurium	23	2 25	1 21	1	1 51	3 14	2 36	17	1 20	223	14	12	9	10	25	2	2	13 313
Typhoid and paratyphoid	5	2	3	3	3	1	1	-	-	-	2	3	-	-	-	2	-	25