

## Supplementary Material

### Suppression of Sertoli cell tumour development during the first wave of spermatogenesis in inhibin $\alpha$ -deficient mice

*Jenna T. Haverfield<sup>A,B</sup>, Peter G. Stanton<sup>A,C</sup>, Kate L. Loveland<sup>A,B,C</sup>, Heba Zahid<sup>A,D</sup>, Peter K. Nicholls<sup>A,C</sup>, Justine S. Olcorn<sup>A,C</sup>, Yogeshwar Makanji<sup>A</sup>, Catherine M. Itman<sup>E</sup>, Evan R. Simpson<sup>A,C</sup> and Sarah J. Meachem<sup>A,B,F</sup>*

<sup>A</sup>Hudson Institute of Medical Research, 27–31 Wright Street, Clayton, Vic. 3168, Australia.


<sup>B</sup>Department of Anatomy and Developmental Biology, Monash University, Wellington Road and Blackburn Road, Clayton, Vic. 3800, Australia.

<sup>C</sup>Department of Biochemistry and Molecular Biology, Monash University, Wellington Road and Blackburn Road, Clayton, Vic. 3800, Australia.

<sup>D</sup>Faculty of Applied Medical Science, Taibah University, Al Madinah Al Monawarah, Universities Road, Medina, 30001, Saudi Arabia.

<sup>E</sup>Priority Research Centres for Reproductive Science and Chemical Biology, School of Environmental and Life Sciences, Faculty of Science and Information Technology, University of Newcastle, University Drive, Callaghan, NSW 2308, Australia.

<sup>F</sup>Corresponding author. Email: sarah.meachem@princehenrys.org



**Fig. S1.** Effect of acute FSH suppression during week 1, 2 or 4 at day 35 on relative mRNA expression of *Star*, *Hsd17b1*, *Igfbp3* and *Smad3*. Data are expressed as mean  $\pm$  s.d. Statistically significant differences across groups are denoted by different letters.