

### Characteristics of gay, bisexual and other men who have sex with men testing and retesting at Australia's first shop-front rapid point-of-care HIV testing service

*Kathleen Ryan<sup>A,B,E</sup>, Anna Wilkinson<sup>A,B</sup>, David Leitinger<sup>A</sup>, Carol El-Hayek<sup>A</sup>, Claire Ryan<sup>A</sup>, Alisa Pedrana<sup>A,B,C</sup>, Margaret Hellard<sup>A,B,D</sup> and Mark Stoové<sup>A,B</sup>*

<sup>A</sup>Centre for Population Health, Burnet Institute, 85 Commercial Road, Melbourne, Vic. 3004, Australia.

<sup>B</sup>School of Public Health and Preventive Medicine, Monash University, Alfred Hospital, Commercial Road, Melbourne, Vic. 3004, Australia.

<sup>C</sup>Department of Nutrition, Harvard T.H. Chan School of Public Health, Huntington Avenue, Cambridge, MA, 02115, USA.

<sup>D</sup>Alfred Health, Infectious Disease Department, Alfred Hospital, Commercial Road, Melbourne, Vic. 3004, Australia.

<sup>E</sup>Corresponding author. Email: kathleen.ryan@burnet.edu.au

#### Supplementary Table 1 – Descriptive characteristics of clients who did and did not return within six months of first test

	No return		Return within 6 months	
	n	%	n	%
	841	76.8	254	23.2
<b>Demographics</b>				
Age group				
16-29	401	47.7	119	46.9
30-39	254	30.2	85	33.5
40+	186	22.1	50	19.7
Median	30.6 (26.1-38.6)		30.5 (25.7-37.1)	
Australian born	526	62.6	134	53.2
Aboriginal &/or Torres Strait Islander	37	4.5	8	3.2
<b>Reason to test</b>				
Regular check up	558	66.4	180	70.9
Recent risk exposure	225	26.8	69	27.2
Follow up for window period	43	5.1	15	5.9

Other	97	11.5	23	9.1
<b>Sexual risk</b>				
Number of anal sex partners				
1 - 10 men	740	88.0	221	87.1
11+ men	101	12.0	33	13.0
Condom use with casual partner(s)				
Consistent	406	48.3	116	45.7
Inconsistent	327	38.9	113	44.5
No casual anal sex/missing	108	12.8	25	9.8
Condom use with regular partner(s)				
Consistent	185	22.0	52	20.5
Inconsistent	338	40.2	103	40.6
No regular anal sex/missing	318	37.8	99	39.0
Group sex with two or more men				
Yes	282	33.7	95	37.6
Recreational drug use before or during sex				
Yes	117	14.0	44	17.5
<b>Testing history</b>				
Never tested	133	16.1	35	14.1
<= 6 months ago	251	30.4	62	49.0
> 6 months ago	330	40.0	122	24.9
Tested, but no details provided	112	13.6	30	12.1

Restricted to men reporting at least one anal sex partner in the six months before the test  
a- Other includes HIV symptoms, STI symptoms, other

**Supplementary Table 2. Descriptive characteristics of clients at first test at PRONTO! reporting no or yes previous HIV testing**

	Previously tested		Never tested	
	n	%	n	%
	973	82.4	208	17.6
<b>Demographics</b>				
Age group				
16-29	423	43.4	139	66.8
30-39	336	34.5	29	13.9
40+	214	22.0	40	19.2
Median (IQR)	31.0 (26.8-38.7)		26.3 (22.8-34.4)	
Australian born	591	60.7	121	58.2
Aboriginal &/or Torres Strait Islander	44	4.5	8	3.9
<b>Reason to test</b>				
Regular check up	673	69.2	107	51.4
Recent risk exposure	249	25.6	71.0	34.13
Other <sup>a</sup>	106	10.9	27	13.0
<b>Sexual risk</b>				
Number of anal sex partners				
None/missing	66	6.8	40	19.2
1-10men	783	80.5	158	76.0
11+ men	124	12.7	10	4.8
Condom use with casual partner				
Consistent	427	43.9	86	41.4
Inconsistent	374	38.4	59	28.4
No casual anal sex/missing	172	17.7	63	30.3
Condom use with regular partner				
Consistent	202	20.8	33	15.9
Inconsistent	380	39.1	55	26.4
No regular anal sex/missing	391	40.2	120	57.7
Group sex with two or more men				
Yes	349	35.9	28	13.5
Recreational drug use before or during sex				
Yes	141	14.5	21	10.1
<b>Return to PRONTO!</b>				
Didn't return in 6m	752	77.3	169	81.3
Return in 6m	221	22.7	39	18.8

24 GBM did not provide testing history details and were excluded

a- Other includes HIV symptoms, STI symptoms, other