

Accessory publication**Fuel loading and flammability in the Mediterranean Basin woody species with different post-fire regenerative strategies**

S. Saura-Mas^{A,D}, S. Paula^B, J. G. Pausas^C and F. Lloret^A

^ACREAF (Center for Ecological Research and Forestry Applications) and Unit of Ecology, Department of Animal and Plant Biology and Ecology, Autonomous University of Barcelona, E-08193 Bellaterra, Barcelona, Spain.

^BCEAM (Fundación Centro de Estudios Ambientales del Mediterráneo), C/ Charles R. Darwin 14, Parc Tecnologic, E-46980 Paterna, Valencia, Spain.

^CCIDE, CSIC, Camí de la Marjal s/n, Albal, E-46470 Valencia, Spain.

^DCorresponding author. Email: s.sauramas@creaf.uab.es

Table A1. Mean value per species and site (standard error in parentheses) of the live and dead fuel relative to the total plant dry mass, separated by fuel diameter and overall mean value

Leaves are included in the live fine-fuel fraction

Seeder	Fine (< 6mm)				Coarse (> 6 mm)				Total			
	Live		Dead		Live		Dead		Live		Dead	
	Montgri	Murta	Montgri	Murta	Montgri	Murta	Montgri	Murta	Montgri	Murta	Montgri	Murta
<i>Calicotome spinosa</i>	0.36 (0.05)	0.24 (0.01)	0.29 (0.04)	0.24 (0.02)	0.11 (0.08)	0.51 (0.01)	0.39 (0.07)	0.01 (0.003)	0.61 (0.07)	0.75 (0.02)	0.39 (0.07)	0.25 (0.02)
<i>Cistus albidus</i>	0.44 (0.07)	0.46 (0.05)	0.18 (0.06)	0.18 (0.02)	0.31 (0.05)	0.35 (0.04)	0.07 (0.04)	0.02 (0.02)	0.75 (0.10)	0.80 (0.02)	0.25 (0.10)	0.20 (0.02)
<i>Cistus monspeliensis</i>	0.56 (0.11)	0.41 (0.04)	0.17 (0.06)	0.12 (0.02)	0.20 (0.05)	0.46 (0.03)	0.07 (0.07)	0.01 (0.01)	0.76 (0.12)	0.87 (0.03)	0.24 (0.12)	0.13 (0.03)
<i>Dorycnium hirsutum</i>	0.88 (0.03)		0.12 (0.03)		0		0		0.88 (0.03)		0.12 (0.03)	
<i>Dorycnium pentaphyllum</i>	0.74 (0.03)		0.25 (0.04)		0.01 (0.01)		0		0.75 (0.04)		0.25 (0.04)	
<i>Fumana ericoides</i>	0.88 (0.04)	0.89 (0.02)	0.11 (0.03)	0.11 (0.02)	0.01 (0.01)	0	0	0	0.89 (0.03)	0.89 (0.02)	0.11 (0.03)	0.11 (0.02)
<i>Fumana laevisipes</i>	0.97 (0.01)		0.03 (0.01)		0		0		0.97 (0.01)		0.03 (0.01)	
<i>Fumana thymifolia</i>	0.95 (0.03)	0.77 (0.02)	0.01 (0.01)	0.23 (0.02)	0.03 (0.03)	0	0	0	0.99 (0.01)	0.77 (0.02)	0.01 (0.01)	0.23 (0.02)
<i>Globularia alypum</i>	0.57 (0.09)	0.69 (0.02)	0.22 (0.05)	0.15 (0.01)	0.18 (0.03)	0.17 (0.03)	0.03 (0.02)	0	0.75 (0.07)	0.85 (0.01)	0.25 (0.07)	0.15 (0.01)
<i>Helichrysum stoechas</i>	0.72 (0.06)		0.19 (0.04)		0.08 (0.03)		0.01 (0.01)		0.81 (0.05)		0.19 (0.05)	
<i>Rosmarinus officinalis</i>	0.48 (0.06)	0.52 (0.03)	0.12 (0.03)	0.05 (0.01)	0.38 (0.03)	0.44 (0.03)	0.01 (0.01)	0	0.86 (0.03)	0.95 (0.01)	0.14 (0.03)	0.05 (0.01)
<i>Teucrium polium</i>	0.80 (0.05)	0.66 (0.05)	0.16 (0.04)	0.34 (0.05)	0.05 (0.02)	0	0	0	0.84 (0.04)	0.66 (0.05)	0.16 (0.04)	0.34 (0.05)
<i>Thymus vulgaris</i>	0.83 (0.03)	0.80 (0.04)	0.07 (0.04)	0.18 (0.03)	0.10 (0.02)	0.02 (0.02)	0	0	0.93 (0.04)	0.82 (0.03)	0.07 (0.04)	0.18 (0.03)
Non-seeder												
<i>Coronilla minima</i>	0.79 (0.04)	0.73 (0.03)	0.17 (0.03)	0.26 (0.02)	0.04 (0.02)	0.01 (0.01)	0	0	0.83 (0.03)	0.74 (0.02)	0.17 (0.03)	0.26 (0.02)
<i>Daphne gnidium</i>	0.55 (0.09)	0.30 (0.07)	0.07 (0.03)	0.10 (0.02)	0.36 (0.06)	0.51 (0.04)	0.02 (0.01)	0.08 (0.04)	0.91 (0.04)	0.81 (0.06)	0.09 (0.04)	0.19 (0.06)
<i>Osyris alba</i>	0.68 (0.06)	0.49 (0.01)	0.24 (0.05)	0.06 (0.01)	0.07 (0.05)	0.45 (0.01)	0	0	0.76 (0.05)	0.94 (0.01)	0.24 (0.05)	0.06 (0.01)
<i>Phillyrea angustifolia</i>	0.39 (0.03)	0.47 (0.03)	0.03 (0.01)	0.09 (0.01)	0.57 (0.02)	0.44 (0.02)	0.01 (0.003)	0.003 (0.002)	0.96 (0.01)	0.90 (0.01)	0.04 (0.01)	0.10 (0.01)
<i>Pistacia lentiscus</i>	0.39 (0.06)	0.47 (0.01)	0.07 (0.01)	0.09 (0.004)	0.50 (0.05)	0.42 (0.02)	0.04 (0.02)	0.01 (0.003)	0.89 (0.03)	0.89 (0.01)	0.11 (0.03)	0.11 (0.01)
<i>Quercus coccifera</i>	0.50 (0.01)		0.11 (0.01)		0.39 (0.01)		0		0.89 (0.01)		0.11 (0.01)	
<i>Rhamnus alaternus</i>	0.36 (0.05)	0.43 (0.03)	0.07 (0.02)	0.12 (0.01)	0.55 (0.04)	0.45 (0.02)	0.01 (0.01)	0.005 (0.003)	0.91 (0.03)	0.87 (0.02)	0.09 (0.03)	0.13 (0.02)
<i>Rhamnus lycioides</i>		0.35 (0.03)		0.18 (0.02)				0.06 (0.01)		0.76 (0.03)		0.24 (0.03)